

r_Patient_Undergoing_Cancer_Therapy_A_Systematic_Review__1_2.pdf

by

Submission date: 27-Nov-2021 07:26PM (UTC+0700)

Submission ID: 1713589087

File name: r_Patient_Undergoing_Cancer_Therapy_A_Systematic_Review__1_2.pdf (283.78K)

Word count: 6595

Character count: 35287

Effectiveness of Oral Cryotherapy for Oral Mucositis on Cancer Patient Undergoing Cancer Therapy: A Systematic Review

ABSTRACT

Background: Oral mucositis is one of the complications in patients undergoing cancer therapy. Oral cryotherapy is an intervention to reduce the severity of oral mucositis.

Aim: To identify, analyze and evaluate articles regarding the effectiveness of cryotherapy for oral mucositis in cancer patients undergoing cancer therapy.

Methods: We performed searching on seven databases (PubMed, CENTRAL, Wiley, CANCELIT, Science Direct, EBSCO, and SpringerLink). The investigation focused on English-language articles, intervention or observational study reporting on the effectiveness of oral cryotherapy against oral mucositis in patients undergoing chemotherapy, and published between October 2015 – October 2020.

Results: Eleven articles met the eligibility criteria for inclusion consisting of 5 RCTs, 3 Quasi-Experiment studies, 2 Cohort studies and one pre-experimental study. The majority of the studies show that oral cryotherapy is an effective intervention to reduce the degree and severity of oral mucositis in patients undergoing cancer therapy (Hematopoietic Stem Cell Transplantation and chemotherapy).

Conclusion: Oral cryotherapy is practical, low-cost, and relatively safe, and can be an alternative therapy in reducing oral mucositis as a side effect of cancer therapy.

Keywords: *Cryotherapy, oral mucositis, cooling therapy, cancer, cancer therapy*

Introduction

Cancer is projected as the top rank among the majority of diseases that have caused the highest incidence of death in the world. Global Burden Cancer (GLOBOCAN) notes that the million cases in 2018 [1]. In the United States, an estimated 1.8 million new cancer cases, with a death rate of more than 600 thousand cases by 2020 [2]. The prevalence of the cases demands the implementation of cancer therapy which can prolong survival and improve patient quality of life.

Currently, combined cancer therapies for example surgery, chemotherapy and radiotherapy have shown promising results on patients survival, however, the cancer therapy can also cause side effects which affect the patients quality of life [3]. One of the most common side effects of cancer therapy is oral mucositis [4,5], which were experienced by 20-80% of patients receiving chemotherapy, 73.2% of patients undergoing Hematopoietic Stem Cell Transplantation (HSCT) [6], and by almost all patients that receive radiation therapy in the head and neck area [7].

Oral mucositis is characterized by inflammation and damage to the mucous membrane of the mouth [8]. Oral mucositis conditions can cause pain in the mouth, decreased/changed sense of taste, dry mouth, decreased saliva production, difficulty swallowing, decreased food intake, and secondary infections [9,10]. This condition can have a negative physical and psychological impact on the quality of life of cancer patients [11]. Thus, prevention and treatment of oral mucositis as a result of cancer therapy requires appropriate intervention.

Complementary and alternative medicine (CAM) therapies have been used by patients to treat various symptoms [12]. One of them is oral cryotherapy for managing oral mucositis is cryotherapy [13,14]. Oral Cryotherapy (OC) is the application of ice chips before, during and after therapy which is useful in preventing and relieving oral mucositis [15,16]. This intervention is easy to be applied and cause no severe side effects [14].

There are several studies that have been conducted to evaluate cryotherapy as treatment and prevention risk of oral mucositis, more focused on kind of cancer treatment [17–19]. However, there is limited studies that discussed tools to determine level of mucositis, how to apply this intervention, and effectiveness of cryotherapy its not only for oral mucositis, but also for another side effect of cancer therapy. Therefore, this study aimed systematically identify, analyze and evaluate articles regarding the effectiveness of OC to oral mucositis in cancer patients undergoing cancer therapy, focusing on assessment of mucositis, procedure, and effectiveness of an intervention.

Methods

This study is a systematic review. We refer to the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) checklist-guideline [20], in assessing and evaluating the studies in this review. We can freely be assessed in <http://www.prisma-statement.org/>.

Search Strategy

Literature searches were conducted from 15 September to 23 October 2020, on seven databases: PubMed, CENTRAL, Wiley, CANCERLIT, Science Direct, EBSCO, and SpringerLink. PICO (patient, intervention, comparison and outcome) framework was used to

facilitate the search strategy, for instance, P: cancer OR neoplasms, I: cryotherapy OR cooling therapy, C: No comparison, O: oral mucositis. PICO and the usage of the keywords depend on the database, which was combined with the use of Boolean "AND" and "OR" (Table 1). In addition, we also searched for articles outside the databases as a secondary search, namely on Google Scholar.

⁶ Eligibility Criteria

Studies were included in this review if they met the following criteria: 1) Article¹⁴ published in English; 2) Intervention or observational studies; 3) Articles reporting on the effectiveness of cryotherapy on oral mucositis experienced by cancer patients undergoing cancer therapy; 4) Articles published from October 2015 – October 2020. Review articles, editorial reviews, qualitative studies, abstracts, and conference proceedings were excluded.

Study Selection

⁵¹ Titles and abstracts were screened based on predetermined inclusion and exclusion criteria. Furthermore, we held panel discussion until an agreement was reached in determining which articles were suitable for inclusion in this study. There were 1687 articles identified in the database, and ten articles on secondary searches outside the databases. Subsequently, papers that had duplicates were excluded, so that 1564 articles were obtained. Subsequent screening⁶¹ eliminated 1475 items and finally only 89 articles remained for assessment (5.69%). We assessed the full-text articles to determine the eligibility of the studies. After the⁴¹ consensus of the review team, we excluded 78 articles and only 11 studies meet eligibility criteria for this review (Table 1). The study inclusion process is illustrated in a PRISMA flow chart (Figure 1).

Table 1. Search Strategy of Electronic Databases


Database	Search Terms	Result	Search Screening	Screening Result	Include Studies
PubMed	(((((cancer[MeSH Terms]) OR (neoplasms[MeSH Terms]) AND (cryotherapy[MeSH Terms]) OR (cryotherapies[MeSH Terms]) OR (cold therapy[MeSH Terms]) OR (cold therapies[MeSH Terms])) AND (oral mucositis[MeSH Terms]))	88	5 years: 26 Full text: 25	25	4
CENTRAL	cancer in Title Abstract Keyword OR neoplasms in Title Abstract Keyword AND cryotherapy in Title Abstract Keyword OR cooling therapy in Title Abstract Keyword AND oral mucositis in Title Abstract Keyword - -	156442	2016-2020: 58474 1000 articles most relevant: 22	22	1
Wiley	"cancer" in Abstract and "chemotherapy OR radiotherapy" in Abstract and "oral cryotherapy" in Abstract and "oral mucositis" in Abstract	13	Last 5 Years: 3	3	0
CANCERLIT	cancer OR neoplasm AND cryotherapy OR cooling therapy AND oral mucositis	102	Last 5 years:50 English: 49	49	1
Science Direct	cancer OR neoplasms AND cryotherapy OR cryotherapies OR cooling therapy OR cooling therapies AND oral mucositis title: mucositis	8618	2016-2020: 1721 Research article: 828	828	0
EBSCO	cancer OR neoplasm AND (cryotherapy or ice therapy or cold therapy) AND oral mucositis	6235	5 years: 946 Academic Journals: 639	639	0
SpringerLink	cancer OR neoplasms AND cryotherapy OR cooling therapy AND oral mucositis!	571	2016-2020:221 Articles:111	111	0
Secondary Search outside of the database				10	5
Total include studies					11

Identification

Screening 10

Eligibility

Inclusion



32
Figure 1. PRISMA Flow Chart

Evaluation of quality of articles

58

We used the Critical Appraisal Skills Program (CASP) tools [21,22], to assess the feasibility of the included studies (Table 2).

Data Extraction

Data extraction process was carried out and developed in the form of a synthesis grid (table 3). The data includes researcher, year, study design, objective, type of cancer, mucositis assessment, outcome measurement, OC procedures, and research results. The accuracy of data extraction is ensured by checking the data 3 times by reviewer panels.

Table 2 Critical Appraisal

No.	Appraisal Checklist Experimental Studies	Johanssen et al. (2019)	Lu et al. (2019)	Nawi et al. (2018)	Askarifar et al. (2016)	Soliman (2019)	Silaban et al. (2020)	Parajuli et al. (2016)	Turkeli et al. (2017)	Bai (2019)	No	Appraisal Checklist Cohort Studies	Okamoto et al. (2019)	Chen et al. (2015)
1	Focused issue	Y	Y	Y	Y	Y	Y	Y	Y	Y	1	Focused issue	Y	Y
33	Randomization	Y	Y	Y	Y	Y	Y	Y	N	N	2	Recruited in an acceptable way	Y	Y
3	All of the patients properly accounted for at its conclusion	Y	Y	Y	Y	Y	Y	?	?	N	3	Exposure accurately measured	Y	Y
4	Blind	N	?	N	Y	?	?	?	?	N	4	Outcome accurately measured	Y	Y
5	The groups similar at the start of the trial	Y	Y	Y	Y	Y	Y	Y	Y	N	5	Identified all-important confounding factors	?	N
6	the groups treated equally	Y	Y	Y	Y	Y	Y	Y	Y	Y	6	The follow up of subjects complete and long enough	Y	Y
7	Intervention effect accounted	Y	Y	Y	Y	Y	Y	?	Y	N	7	Reported the rate or the proportion between the exposed/unexposed, the ratio/rate difference	?	N
8	The accuracy of the estimated effect of the intervention can be accounted	Y	Y	?	Y	Y	Y	?	?	?	8	How precise are the results	Y	Y
5	the results can be applied to the local population, or in your context	?	?	Y	?	Y	Y	?	?	?	9	Believe the results	Y	Y
10	all clinically important outcomes were considered	Y	Y	Y	Y	Y	Y	?	Y	?	10	Can the results be applied to the local population	Y	Y
48	Are the benefits worth the harms and costs	?	Y	?	Y	?	Y	?	Y	?	11	Fit with other available evidence	Y	Y
11	Level of evidence	1c	1b	2b	1b	2b	2b	2c	2b	2c	12	The implication of nursing practice	Y	46
	Grade of recommendation	A	A	B	A	B	B	B	B	C		Grade of recommendation	B	B

Y: Yes, N: No, ? Can't Tell

Table 3 Synthesis Grid

Researcher; Country	Design study	Aim	Type cancer	Cancer Therapy	Assessment mucositis	Another Outcome measurement	Procedure Oral Cryotherapy	Result
Johansson et al. (2019); Sweden [23]	RCT	To investigate whether 2 hours of cryotherapy is as useful as 7 hours of cryotherapy in treating oral mucositis in patients receiving high doses of Melphalan as auto SCT in myeloma patients	Multiple Myeloma	(56)motherapy (high-dose Melphalan) and autologous stem cell transplantation	WHO and OMAS scale	Infectious haematological toxicity Analgesic needs TPN	Ice cube is given 30 minutes before the drug is given (47) he patient did not take any other oral intake during cryotherapy and were instructed to n(74) about one tablespoon of shaved ice in their mouth and immediately replenished. Duration: 2 hour and 7 hours of cryotherapy The oral mucositis assessment(50) was performed before cryotherapy (day 2), just before stem cell infusion (day0), and every day it was started +4 days until the resolution of the mucositis.	Primary Outcome: No significant difference was observed in reference to the proportion of patients presenting with grade 3 and 4 toxicity according to the WHO Scale (2.1 and 4.3% over 2 and 7 hours, respectively; 95% CI - 0.09 to 0.049; p = 0.98) between the two groups. Secondary Outcome: There is no significant difference between IC and CG regarding haematological, infectious and toxicity need for analgesics and TPN.
Lu et al. (2019); China [21]	RCT	To determine the best time to administer oral cryotherapy to prevent oral mucositis in patients undergoing HSCT	Hematology Cancer	HSCT	NCI-CTC version 2 OMDQ	Duration and recovery time for severe oral mucositis	Patients were given an ice cube with (46)ound angle, measuring (3.2 cm x 3.3 cm x 1 cm), comparing the administration of ice cube to 3 groups who received oral cryotherapy intervention, some were given early initiation of therapy, in the middle, and administration of ice cube	Primary Outcome: Both groups of patients who were given OC before and at the time of receiving HSCT had a low incidence of mucositis and a short duration of (19)cositis (≥ level 3), although there was no significant difference. The highest incidence of mucositis was in the group given OC after the HSCT procedure.

minutes 2 times a day. **Secondary Outcome:** OC administration can reduce the duration and recovery time of severe mucositis

Patients can have a pause of 60-180 seconds if they feel uncomfortable. Mucositis assessments were carried out daily for up to 15 days after the completion of HSCT

71

Okamoto et al. (2019); Japan [25]

Retrospective cohort study
Evaluating the role of oral cryotherapy as a prophylaxis in patients receiving DCF therapy

Esophageal Cancer

Chemotherapy (DCF)

Grading Oral mucositis, but it was not mentioned what kind of assessment

Anorexia level

Ice cube is given to patients 10 minutes before DCF is given until the completion of therapy. The total water content in the ice provided is 200-300 ml. The chunks of ice are round on each side.

Primary Outcome: The incidence of oral mucositis in all grades and grade 3 was significantly lower in the OC group than in the non-OC group (24.1% vs 71.4%, $p < 0.001$ and 0% vs. 28.6%, $p = 0.001$ respectively)

Secondary Outcome: The incidence of anorexia in all grades and grade 3 was significantly lower in the OC group (22.4% vs. 57.1%, $p = 0.037$ and 0% vs. 28.6%, $p = 0.010$, respectively)

Nawi, et al. (2018); Malaysia [26]

RCT

Evaluating the effectiveness of oral cryotherapy in the prevention of oral mucositis and pain in colorectal cancer patients undergoing fluorouracil chemotherapy

Colorectal Cancer

Chemotherapy

WHO Mucositis Scale

Pain associated with VAS scale

Giving ice chips for 30 minutes while the chemo drug is given

Primary Outcome: The majority of CG participants exhibited grade 2 mucositis levels and more significant grading compared to IG ($p < 0.001$).

Secondary Outcome: The majority of participants in IG reported no pain

Silaban et al. (2020); Indonesia [27]

Quasi-experiment

To identify the effectiveness of oral cryotherapy in the prevention of oral mucositis in cancer

Solid Cancer

Chemotherapy

OAG

-

There is no specific mention of the oral cryotherapy procedure. However, it is described that the oral mucositis assessment was performed

Primary Outcome: There is a difference in the value of mucositis after being given OC treatment (post) where 29 respondents did not experience mucositis, and three respondents

patients undergoing chemotherapy

at 7 and 14 days after the intervention

experienced mucositis. While in CG, 18 respondents experienced mucositis and did not experience mucositis 12. The degree of mucositis of patients in the two treatment groups was found to be significant⁵⁴

Primary Outcome: There was a decrease in the degree of mucositis of patients who were given OC compared to those who did not get OC

Ice cubes were given to IG for five days, 5 minutes before, two times during chemo, and 5 minutes after chemotherapy. Patients in both treatment groups had Grade I and Grade II⁴⁹ cositis.

WHO Mucositis Scale

Chemotherapy

All-type cancer

10 evaluate the effectiveness of oral cryotherapy in treating oral mucositis by patients undergoing chemotherapy

Quasi-experiment

Parajuli et al. (2016); India [28]

To assess the degree of oral mucositis of patients receiving chemotherapy before and after oral cryotherapy

Pre-experiment study

Bai (2019); India [29]

OAG

Chemotherapy

Solid Cancer

Ice cubes were applied to the oral mucosa for 5 minutes before the chemotherapy session and 20 minutes after the session. The post-test assessment using OAG was carried out on the 7th day.

Pre-experiment study

20

Primary Outcome: There was a significant difference between **72** pre-test and post-test scores of oral mucositis levels among cancer patients receiving chemotherapy.

Secondary Outcome: OC was significantly effective in reducing the degree of oral mucositis in patients receiving chemotherapy.

There is a significant relationship between the degree of oral mucositis after OC in patients receiving chemotherapy with demographic variables such as age, gender, religion, educational status, marital status, and causes of disease

Askarifar et al. (2016); Iran [30]

RCT

1 to determine the effectiveness of oral cryotherapy in patients who receive continuous chemotherapy with oral mucositis while undergoing bone marrow transplantation

Multiple Myeloma, Hodgkin's and Non-Hodgkin's Cancer

Chemotherapy WHO Mucositis Scale

Neutrophil Rate

55 cubes were applied 5 minutes before, during, and after each dose of chemotherapy drug. Initially, ice cubes were used continuously, once, for thirty minutes. Then, with a maximum break of twenty minutes, ice use is restarted for the next thirty minutes and continues up to five minutes after the completion of chemotherapy. However, a hot sensation in the mouth of fewer than twenty minutes is considered a factor in reusing ice. Ice cubes as ice moulds measured approximately three by four centimeters in a sterile container were prepared for the patients. Mucositis assessments were carried out on days 3, 7, 14 and 21 after chemotherapy

34 **Primary Outcome:** OC was more effective than normal saline in reducing the severity of mucositis in patients receiving chemotherapy (1.81 < 2.54 and 0.13 < 0.92, respectively) on days 7 and 14 ($p < 0.164$)
Secondary Outcome: There was no difference in neutrophil values in the two treatment groups.

Soliman (2019); Egypt [31]

RCT

1 To determine the effectiveness of cryotherapy in dealing with the severity of oral mucositis caused by chemotherapy and pain due to oral mucositis in patients receiving combination chemotherapy

Gastrointestinal Cancer

Chemotherapy WHO Mucositis Scale

The pain was assessed with the NRS

39 Ice cubes are given 5 minutes before, during chemo, and 5 minutes after chemotherapy, a total time of 20-25 minutes. If the ice cubes melt, they must be replaced immediately. If the patient is uncomfortable with ice cubes in the mouth, he can

1 **Primary Outcome:** In the majority of patients in the cryotherapy group, oral mucositis was undetectable (Grade 0) on days 7, 14 and 21. The incidence of similar grades 1 and grade 2 oral mucositis in the cryotherapy group was significantly reduced when compared with the control group where $p < 0.001$.

(Fluorouracil and Leucovorin).

rest for no more than 30 minutes. The ice cubes used have round corners. The ice used must be the right size and can be moved in the mouth. Mucositis assessment was carried out on day 7, 14 and 21 days

Secondary Outcome: patients in CG showed a significantly higher level of oral discomfort ($p = 0.001$).

7

Turkeli et al. (2016); Turkey [32]

To determine whether cryotherapy can prevent mucositis caused by 5-FU therapy through oral mucosal smears by looking at cytological changes

Gastrointestinal Cancer

Chemotherapy

Exfoliative cytology of oral mucosal swabs

Grading Oral mucositis but didn't mention kind of assessment

Primary Outcome: The nuclear and cytoplasmic volumes in cells decreased significantly in patients receiving cryotherapy compared to pre-cryotherapy (0.047 and 0.042, respectively). The results of the histomorphometry estimation show that cryotherapy can be used to prevent oral mucosal damage and can reduce the frequency and duration of oral mucositis caused by 5-FU.

7

Secondary Outcome: Grade I and II mucositis decreased significantly in patients receiving cryotherapy compared to patients without cryotherapy ($p = 0.03$)

37

Chen et al. (2015); Canada [33]

to examine the effectiveness of the cryotherapy protocol applied in a hematopoietic stem cell transplant program.

Multiple Myeloma

HSCT

Duration of oral mucositis, length of stay in the hospital, use of narcotics and parenteral nutrition.

WHO mucositis scale

The patient is instructed to hold the ice flakes inside mouth for 5 minutes before high-dose melphalan infusion, for 30 minutes and 30 minutes after completion of the infusion.

Primary Outcome: The incidence and severity of mucositis were found to be significantly lower in the cryotherapy group. 32.71.4% had mucositis in the OC group compared with 95.7% in CG ($p < 0.001$). The median mucositis severity, assessed using the WHO oral toxicity

scale from 0–4 levels, experienced in CG was 2.5 vs. 2 in the OC group ($p = 0.03$).

Secondary Outcome: The duration of oral mucositis and use of parenteral narcotics was also significantly reduced. Duration of hospitalization and use of parenteral nutrition were similar between the two groups.

52

Abbreviations: 5-FU: 5-fluorouracil; DCF: docetaxel; cisplatin, and fluorouracil; CG: Control Group; HSCT: Haematopoietic Stem Cell Transplantation; IG: Intervention Group; KS: Numeric Rating Scale; OAG: Oral Assessment Guide; OC: Oral Cryotherapy; OMAS: Oral Mucositis Assessment Score; OMDQ: Oral Mucositis Daily Questionnaire; RCT: Randomized Control Trial; TPN: Total Parenteral Nutrition; VAS: Visual Analogue Scale; WHO: World Health Organization

Results

Assessment of Oral Mucositis

Eleven studies included in this review used different instruments to assess oral mucositis in cancer patients undergoing cancer therapy. The instruments were the WHO mucositis scale [23,28,30,31,33], the Oral Assessment Guide (OAG) [15,29], the Oral Mucositis Assessment Scale (OMAS) [23], the National Cancer Institute Common Toxicity Criteria (NCI-CTC) [24], and Oral Mucositis Daily Questionnaire (OMDQ) [24,26]. The OMDQ is a self-assessed questionnaire, whereas other tools are used by health practitioners. However, there are two studies that did not specifically mention the instrument used to assess the oral mucositis in their studies.

OC Procedure

Several studies have shown the time to assess the degree of mucositis in cancer patients receiving cancer therapy, but it is still varied. The principle of applying ice cube into the mouth as a core part of the cryotherapy procedure was relatively similar among the reviewed studies. However, there was a slight difference in the time of administration. The majority of studies applied OC procedures before cancer therapy interventions were carried out. Six studies started giving ice cube 5 minutes before cancer therapy was started [28–33]. Another study started 10 minutes [25], and 30 minutes before cancer therapy was given [23].

Beside time of OC application, duration during therapy was also varied. Some studies started OC when the treatment was begun, mid until the treatment is finished with a period ranging from 5-30 minutes, and some even reach 2-7 hours, some are once or twice. Some are continuous until the therapy is completed, followed by a pause if the patient experiences inconveniences. The majority of the research evaluated the degree of mucositis 1–2 weeks after the intervention.

Regarding the size of the ice cube, only four studies that describe the ice cube criteria were used: the OC [24,25,30,31]. In detail, Lu et al. stated that the size of the ice cube was 3.2 cm × 3.3 cm × 1 cm [24]. Another study describes that the ice cube made from ice moulds measuring three by four centimeters in a sterile container [30]. Okamoto et al. further explained the total water content for making ice cube, which was around 200-300 mL for making ice chunks [25]. Meanwhile, Soliman et al. did not explain size details, but the criteria for the ice cube must be suitable and moveable in the mouth [31]. Although the size criteria of the ice cube were different, these studies agreed that the ice cube used were rounded corners to prevent mouth irritation.

The effectiveness of OC to oral mucositis in patients undergoing cancer therapy

The majority of the reviewed studies described that OC intervention was effective in reducing the degree and severity of oral mucositis in patients undergoing cancer therapy, either HSCT or chemotherapy. Based on the mucositis grading assessment instruments, these studies found mucositis grade reduction in the OC group [25–32]. Meanwhile, three studies reported that the incidence of oral mucositis on different grades was significantly lower in the group of patients receiving OC than the non-OC group [25,26,31].

Similar results were also found in patients who underwent HSCT in which the severity of mucositis was significantly lower in the group receiving OC [30,33]. Additional studies conducted by Lu et al. found that patients who were given OC before and while receiving HSCT had a lower incidence of mucositis or a short duration of mucositis (\geq level 3), although there was no significant difference statistically [24].

The effectiveness of OC on other outcomes in patients undergoing cancer therapy

Ten other outcomes were assessed regarding the effectiveness of OC. It was found that the OC decreased pain level of the patients oral mucositis [26,31], shorter the duration and recovery of severe mucositis [24,33], reduced the incidence of anorexia [25], and usage of parenteral narcotics [33]. Furthermore, the exfoliative cytology results on the oral mucosal layer showed a decrease in the nuclear and cytoplasmic volume in cells after the OC [32]. However, there were no significant differences between OC and controlled groups in term of the level of toxicity of infectious hematology, analgesic need, TPN [23], the neutrophil rate [30], and length of hospitalization [33].

Discussion

We conducted a review on relevant studies, to assess the effectiveness of OC for oral mucositis on cancer patient undergoing cancer therapy. This review focuses on the assessment of oral mucositis, the OC procedure, and the effectiveness of the OC. Assessment of the degree of patient mucositis is an essential component in evaluating the effectiveness of the OC intervention. This review found several instruments used in the included studies.

This diversity of instruments is a clinical-based assessment tool that can be used to assess signs of oral mucositis. The OMDQ is a self-reported instruments that can be an alternative clinical assessment if the patient refuses to do an oral examination by health practitioners [34]. A research conducted by Wardill et al. reported that the NCI-CTC is an instrument that is more widely used in assessing oral mucositis in cancer patients, followed by the Radiation Oncology Research Group (RTOG), WHO mucositis scale, OMAS, Oral Mucositis Index (OMI), and OAG [35]. This research does not specify which one is the most effective instrument and more emphasize the importance of using a validated instrument in assessing mucositis. The usage of the tool must follow the principles of validity and must be able to assist in determining the degree of mucositis so that the appropriate treatment can be applied. In addition, in assessing the degree of mucositis, two instruments can be combined, such as assessment instruments from health practitioners and self-reported instruments.

The OC implementation protocol in each of the reviewed studies have a relatively same regarding the application principle, but difference in the procedural time and duration. Previous research, in patients undergoing chemotherapy, Ismael et al. stated that ice cube administration was given 10 minutes before therapy, continued 15 minutes each while treatment was taking place and after the chemotherapy infusion was complete [36]. Meanwhile, related to duration of OC administration, Research from Johanssen et al. proved that the duration of 2 hours or 7 hours of OC there was no difference in results in determining the length of intervention in the treatment of oral mucositis.

The Multinational Association of Supportive Care in Cancer and the International Society of Oral Oncology (MASCC/ISO) also gave recommendations, not much different from previous years guidelines, it is stated that the recommended duration of OC administration is 30 minutes, because of more than 30 minutes can cause a sensation of discomfort [5,37]. MASCC/ISO guidelines reveal that oral cryotherapy can be a modality to prevent oral mucositis in patients undergoing autologous HSCT accompanied by high doses of Melphalan, and in solid cancer patients receiving 5-FU therapy [37]. This means that the effectiveness of the duration of OC administration should take into account the patient's comfort during the procedure.

Regarding the size and volume of the ice cube/ice chip used, we have not found any definite size guidelines regarding its use and its effect on OC. From this review, we get a summary that the use of ice cube should prioritize patient comfort so that the ice cube used has rounded corners, is suitable and can be moved in the mouth.

In this systematic review, we found that OC can reduce the incidence and severity of oral mucositis in cancer patients undergoing cancer therapy. This review is in line with the management guidelines from MASCC / ISO [5], declared that OC is included as one of the palliative management of patients undergoing cancer therapy. OC is easy to apply, low cost, and relatively safe. OC is characterized by the use of an ice cube/ice chip which is readily available [31]. A study by Walladbegi et al. stated that the procedure was consistently lower the temperature in oral mucosa during the administration of cytotoxic drugs that can reduce the release of pro-inflammatory cytokines [38]. Thus, the effect of tissue damage can be minimized [38].

In addition, this review shows that OC is significantly shorter the pain level, duration and recovery of severe mucositis, a lower incidence of anorexia, decreased parenteral narcotic use, and decreased nuclear and cytoplasmic volume in the oral mucosal layer as a result of exfoliative cytology. Meanwhile, the toxicity level of infection hematology, analgesic requirement, neutrophil rate, length of stay and TPN were not significant. Ulceration of mucositis can cause pain [14]. However, through the cooling process in the oral cavity, it can cause a decrease in the release of inflammatory cytokines (IL-6 and TNF- α) [38], and reduce pain levels [39]. The pain decreases; the appetite increases so that the incidence of anorexia can be suppressed. OC can be a promising intervention in dealing with the side effects of cancer therapy.

Implication of Nursing

Nurses can use the OC method as a nursing implementation in reducing oral mucositis in patients undergoing cancer therapy. Besides, nurses can use the grading mucositis assessment as a routine activity in evaluating the oral mucosa of cancer patients.

Conclusion

OC can be an intervention in preventing the incidence and reducing the degree and severity of oral mucositis in patients undergoing cancer therapy, either

HSCT or chemotherapy. OC is practical, low-cost, and relatively safe to use. Oral cryotherapy can be part of the management of oral mucositis by following appropriate procedural guidelines.

Acknowledgment

²¹ This work was supported by The Directorate of Research and Community Service – The Indonesian Ministry of Research, Technology and Higher Education, (Number: 1516/UN4.22/PT.01.03/2020). The authors would like to thanks to Moh. Syafar Sangkala for the suggestion in improving this manuscript.

References

1. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018 : GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018;68(6):394–424. <https://doi.org/10.3322/caac.21492>. PMID: 30207593
2. Siegel RL, Miller KD, Jemal A. Cancer statistics , 2020. *CA Cancer J Clin.* 2020;0(0):1–24. <https://doi.org/10.3322/caac.21590>. PMID: 31912902
3. Gegechkori N, Haines L, Lin JJ. Long-term and latent side effects of specific cancer types. *Med Clin North Am.* 2017;101(6):1053–73. <https://doi.org/10.1016/j.mcna.2017.06.003>. PMID: 28992854
4. Berger K, Schopohl D, Bollig A, Strobach D, Rieger C, Rublee D, et al. Burden of Oral Mucositis: A Systematic Review and Implications for Future Research. *Oncol Res Treat.* 2018;41(6):399–405. <https://doi.org/10.1159/00048708>. PMID: 29734184
5. Elad S, Cheng KKF, Lalla R V., Yarom N, Hong C, Logan RM, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer.* 2020;1–9. <https://doi.org/10.1002/cncr.33100>. PMID: 32786044
6. Chaudhry HM, Bruce AJ, Chb MB, Wolf RC, Rph P, Litzow MR, et al. The incidence and severity of oral mucositis among allogeneic hematopoietic stem cell transplantation patients: a systematic review. *Biol Blood Marrow Transplant.* 2015;22(4):p605–616. <https://doi.org/10.1016/j.bbmt.2015.09.014>. PMID: 26409924
7. Maria OM, Eliopoulos N, Muanza T. Radiation-Induced Oral Mucositis. *Front Oncol.* 2017;7(MAY). <https://doi.org/10.3389/fonc.2017.00089>. PMID: 28589080
8. Gupta A, West HJ. Mucositis (or Stomatitis). *JAMA Oncol.* 2016;2(10):1379. <https://doi.org/10.1001/jamaoncol.2016.2103>. PMID:

27584707

9. Çakmak S, Nural N. Incidence of and risk factors for development of oral mucositis in outpatients undergoing cancer chemotherapy. *Int J Nurs Pract*. 2019;25(1):1–9. <https://doi.org/10.1111/ijn.12710>. PMID: 30461128
10. Mallick S, Benson R, Rath GK. Radiation induced oral mucositis: a review of current literature on prevention and management. *Eur Arch Otorhinolaryngol*. 2016;273(9):2285–93. <https://doi.org/10.1007/s00405-015-3694-6>. PMID: 26116012
11. Jehn P, Stier R, Tavassol F, Dittman J, Zimmerer R, Gellrich N-C, et al. Physical and psychological impairments associated with mucositis after oral cancer treatment and their impact on quality of life. *Oncol Res Treat*. 2019;42(6):342–9. <https://doi.org/10.1159/000499720>. PMID: 30970370
12. Rasheed Z, Al-Rashidy M, Al-Raqibah A, Al-Moqbel A, Al-Bakri A, Al-Harbi A, et al. Experience of cancer patients for the usage of complementary and alternative medicine therapy. *Open Access Maced J Med Sci*. 2020;8:521–7.
13. Daugėlaitė G, Užkuraitytė K, Jagelavičienė E, Filipauskas A. Prevention and treatment of chemotherapy and radiotherapy induced oral mucositis. *Med*. 2019;55(2):25. <https://doi.org/10.3390/medicina55020025>. PMID: 30678228
14. Riley P, Am G, Hv W, Littlewood A, Je C, Mg M. Interventions for preventing oral mucositis in patients with cancer receiving treatment : oral cryotherapy. *Cochrane Database Syst Rev*. 2015;(12):CD011552. <https://doi.org/10.1002/14651858.CD011552.pub2>. PMID: 26695736.
15. Baysal E, Sari D. Effects of cryotherapy in preventing oral mucositis in hematopoietic stem cell transplantation patients: Literature review. *Complement Ther Clin Pract* [Internet]. 2019;35(March):342–7. Available from: <https://doi.org/10.1016/j.ctcp.2019.03.017>. PMID: 31003680
16. Riley P, McCabe MG, Glenny A. Oral cryotherapy for preventing oral mucositis in patients receiving cancer treatment. *JAMA Oncol*. 2016;2(10):1365–6. <https://doi.org/10.1001/jamaoncol.2016.2680>. PMID: 27583808.
17. Maza M, Figueroa R, Laskow B, Juárez A, Alfaro K, Alonzo TA, et al. Effects of Maintenance on Quality of Performance of Cryotherapy Devices for Treatment of Precancerous Cervical Lesions. *J Low Genit Tract Dis*. 2018;22(1):47–51. <https://doi.org/10.1097/LGT.0000000000000359>. PMID: 29271857
18. Heydari A, Sharifi H, Salek R. Effect of Oral Cryotherapy on Combination Chemotherapy-induced Oral Mucositis: A Randomized Clinical Trial. *Middle East J Cancer* [Internet]. 2012;3:55–64. Available from: <http://mej.sums.ac.ir/index.php/mejc/article/view/42>

19. Kakoei S, Ghassemi A, Nakhaee N. Effect of cryotherapy on oral mucositis in patients with head and neck cancers receiving radiotherapy. *Int J Radiat Res* [Internet]. 2013;11(2):117–20. Available from: <http://ijrr.com/article-1-1037-en.html>
20. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann C, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372(n71):1–9. <https://doi.org/10.1136/bmj.n71>. PMID: 33782057
21. CASP. CASP Cohort Study Check List [Internet]. 2018 [cited 2018 Sep 23]. Available from: http://casp-uk.net/wp-content/uploads/2018/01/CASP-Cohort-Study-Checklist_2018.pdf
22. CASP. CASP Randomised Controlled Trial Checklist [Internet]. 2018 [cited 2018 Sep 23]. Available from: <https://casp-uk.net/wp-content/uploads/2018/01/CASP-Randomised-Controlled-Trial-Checklist.pdf>
23. Johansson JE, Bratel J, Hardling M, Heikki L, Mellqvist UH, Hasséus B. Cryotherapy as prophylaxis against oral mucositis after high-dose melphalan and autologous stem cell transplantation for myeloma: a randomised, open-label, phase 3, non-inferiority trial. *Bone Marrow Transplant* [Internet]. 2019;54(9):1482–8. Available from: <http://dx.doi.org/10.1038/s41409-019-0468-6>
24. Lu Y, Zhu X, Ma Q, Wang J, Jiang P, Teng S, et al. Oral cryotherapy for oral mucositis management in patients receiving allogeneic hematopoietic stem cell transplantation: a prospective randomized study. *Support Care Cancer*. 2020;28(4):1747–54. <https://doi.org/10.1007/s00520-019-04966-z>. PMID: 31302767
25. Okamoto K, Ninomiya I, Yamaguchi T, Shira S, Nakanuma S, Tajima H, et al. Oral cryotherapy for prophylaxis of oral mucositis caused by docetaxel, cisplatin, and fluorouracil chemotherapy for esophageal cancer. *Esophagus*. 2019;16(2):207–13. <https://doi.org/10.1007/s10388-018-00655-8>. PMID: 30600487
26. Nawi RIM, Chui PL, Wan Ishak WZ, Hsien Chan CM. Oral cryotherapy: Prevention of oral mucositis and pain among patients with colorectal cancer undergoing chemotherapy. *Clin J Oncol Nurs*. 2018;22(5):555–60.
27. Silaban NY, Nasution SS, Siregar CT. The Influence of oral cryotherapy on prevention from mucositis in cancer patients under chemotherapy in RSUP H . Adam Malik, Medan. *Indian J Public Heal Res Dev*. 2020;11(6):1380–6.
28. Parajuli S, Rani S, Asha T. An experimental study to evaluate the effectiveness of cryotherapy on occurrence of oral mucositis among patients receiving cancer chemotherapy in selected hospitals of new delhi. *Int J Nurs Midwif Res* 2016. 2016;3(4):27–31.
29. Bai RR. Effectiveness of Oral Ice Cube Application on Prevention of Oral

Mucositis among Cancer Patients Who Receive Chemotherapy. *Innov J Nurs Healthc.* 2019;5(3):38–43.

30. Askarifar M, Lakdizaji S, Ramzi M, Rahmani A, Jabbarzadeh F. The effects of oral cryotherapy on chemotherapy-induced oral mucositis in patients undergoing autologous transplantation of blood stem cells : a clinical trial. *Iran Red Crescent Med J.* 2016;18(4):e24775.
31. Soliman HMM. The effect of cryotherapy on chemotherapy induced oral mucositis in Egyptian cancer patients: A randomized controlled trial. *J Nurs Educ Pract.* 2019;9(11):63.
32. Turkeli M, Aldemir MN, Bingol F, Dogan C, Kara A. A morphometric study of the protective effect of cryotherapy on oral mucositis in cancer patients treated with 5-fluorouracil. *Biotech Histochem.* 2016;91(7):465–71. <https://doi.org/10.1080/10520295.2016.1220020>. PMID: 27644112
33. Chen J, Seabrook J, Fulford A, Rajakumar I. Icing oral mucositis : oral cryotherapy in multiple myeloma patients undergoing autologous hematopoietic stem cell transplant. *J Oncol Pharm Pr.* 2015;23(2):116–120.
34. Ágnes J, János S, Katalin G, Judit N, Csongor K, J MI. Patient-reported measurements of oral mucositis in pediatric patients with cancer. *Oral Health Dent Manag.* 2016;15(6):1–10.
35. Wardill HR, Sonis ST, Blijlevens NMA, Sebille YZA Van, Ciorba MA, Loeffen EAH. Prediction of mucositis risk secondary to cancer therapy : a systematic review of current evidence and call to action. *Support Care Cancer.* 2020;28(11):5059–73. <https://doi.org/10.1007/s00520-020-05579-7>. PMID: 32592033
36. Ameen BIH, Alzubaidee AF, Majid S. Comparative evaluation of the efficacy of ice cubes versus sodium bicarbonate mouthwash both as prophylactic measure and as treatment of oral mucositis induced by systemic anticancer therapies. *Tabari Biomed Student Res J.* 2019;1(4):13–7. <https://doi.org/10.18502/tbsrj.v1i4.2244>
37. Correa MEP, Cheng KKF, Chiang K, Kandwal A, Loprinzi CL, Mori T, et al. Systematic review of oral cryotherapy for the management of oral mucositis in cancer patients and clinical practice guidelines. *Support Care Cancer.* 2020;28(5):2449–56. <https://doi.org/10.1007/s00520-019-05217-x>. PMID: 31836937
38. Walladbegi J, Smith SA, Grayson A, Murdoch C, Jontell M, Colley HE. Cooling of the oral mucosa to prevent adverse effects of chemotherapeutic agents: an in vitro study. *J Oral Pathol Med.* 2018;47:477–83. <https://doi.org/10.1111/jop.12696>. PMID: 29469972
39. Oliveira Júnior JO de, Portella Junior CSA, Cohen CP. Inflammatory mediators of neuropathic pain. *Rev Dor São Paulo.* 2016;17(Suppl 1):35–42. <https://doi.org/10.5935/1806-0013.20160045>.



ORIGINALITY REPORT

28%

SIMILARITY INDEX

25%

INTERNET SOURCES

22%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

1	pdfs.semanticscholar.org Internet Source	2%
2	ircmj.neoscriber.org Internet Source	2%
3	discovery.dundee.ac.uk Internet Source	1%
4	www.cybermedlife.eu Internet Source	1%
5	internationaljournalofcaringsciences.org Internet Source	1%
6	link.springer.com Internet Source	1%
7	www.tandfonline.com Internet Source	1%
8	pubmed.ncbi.nlm.nih.gov Internet Source	1%
9	Submitted to The Royal Marsden School of Cancer Nursing and Rehabilitation Student Paper	1%

10	Submitted to Chester College of Higher Education Student Paper	1 %
11	Eli Zuraida, Andi Masyitha Irwan, Elly Lilianty Sjattar. "Self-management education programs for patients with heart failure: a literature review", Central European Journal of Nursing and Midwifery, 2021 Publication	1 %
12	Joey Chen, Jamie Seabrook, Adrienne Fulford, Irina Rajakumar. "Icing oral mucositis: Oral cryotherapy in multiple myeloma patients undergoing autologous hematopoietic stem cell transplant", Journal of Oncology Pharmacy Practice, 2016 Publication	1 %
13	Submitted to St George's Hospital Medical School Student Paper	1 %
14	medical.adrpublications.in Internet Source	1 %
15	www.ncbi.nlm.nih.gov Internet Source	<1 %
16	Submitted to King's College Student Paper	<1 %
17	Submitted to University Of Tasmania Student Paper	<1 %

18	Ana Myriam Lavín - Pérez, Daniel Collado - Mateo, Xián Mayo, Liam Humphreys et al. "High - intensity exercise to improve cardiorespiratory fitness in cancer patients and survivors: A systematic review and meta - analysis", Scandinavian Journal of Medicine & Science in Sports, 2020 Publication	<1 %
19	zenodo.org Internet Source	<1 %
20	bib.mariehaps.vinci.be Internet Source	<1 %
21	smujo.id Internet Source	<1 %
22	Submitted to University of Sydney Student Paper	<1 %
23	Submitted to University of Colorado, Denver Student Paper	<1 %
24	mafiadoc.com Internet Source	<1 %
25	www.jcdr.net Internet Source	<1 %
26	Sapna Oberoi, Gabriele Zamperlini-Netto, Joseph Beyene, Nathaniel S. Treister, Lillian Sung. "Effect of Prophylactic Low Level Laser	<1 %

Therapy on Oral Mucositis: A Systematic Review and Meta-Analysis", PLoS ONE, 2014

Publication

27

Submitted to University of North Carolina -
Wilmington

Student Paper

<1 %

28

commonrepo.um.edu.my

Internet Source

<1 %

29

Submitted to University of Liverpool

Student Paper

<1 %

30

doi.org

Internet Source

<1 %

31

www.researchsquare.com

Internet Source

<1 %

32

Ángel López-González, Marta García-
Quintanilla, Carmen María Guerrero-Agenjo,
Jaime López Tendero et al. "Efficacy of
Cryotherapy in the Prevention of Oral
Mucositis in Adult Patients with
Chemotherapy", International Journal of
Environmental Research and Public Health,
2021

Publication

<1 %

33

Submitted to Cox College of Nursing and
Health Sciences

Student Paper

<1 %

34	Submitted to UT Health Science San Antonio Student Paper	<1 %
35	pericles.pericles-prod.literatumonline.com Internet Source	<1 %
36	repositorio.unb.br Internet Source	<1 %
37	www.science.gov Internet Source	<1 %
38	Submitted to The University of Memphis Student Paper	<1 %
39	Submitted to University of Hong Kong Student Paper	<1 %
40	Submitted to University of Saint Francis Student Paper	<1 %
41	journals.plos.org Internet Source	<1 %
42	www.nice.org.uk Internet Source	<1 %
43	www.scribd.com Internet Source	<1 %
44	Submitted to Kaplan University Student Paper	<1 %
45	journal.uui.ac.id Internet Source	<1 %

46

www.bjorl.org

Internet Source

<1 %

47

www.nature.com

Internet Source

<1 %

48

Bowling, Ann, Ebrahim, Shah. "EBOOK: Handbook of Health Research Methods: Investigation, Measurement and Analysis", EBOOK: Handbook of Health Research Methods: Investigation, Measurement and Analysis, 2005

Publication

<1 %

49

Ebru Baysal, Dilek Sari, Filiz Vural, Seçkin Çağırğan et al. "The Effect of Cryotherapy on the Prevention of Oral Mucositis and on the Oral pH Value in Multiple Myeloma Patients Undergoing Autologous Stem Cell Transplantation", Seminars in Oncology Nursing, 2021

Publication

<1 %

50

Jan-Erik Johansson, John Bratel, Mats Hardling, Lena Heikki, Ulf-Henrik Mellqvist, Bengt Hasséus. "Cryotherapy as prophylaxis against oral mucositis after high-dose melphalan and autologous stem cell transplantation for myeloma: a randomised, open-label, phase 3, non-inferiority trial", Bone Marrow Transplantation, 2019

Publication

<1 %

51

Su Htet Htet Aung, Kate White, Jacqueline Bloomfield. "The Experiences and the Needs of Caregivers of Patients With Head and Neck Cancer", *Cancer Nursing*, 2021

Publication

<1 %

52

[researchprotocols.org](https://www.researchprotocols.org)

Internet Source

<1 %

53

Java Walladbegi, Sarah A. Smith, Amy K. Grayson, Craig Murdoch, Mats Jontell, Helen E. Colley. "Cooling of the oral mucosa to prevent adverse effects of chemotherapeutic agents: An in vitro study", *Journal of Oral Pathology & Medicine*, 2018

Publication

<1 %

54

Joyce Ohana de Lima Martins, Marcela Maria Fontes Borges, Malta Cássia Nóbrega, Anna Clara Aragão Matos Carlos et al. "Risk Factors for Oral Mucositis During Chemotherapy Treatment for Solid Tumors: A Retrospective STROBE-Guided Study", *Research Square Platform LLC*, 2021

Publication

<1 %

55

Submitted to University of South Florida

Student Paper

<1 %

56

reins.tmd.ac.jp

Internet Source

<1 %

www.frontiersin.org

57	Internet Source	<1 %
58	www.internationaljournalofcaringsciences.org Internet Source	<1 %
59	Submitted to Cardiff University Student Paper	<1 %
60	HV Worthington. "Interventions for treating oral mucositis for patients with cancer receiving treatment", Cochrane Database of Systematic Reviews Reviews, 04/19/2004 Publication	<1 %
61	ajod.org Internet Source	<1 %
62	dergipark.org.tr Internet Source	<1 %
63	ir.ymlib.yonsei.ac.kr Internet Source	<1 %
64	jenci.springeropen.com Internet Source	<1 %
65	jnccn.org Internet Source	<1 %
66	nrl.northumbria.ac.uk Internet Source	<1 %
67	onlinelibrary.wiley.com Internet Source	<1 %

- | | | |
|----|--|------|
| 68 | worldwidescience.org
Internet Source | <1 % |
| 69 | www.mdpi.com
Internet Source | <1 % |
| 70 | www.sciedupress.com
Internet Source | <1 % |
| 71 | Li Wang, Zhenyang Gu, Ruiren Zhai, Shasha Zhao et al. "Efficacy of Oral Cryotherapy on Oral Mucositis Prevention in Patients with Hematological Malignancies Undergoing Hematopoietic Stem Cell Transplantation: A Meta-Analysis of Randomized Controlled Trials", PLOS ONE, 2015
Publication | <1 % |
| 72 | Paras Ahmad, Usman Akhtar, Ahmed Chaudhry, Usman Rashid, Sarmad Saif, Jawaad Ahmed Asif. "Treatment and prevention of oral mucositis: A literature review", European Journal of General Dentistry, 2021
Publication | <1 % |
| 73 | Wanessa Miranda-Silva, Wagner Gomes-Silva, Yehuda Zadik, Noam Yarom et al. "MASCC/ISOO clinical practice guidelines for the management of mucositis: sub-analysis of current interventions for the management of | <1 % |

oral mucositis in pediatric cancer patients",
Supportive Care in Cancer, 2020

Publication

74

YK Cho, DW Sborov, M Lamprecht, J Li et al.
"Associations of High-Dose Melphalan
Pharmacokinetics and Outcomes in the
Setting of a Randomized Cryotherapy Trial",
Clinical Pharmacology & Therapeutics, 2017

Publication

<1 %

75

livrepository.liverpool.ac.uk

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On