

# 8.\_9621-Article\_Text-56033-2- 10-20220115.pdf

*by*

---

**Submission date:** 23-Sep-2022 09:44PM (UTC+0700)

**Submission ID:** 1907121534

**File name:** 8.\_9621-Article\_Text-56033-2-10-20220115.pdf (557.83K)

**Word count:** 7370

**Character count:** 36976

## Evaluation of the advanced yield trial on tropical wheat (*Triticum aestivum*) mutant lines using selection index and multivariate analysis

MUH FADLI<sup>1</sup>, MUH FARID<sup>1\*</sup>, AMIR YASSI<sup>2</sup>, NASARUDDIN<sup>2</sup>, MUHAMMAD FUAD ANSHORI<sup>2</sup>,  
AMIN NUR<sup>3</sup>, SURATMAN<sup>4</sup>

<sup>1</sup>Program of Agrotechnology, Graduate School, Universitas Hasanuddin, Jl. Perintis Kemerdekaan Km. 10, Makassar 90245, South Sulawesi, Indonesia

<sup>2</sup>Department of Agronomy, Faculty of Agriculture, Universitas Hasanuddin, Jl. Perintis Kemerdekaan Km. 10, Makassar 90245, South Sulawesi, Indonesia. Tel.: +62-411-587064, \*email: farid\_deni@yahoo.co.id

<sup>3</sup>Assessment Institute for Agriculture Technology of Gorontalo, Jl. Mohamad Van Gobel No. 270, Bone Bolango 96119, Gorontalo, Indonesia

<sup>4</sup>Indonesia Centre for Agricultural Land Resources Research and Development, Ministry of Agriculture, Jl. Ir. H. Juanda No. 98, Bogor 16123, West Java, Indonesia

Manuscript received: 15 October 2021. Revision accepted: 30 December 2021.

**Abstract.** Fadli M, Farid M, Yassi A, Nasaruddin, Anshori MF, Nur A, Suratman. 2021. Evaluation of the advanced yield trial on tropical wheat (*Triticum aestivum*) mutant lines using selection index and multivariate analysis. *Biodiversitas* 23: 540-547. The development of adaptive wheat (*Triticum aestivum* L.) mutant lines in tropical climate is a long-term solution. The advanced yield trial is an important stage in this development, which is carried out using a selection index and multivariate analysis. This study aims to determine the selection effectiveness of tropical wheat mutant lines by using a selection index and multivariate analysis. The study was conducted in three locations, which included two locations for the follow-up yield test in Kelara Village and Ujung Bulu Village (South Sulawesi, Indonesia), and the third location for the selection validation in Padang Magek Village (West Sumatra, Indonesia). This study was designed by a nested Randomized Complete Block Design (RCBD), with replications nested in locations. The samples included 16 wheat genotypes (13 mutant lines and three comparison varieties) that were replicated three times. The result showed that the selection of tropical wheat mutant lines using selection index and multivariate analysis were considered effectively with high determination value of 0.57 (57%). The results of the selection index show that seven lines were recommended for multi-location trials since they had a better selection index potential than Dewata variety.

**Keywords:** Mutant lines, PCA analysis, selection index, *Triticum aestivum*, tropical climate

**Abbreviations:** 1000-GW: 1000 Grain weight, Chl<sub>a</sub>: Chlorophyll density, CV: Coefficient of variation, DF: Days to flowering, DH: Days to harvest, GFT: Grain Filling time, GWP: Grain weight Per panicle, NGP: Number of grain per panicle, NPT: Number of Productive tillers, NT: Number of tillers, PC: Principal component, PCA: Principal component analysis, PEF: Percentage of empty florets, PH: Plant Height, PL: Panicle length, RCBD: Randomized Complete blocked design, S: Spikilet, SD: Stomatal density, Vg: Genetic variability, Vp: Phenotypic variability

### INTRODUCTION

Wheat (*Triticum aestivum* L.) is a primary food source in subtropical regions and a secondary food source in tropical regions, specifically wheat flour (Tharise et al. 2014). The availability of wheat promotes food security in Indonesia, since wheat flour is the second-largest source of carbohydrates after rice (Rafiuddin et al. 2020). Therefore, the increasing population growth of Indonesia has led to an annual increase in the demand for wheat, specifically wheat flour. The demand for wheat flour can be fulfilled by increasing wheat importation. According to Statistics Indonesia (2021), Indonesia imported about 10.29 million tons of wheat flour in 2020. However, this import danger to Indonesia food sovereignty. Therefore, the promotion of wheat farming is a better way of resolving the demand in Indonesia. Generally, wheat plants grow better in the highlands of Indonesia at 1000 m above sea level due to their optimal temperature ranging between 8-10°C (Farid et

al. 2021). However, this condition is rare and only found in small areas of specific regions (Rafiuddin et al. 2020). The development of highland wheat will promote the growth of other products, such as vegetables, fruit, tea, coffee, etc. (Farid et al. 2020), which are considered to be more in demand and profitable than wheat farming (Rafiuddin et al. 2020). Therefore, wheat farming should be limited to the lower plains or the overall tropical climate.

Wheat farming in tropical lands is associated with the development of heat-resistant wheat varieties that require a diversity induction method for the selection process (Bovina et al. 2014). Irradiation mutation is one of the effective methods for developing such varieties randomly (Xiong et al. 2019). Furthermore, this method employs random abrasion of genes and chromosomes, which produces new viable variants (Sun et al. 2006; Farid and Ridwan 2018; Nasaruddin et al. 2018; Nur et al. 2018). Farid et al. (2020) have conducted the breeding procedure until it reached a preliminary yield test on several lines that

are adaptive to the tropical lands. These lines have to continue to the following stage, namely the advanced yield trial. However, the effectiveness of the selection in advanced yield trials must be carried out with precision and accuracy. One method for improving the accuracy is by selecting based on a combination of primary and secondary parameters.

According to Wening et al. (2019), Fadhli et al. (2020), and Anshori et al. (2021), the yield test can be conducted effectively through a combination of primary and secondary characters in the form of a selection index, which is a method of selecting characters based on their weight (Simeao et al. 2015). Subsequently, the combination is substituted into a multiple regression formula. The order of the selection indices for each genotype serves as a guide in the selection process (Rajamani et al. 2016; Islam et al. 2017). According to Sabouri et al. (2008), Figueiredo et al. (2012), and Anshori et al. (2019), the selection index is effective and equivalent to the weight assigned to each character. The determination of the selection index can be conducted by multivariate analysis, which is used to interpret and simplify data containing a large number of variables (Mattjik and Sumertajaya 2011). Additionally, this analysis can aid breeders in determining the characteristics and weighting values for the selection process (Arain et al. 2018; Farid et al. 2020).

The cultivation of wheat in tropical lands requires advanced yield trialing of wheat mutant lines using a selection index based on multivariate analysis. Therefore, this study aims to determine the selection effectiveness of tropical wheat mutant lines by using a selection index and multivariate analysis.

## MATERIALS AND METHODS

### Study area

The study was conducted from July to October 2019 in three locations. The advanced yield trial was conducted in two locations, i.e. Kelara Village and Ujung Bulu Village,

whereas, the validation of the selection was conducted in Padang Magek Village (Table 1).

### Plant material

A nested Randomized Complete Block Design (RCBD) was used in this study, where the replications were nested in the locations. The study included 16 wheat genotypes, which consisted of 13 wheat mutant lines irradiated by gamma radiation (1 line from Dewata variety, 4 lines from Selayar variety, 8 lines from Nias variety) and three control varieties (Dewata, Selayar, and Nias) (Table 2). Each variety was replicated three times at each location, for a total of 96 experimental units, and also each genotype consisted of three rows of plants cultivated in an array. Each experimental unit has ten clumps for observation samples.

### Procedures

The procedure for planting wheat was in accordance with Nasruddin et al. (2018) and Farid et al. (2020). Plots were measured at 2.5x1 m with 0.5 m apart in each replication. Each plot consisted of three rows 0.25 m apart. The seeds were planted in holes, which were formed from the array. Each of the wheat genotypes weighed 7 g and was planted in each row at 20x5 cm apart. The maintenance activities in wheat cultivation include watering, fertilizing, weeding, and controlling plant pest organisms. Watering was conducted every day, while the first and second weeding occurred on the fourth and sixth week after planting, respectively. Fertilization was performed twice on the tenth and thirtieth day after planting using Urea, SP36, and NPK fertilizer (16-16-16) at a dose of 150 kg ha<sup>-1</sup>, 200 kg ha<sup>-1</sup>, and 100 kg ha<sup>-1</sup>, respectively. The second fertilization was conducted using Urea at a dose of 150 kg ha<sup>-1</sup>. Subsequently, chemical pesticides (Deltamethrin 25 g L<sup>-1</sup>, and Profenofos 500 g L<sup>-1</sup>) were used to control plant pest organisms. Harvesting was performed when 80% of the plants in a plot were physiologically ripe, as indicated by yellowing panicles, stems, and leaves of the plant, as well as hardening of the wheat seeds.

**Table 1.** The detailed information of field locations

Aspects	Field locations		
	Kelara Village	Ujung Bulu Village	Padang Magek Village
Detailed location	South Tolo Sub-district, Jenepono District, South Sulawesi, Indonesia	Rumbia Sub-district, Jenepono District, South Sulawesi, Indonesia	Rambatan Sub-district, Tanah Datar District, West Sumatra, Indonesia
Altitude (asl.)	135 m	800 m	400 m
Coordinates	5°34'59" S 119°48'31" E	5°24'56" S 119°54'19" E	0°29'37" S 100°34'04" E

**Table 2.** The detailed information of genotype used in this study

Genotype	Lines	Wild type mutant	Genotype	Lines	Wild type mutant
G1	N 200 2.3.3	Nias variety	G9	410 8.3.1	Selayar variety
G2	N 250 4.6.2	Nias variety	G10	N 200 2.4.6B	Nias variety
G3	D 200	Dewata variety	G11	N 200 2.5.2	Nias variety
10	S 300 7.9.1	Selayar variety	G12	S 300 2.1	Selayar variety
G5	N 350 3.7.1	Nias variety	G13	N 350 3.6.2	Nias variety
62	N 200 1.7.1	Nias variety			Dewata
G7	N 350 3.1.3	Nias variety			Selayar
G8	S 300 3.6.1	Selayar variety			Nias
					Control variety

### Parameters observations

The data used in this study were obtained from field observations in each experimental plot, such as plant height, number of tillers, number of productive tillers, stomatal density, chlorophyll index, days to flowering, days to harvest, grain filling time, panicle length, spikelet, floret, percentage of empty florets, number of grains per panicle, grain weight per panicle, 1000 grain weight, and the yield (Nasruddin et al. 2018; Farid et al. 2020).

### Data analysis

Analysis of variance (ANOVA) and phenotypic-genetic correlation were used to analyze the observed data. The selected characters and correlations from ANOVA were subjected to in-depth analysis using path and principal component analysis (PCA), which served as the determinants of the optimal selection index based on index heritability as an early index validation technique (Anshori et al. 2019). Subsequently, the mutant lines were selected using the best selection index and a rating system. The ANOVA and PCA analysis used the STAR 2.0.1 software by IRRI. Meanwhile, path analysis used excel 2016 version and correlation analysis used META-R by CIMMYT. The selected mutant lines were propagated and validated using regression analysis by MINITAB 17 version (Anshori et al. 2020).

## RESULTS AND DISCUSSION

### Analysis of variance on the character of wheat growth in two locations

The analysis of variance results showed that there was a significant environmental influence on various characteristics, such as number of tillers, chlorophyll index, days to flowering, days to harvest, spikelet, percentage of empty florets, number of grains per panicle, grain weight per panicle, and the yield (Table 3). Genotype variance had a highly significant effect (\*\*) on plant height, number of tillers, number of productive tillers, flowering days, harvest

days, time of grain filling, panicle length, grain, percentage of empty buds, the weight of grain per panicle, and yield. The stomata density, number of grains per panicle, and the weight of 1000 grains have affected significantly (\*) on genotype variance, while chlorophyll index was not significantly affected by its variance (ns). The interaction variance is very significantly affected on plant height, stomatal density, days to flowering, grain filling time, percentage of empty florets, number of grains per panicle, and grain weight per panicle. The panicle length and the yield were just affected significantly by interaction variance. Identification of genetic parameters has focused on heritability. Generally, the result of the heritability analysis showed that there was a high heritability of certain traits, such as the number of tillers, number of productive tillers, days to flowering, days to harvest, grain weight per panicle, and the 1000 grain weight. The panicle length, spikelet, and yield were all classified as having a medium heritability. However, the plant height, stomatal density, chlorophyll index, grain filling time, percentage of empty florets, and the number of grains per panicle of plants were all classified as having a low heritability based on the analysis of variance and the heritability of the yield.

### Establishment of a selection index for lowland wheat yield test

The results of phenotypic and genetic correlations on the yield are presented in Table 4. The results of phenotypic correlation showed that there was a significant relationship between the yield and other characteristics, such as the number of tillers, number of productive tillers, panicle length, stomatal density, spikelet, and grain weight per panicle of 0.81, 0.78, 0.86, 0.69, 0.66, and 0.89, respectively. Meanwhile, the results of genetic correlation showed that there was a significant relationship between yield and other characteristics, such as the number of tillers, the number of productive tillers, spikelet, grain weight per panicle, and 1000 grain weight at a value of 1.00 each, as well as panicle length at 0.65.

**Table 3.** Analysis of variance (ANOVA) of growth characteristics in the test of the advanced yield of wheat mutants

Characters	MS			Genetic parameters			
	Environment (E)	Genotype (G)	E x G	CV	Vg	Vp	h <sup>2</sup>
Plant height (cm)	119.93ns	64.88**	98.73**	7.65	0.00	16.45	0.00
Number of tillers (tillers)	43.34**	4.21**	1.20ns	15.89	0.50	0.70	71.39
Number of productive tillers (tillers)	5.59ns	3.26**	1.22ns	16.61	0.34	0.54	62.53
Stomatal density (mm <sup>2</sup> )	214.05ns	72.09*	75.83**	14.53	0.00	12.64	0.00
Chlorophyll Index (CCI)	9618599.69**	42.17ns	38.69ns	1.33	0.58	7.03	8.25
Days to flowering (day)	2981.29**	28.30**	13.18**	1.89	2.52	4.72	53.45
Days to harvest (day)	3488.11**	10.33**	1.84ns	1.35	1.41	1.72	82.18
Time in filling time (day)	17.23ns	27.35**	27.35**	3.96	0.00	4.56	0.00
Panicle length (cm)	0.02ns	0.80**	0.58*	6.19	0.04	0.13	27.09
Spikelet (seed)	396.74**	3.72**	2.15ns	8.70	0.26	0.62	42.20
Percentage of empty florets (%)	3014.59**	92.49**	185.73**	8.92	0.00	30.95	0.00
Number of grains per panicle (seed)	2408.61**	23.80*	58.64**	15.41	0.00	9.77	0.00
Grain weight per panicle (g)	3.14**	0.10**	0.05**	20.35	0.01	0.02	53.22
1000 grain weight (g)	168.88ns	197.08*	55.31ns	30.93	23.63	32.85	71.94
The yield (ton)	249.39**	5.89**	3.59*	36.50	0.38	0.98	39.05

Note: MS: mean square, CV: coefficient of variance, Vg: genotypic variance, Vp: phenotypic variance, h<sup>2</sup>: broad heritability, \*: significant at level 0.05, \*\*: significant at level 0.01, ns: no significant

The result of the path analysis on the yield is presented in Table 5 there was a very significant positive and direct effect on the yield by the number of productive tillers and the grain weight per panicle at 0.78 and 0.92, respectively. Conversely, the number of tillers had a very significant negative and direct effect on the yield at -0.78. Meanwhile, there was a high indirect effect by the number of productive tillers and grain weight per panicle on other characteristics related to the yield at 2.50 and 2.86, respectively. Lastly, the number of tillers had the most negative effect on other characteristics related to the yield at -2.61.


The results of the principal component analysis showed that four principal components had an eigen-value greater than 1 at a cumulative proportion of 0.8239 (Table 6). The greatest variance was observed in the principal component 1 (PC1) at 0.4707 (47.07%), which also showed the greatest variance of yield eigenvectors at -0.3137. Therefore, PC1 could be used to determine the weight of the selection index. The weight of the modeled selection index consists of three formulations, which include: (i) Path analysis index = the yield + 0.78 number of productive tillers + 0.92 grain weight per panicle. (ii) PCA index = 0.314 the yield + 0.332 number of productive tillers + 0.342 grain weight per panicle. (iii) Path - PCA index = 0.314 the yield + 0.259 number of productive tillers + 0.315 grain weight per panicle

**Selection and validation selection index on wheat yield test in lowland**

The actual value and standard deviation as the basis in forming the index value (Table 7). The results of selection index analysis showed that the PCA index has the highest heritability index (48.16%). Meanwhile, the path index and path-PCA have heritability index of 46.26 and 46.64, respectively. Therefore, the PCA index was used in selecting the tropical wheat genotypes. Based on this the PCA index selection, there were seven promising wheat lines, namely N 250 4.6.2 (G2), D 200 (G3), S 300 7.9.1

(G4), N 350 3.7.1 (G5), N 350 3.1.3 (G7), N 200 2.5.2 (G11), and N 350 3.6.2 (G13), which were better than the comparison Dewata variety and had a positive index value.

The validation results were analyzed using regression analysis (Figure 1). According to the analysis, the PCA index could be predicted the yield in Padang Magek Village with a formula as  $2.126 + 0.2971 \text{ PCA index}$ . Although the b or gradient value was not close to 1, the PCA index has a good determination value of 57.2% (Figure 1). Besides that, all the samples were within the prediction interval (PI) or the confidence interval (CI), whereas all mutant lines fell within or close to the CI. Therefore, the PCA index has been considered to have good validation.



**Figure 1.** Regression analysis in index value on the yield in Padang Magek Village, Rambatan Sub-district, Tanah Datar District, West Sumatra Province, Indonesia (note: Y\_Pad: Yield in Padang Magek Village)

**Table 4.** Phenotypic and genotypic correlations for all wheat growth characters in the lowlands

Traits	PH	NT	NPT	SD	CD	FA	HA	GFT	PL	S	PEF	NGP	GWP	1000-GW
<b>Phenotypic correlation</b>														
The yield	0.50	0.81**	0.78**	0.86**	0.35	0.15	0.06	-0.12	0.69**	0.66**	-0.28	0.48	0.89**	0.42
<b>Genetic correlation</b>														
The yield	NA	1.00**	1.00**	NA	-0.4	0.51	0.22	-0.55	0.65**	1.00**	NA	NA	1.00**	1.00**

Note: PH: plant height, NT: Number of tillers, NPT: Number of productive tillers, SD: Stomatal density, CD: Chlorophyll density, DF: Days to flowering, DH: Days to harvest, GFT: Grain filling time, PL: Panicle length, S: Spikelet, PEF: Percentage of empty florets, NGP: Number of grain per panicle, GWP: Grain weight Per panicle, 1000-GW: 1000 Grain weight, \*\*significant at level 0.01.

**Table 5.** Path analysis of wheat the yield based on characters that are significantly correlated with the yield

Characters	Direct effect	Indirect influence						Residual
		NT	NPT	PL	S	GWP	1000-GW	
Number of tillers	<b>-0.78**</b>		0.75	0.26	-0.25	0.8	0.01	0.13
Number of productive tillers	<b>0.78**</b>	-0.75		0.23	-0.25	0.76	0.01	0.13
Panicle length	0.37	-0.56	0.49		-0.21	0.59	0.02	0.13
Spikelet	-0.32	-0.62	0.62	0.24		0.71	0.01	0.13
Grain weight per panicle	<b>0.92**</b>	-0.68	0.64	0.23	-0.24		0.02	0.13
1000 grain weight	0.4	-0.19	0.12	0.15	-0.1	0.04		0.13
Total		-2.61	2.50	0.96	-0.95	2.86	0.07	

Note: NT: Number of tillers, NPT: number of productive tillers, PL: panicle length, S: spikelet, GWP: grain weight per panicle, 1000-GW: 1000 grain weight, \*\* significant at level 0.01

**Table 6.** Analysis of the principal components of the growth character of wheat in the advanced yield trial

Characters	PC1	PC2	PC3	PC4	PC5
Plant height	-0.2051	-0.3049	0.3754	-0.2537	0.0580
Number of tillers	-0.3450	0.0123	-0.0732	-0.0447	0.2094
Number of productive tillers	-0.3322	-0.0149	-0.1458	-0.0086	0.1557
Stomatal density	-0.2499	-0.1964	0.1420	0.2229	-0.3218
Chlorophyll index	-0.2178	-0.3166	-0.2132	-0.1263	0.4796
Days to flowering	-0.0465	0.5253	0.2574	-0.2269	0.2459
Days to harvest	-0.0215	-0.0098	0.2470	-0.7761	-0.0535
Grain filling time	0.0421	-0.5514	-0.1025	-0.2453	-0.2707
Panicle length	-0.2796	0.1704	0.1915	0.0774	0.0530
Spikelet	-0.3268	-0.0405	-0.0958	0.0974	0.0437
Empty florets	-0.3268	-0.0405	-0.0958	0.0984	0.0435
Percentage of empty florets	0.1404	-0.3166	0.3149	0.2243	0.5826
Number of grains per panicle	-0.2627	0.2299	-0.3627	-0.1761	0.0504
Grain weight per panicle	-0.3415	-0.0145	0.0025	0.0206	-0.1936
1000 grain weight	-0.1371	0.0410	0.5566	0.2113	-0.0576
The yield	-0.3137	0.0227	0.1830	0.0484	-0.2625
Standard deviation	2.7442	1.6005	1.2912	1.1929	0.9075
Proportion of Variance	0.4707	0.1601	0.1042	0.0889	0.0515
Cumulative Proportion	0.4707	0.6308	0.735	0.8239	0.8754
EigenValues	7.5305	2.5617	1.6672	1.4229	0.8235

Note: PC: Principal component

**Table 7.** Selection index based on path analysis, PCA, and combination of path analysis and PCA

G	Actual value			Standardization			Index		
	Number of productive tillers	Grain weight per panicle	The yield	Number of productive tillers	Grain weight per panicle	The yield	Path	PCA	Path-PCA
G7	6.45	0.92	5.05	1.33	1.97	1.52	4.37	1.59	1.44
G5	6.39	0.84	5.25	1.26	1.38	1.73	3.98	1.43	1.3
G13	6.61	0.75	4.37	1.56	0.68	0.84	2.68	1.01	0.88
G4	6.39	0.76	4.05	1.25	0.8	0.52	2.23	0.85	0.74
G11	5.45	0.78	4.03	-0.03	0.93	0.49	1.33	0.46	0.44
G2	6.11	0.68	3.49	0.88	0.15	-0.05	0.78	0.33	0.26
G3	5.78	0.69	3.73	0.43	0.23	0.19	0.74	0.28	0.24
Dewata	5.22	0.65	3.87	-0.33	-0.05	0.34	0.03	-0.02	0
Nias	5.39	0.62	3.52	-0.1	-0.35	-0.02	-0.42	-0.16	-0.14
G10	4.89	0.58	4.27	-0.78	-0.62	0.73	-0.44	-0.24	-0.17
G6	5	0.64	3.08	-0.63	-0.15	-0.47	-1.1	-0.41	-0.36
G9	4.84	0.57	2.76	-0.85	-0.72	-0.79	-2.12	-0.78	-0.69
Selayar	4.39	0.63	2.79	-1.46	-0.24	-0.75	-2.12	-0.8	-0.69
G12	5.14	0.55	2.37	-0.44	-0.87	-1.18	-2.32	-0.81	-0.76
G1	5	0.52	2.22	-0.63	-1.08	-1.33	-2.81	-0.99	-0.92
G8	4.39	0.4	1.76	-1.46	-2.04	-1.79	-4.81	-1.74	-1.58
Heritability index							46.26	48.16	46.64

## Discussion

Analysis of variance is used as a basis for evaluating the influence of genetic, environmental, and the interaction between both components in a population of genotypes in plant breeding (Syukur et al. 2015; Farid et al. 2020; Anshori et al. 2021). There are many pattern responses among genetic, environmental, and the interaction effect in ANOVA. The interactive pattern has in group 1 consists of plant height, stomatal density, grain filling time, and panicle length, and group 2 consists of the number of tillers, days to harvest, and spikelet. The first group was significantly affected by genetics and the interaction effect. However, it does not have an environmental variance. It means that the average environment of this group character relatively same, but there was a dynamic response pattern among genotypes in two environments. Group 2 is not significantly affected by interaction; however, the environment and genotype of these characters have a significant effect. It means that the changed response between environments is stable (Anshori et al. 2019). Although, the focus of analysis in advanced yield trials is the pure genotype effect that frees from the interaction effect (Hidayatullah et al. 2018; Liu et al. 2019; Akbar et al. 2021).

The genotype provides an early indication of the effectiveness of plant breeding experiments. According to Syukur et al. (2015), diverse genotypes facilitate the selection in plant breeding. This study showed that genotype had a significant effect on almost all parameters, which gave rise to effective selection. Selection in certain environments requires the existence of effective selection characteristics due to environmental factors interactions between the genotype and environment (Anshori et al. 2019; Anshori et al. 2020; Farid et al. 2020). The yield is the main trait, which exhibits a significant response to the two sources of diversity. This shows that the selection process is based on relatively unstable yield, as supported by the heritability analysis, which indicates a moderate degree of inheritance. Heritability analysis is a type of genetic analysis, which is frequently used to determine the magnitude of the genetic influence on the observed phenotype diversity. According to Syukur et al. (2015), line selection is effective when carried out on highly heritable characteristics of populations. Therefore, the selection of the yield-enhancing and the yield-supporting characteristics is effective.

The application of multivariate analysis techniques, such as phenotypic and genetic correlation, path analysis, and PCA is an effective assessment package for character selection. These methods have been applied by Anshori et al. (2019) and Anshori et al. (2021) on salinity-tolerant rice, as well as Fadhlil et al. (2020) on drought-tolerant rice and maize, respectively, and lastly, Farid et al. (2021) on hydroponic wheat selection. Correlation analysis was the fundamental concept of this analysis. Phenotypic-genetic correlation analysis was important in the selection process, specifically when genotypes are evaluated in several environments. Additionally, this was consistent with Lozada et al. (2019), which stated the critical importance of phenotypic-genetic correlation in the selection process,

specifically when based on secondary characters. Generally, genetic analysis can separate environmental influences and interactions in the correlation analysis, such that the difference between phenotypic and genetic correlation becomes the basis for assessing environmental interactions (Singh and Chaudhary 2007; Anshori et al. 2019; Baye et al. 2020). Environmental interactions will affect the correlation analysis when there is a difference between phenotypic and genotypic correlation. Therefore, this concept often serves as the foundation for more detailed analysis (Anshori et al. 2019). Although, characters with significant genetic correlations are still included in the in-depth analysis. According to phenotypic-genetic correlation analysis, potential characteristics for further analysis included the number of tillers, the number of productive tillers, panicle length, the number of spikelets, grain weight per panicle, and 1000-grain weight.

The path analysis is quantified the direct effect of a certain trait on the diversity of its main characters or the yield (Singh and Chaudhary 2007). This analysis is very important in assessing selection characteristics due to the insufficiency of correlation analysis in this study. Fadhlil et al. (2020) demonstrated the effectiveness of this analysis on maize during drought stress testing, while Baye et al. (2020), Bisht et al. (2020), Nada et al. (2021), and Farid et al. (2021) demonstrated the effectiveness of this analysis on wheat mutant. The number of productive tillers and grain weight per panicle were identified as characteristics with the potential to support the yield through path analysis. Therefore, both characteristics could be used to construct the selection index.

The number of productive tillers and grain weight per panicle were effective secondary characters for wheat and rice. This is also supported by Anshori et al. (2019), Akbar et al. (2021) and Anshori et al. (2021) for rice and Arya et al. (2017), Ibrahim (2019), Farid et al. (2020), and Farid et al. (2021) for wheat. Wheat and rice, as well as some Poaceae cereal crops, increase their growth area through the use of tillers or axillary shoots (Hussien 2014). Tiller growth becomes important in wheat and rice production since each tiller has the potential to produce grains (Sidwell et al. 1976; Kato et al. 2000; Huang et al. 2003; Kumar et al. 2007). Meanwhile, the yield potential of cereal crops, such as corn and sorghum is largely determined by the size of the cob and panicle since they rely on apical dominant growth (Dixit et al. 2014). According to Nukasani et al. (2013), Ayer et al. (2017), Bisht et al. (2020), Nada et al. (2021), and Farid et al. (2021), grain weight per panicle is another important trait that contributes to the wheat yield. The combination of these two characteristics with the yield is an optimal choice in developing the selection index. However, the selection index should be considered based on the priority value or diversity of the characteristics. This can be accomplished using the path analysis, principal component analysis (PCA), or a combination of PCA and path analysis (Path-PCA) (Table 5), which shows genotypic selection index. The selection index of the principal component is based on eigenvalues, which denote negative values. However, the eigenvalues of the three selection characteristics can be

weighted by a positive sign since the negative sign is restricted to the direction of variance and not to the absolute value (Jolliffe 2002). The effectiveness of the three indices was determined using the heritability analysis of the selection index, which shows 25% effectiveness of genetically weighted combinations (Ramos et al. 2014; Gazal et al. 2017). According to Table 3 and 7, the heritability of the selection index is greater than the heritability of the yield in isolation. Furthermore, the combination of several secondary characteristics with the yield can increase the genetic potential of the main characteristics, such as the yield, which increases the effectiveness of the selection over using only the yield. Farid et al. (2020) and Farid et al. (2021) reported similar effectiveness on wheat selection, Anshori et al. (2019) on rice under salinity stress, Rahajeng and Rahayuningsih (2017) on sweetpotato in two environments.

The selection index was then validated using regression analysis, which is used to determine the effectiveness of the model. In regression analysis, the determination value is an important component of the validation assessment (Al-Salim et al. 2017). The model is more suited to the selection accuracy at a higher determination value (Chun et al. 2021). The concept for the assessment was developed by Farid et al. (2021) and Anshori et al. (2021). The regression determination value shows that the selection index model was quite good since it exceeded 57% (Anshori et al. 2021; Farid et al. 2021). Additionally, none of the genotypes was outside the prediction interval, which is a time interval between two predictions in a model. The amount of selected tropical wheat mutant lines in this study also has the same amount reported by Sari et al. (2016). Besides that, these selected genotypes have a higher yield than the stable tropical wheat lines from Ashari et al. (2012). Therefore, this selection index was considered effective for use in multi-location wheat genotype selection.

In conclusion, the using selection index and multivariate analysis in the selection of tropical wheat mutant lines are considered effectively with high determination validation as 57%. Wheat the yield was strongly supported in this study by secondary characteristics, such as the number of productive tillers and grain weight per panicle. Also, PCA analysis was considered effective in constructing the selection index. The selection index in this analysis was calculated as follows: selection index = 0.314 the yield + 0.332 number of productive tillers + 0.342 grain weight per panicle. Consequently, seven lines were selected based on the yield, 10th included N 250 4.6.2 (G2), D 200 (G3), S 300 7.9.1 (G4), N 350 3.7.1 (G5), N 350 3.1.3 (G7), N 200 2.5.2 (G11), and N 350 3.6.2 (G13). These mutant lines would be recommended in multi-location testing to release tropical wheat varieties in Indonesia.

#### ACKNOWLEDGEMENTS

We would like to acknowledge the Ministry of Research, Technology, and Higher Education of the

Republic of Indonesia for funding and all members of the project who contributed to the implementation of the fieldwork.

#### REFERENCES

- Akbar MR, Bambang SP, Iswari SD, Willy BS, Sugiyanto, Anshori MF. 2021. Agronomic and yield selection of doubled haploid lines of rainfed lowland rice in advanced yield trials. *Biodiversitas* 22 (7): 3006-3012. DOI: 10.13057/biodiv/d220754.
- Al-Salim SHF, Maysoun MS, Ragheb HAA, Abbas LA. 2017. Correlation regression analysis in sorghum under different levels of nitrogen. *J Sci Agric* 1: 69-78. DOI: 10.25081/jsa.2017.v1i031.
- Anshori MF, Purwoko BS, Dewi IS, Ardie SW, Suwarno WB. 2019. Selection index based on multivariate analysis for selecting doubled-haploid rice lines in lowland saline prone area. *SABRAO J Breed Genet* 51: 161-174.
- Anshori MF, Purwoko BS, Dewi IS, Ardie SW, Suwarno WB. 2020. Cluster heatmap for detection of good tolerance trait on doubled-haploid rice lines under hydroponic salinity screening. *IOP Conf. Series: Earth and Environmental Science*. IPB University, Bogor.
- Anshori MF, Bambang SP, Iswari SD, Sintho WA, Willy BS. 2021. A New approach to select doubled haploid rice lines under salinity stress using indirect selection index. *Rice Sci* 28 (4): 368-378. DOI: 10.1016/j.rsci.2021.05.007.
- Araim SM, Sial MA, Jamali KD, Laghari KA. 2018. Grain yield performance, correlation, and cluster analysis in elite bread wheat (*Triticum aestivum* L.) lines. *Acta Agrobot* 71: 1747. DOI: 10.5586/aa.1747.
- Arya VK, Singh J, Kumar L, Kumar R, Kumar P, Chand P. 2017. Genetic variability and diversity analysis for yield and its components in wheat (*Triticum aestivum* L.). *Indian J Agric Sci* 51: 128-134. DOI: 10.18805/ijare.v0i0F.7634.
- Ashari S, Waluyo B, Yulianah I, Kendarini N, Jusuf M. 2012. Stability of wheat genotypes adapted in tropical medium and lowland. *Agrivita* 34 (1): 75-83. DOI: 10.17503/Agrivita-2012-34-1-p075-083.
- Ayer DK, Sharma A, Ojha BR, Paudel A, Dhaka K. 2017. Correlation and path coefficient analysis in advanced wheat genotypes. *SAARC J Agri* 15 (1): 1-12. DOI: 10.3329/sja.v15i1.33155.
- Baye A, Berihun B, Bantayehu M, Derebe B. 2020. Genotypic and phenotypic correlation and path coefficient analysis for yield and yield-related traits in advanced bread wheat (*Triticum aestivum* L.) lines. *Cogent Food Agric* 6: 1752603. DOI: 10.1080/23311932.2020.1752603.
- Boy R, Bronnazzi A, Gasparini G, Sestili F, Palumbieri S, Botticella E, Lafiandra D, Mantovani P, Massi A. 2014. Development of a 52-LING resource in durum wheat for reverse- and forward-genetic uses. *Crop Pasture Sci* 65 (1): 1-14. DOI: 10.1071/CP13226.
- Bisli Dhaka NS, Kumar N, Malik R, Kumar S. 2020. Path analysis studies of EMS-mutagenized mutant population of hexaploid wheat (*Triticum aestivum* L.). *Intl J Chem Stud* 8 (6): 391-394. DOI: 10.22271/chemi.2020.v8.i6f.10799.
- Chu Z, Wang C, Bi RT, Zhu HF, He P, Jing YD, Yang WD. 2021. Winter wheat yield estimation based on assimilated Sentinel-2 images with the CERES-Wheat model. *J Integr Agric* 20 (7): 1958-1968. DOI: 10.1016/S2095-3119(20)63483-9.
- Dixit, AK, Singh, A, Shahi, JP, Kumar P, Singh RK. 2014. Character association studies and path coefficients in maize (*Zea mays* L.) under rabbinic conditions. *Tren dalam Biosains* 7 (17): 2479-2481. [Indonesian]
- Farid M, Farid M, Rafiuddin, Effendi R, Azrai M, Anshori MF. 2020. Multivariate analysis to determine secondary trait in selecting adapted hybrid corn lines under drought stress. *Biodiversitas* 21 (8): 3617-3624. DOI: 10.13057/biodiv/d210826.
- Farid M. 2018. Growth and production of various wheat genotypes at various PEG concentration in hydroponic. *Agrotech J* 3: 21-26.
- Farid M, Ridwan I. 2018. Tolerance limits of Indonesian rice varieties to drought and salinity in germination phase using PEG and NaCl as selection agents. *IOP Conf. Series: Earth and Environmental Science*. Hasanuddin University, Makassar, 2018.
- Farid M, Nasaruddin, Musa Y, Anshori MF, Ridwan I, Hendra J. 2020. Genetic parameters and multivariate analysis to determine secondary

- traits in selecting wheat mutant adaptive on tropical lowlands. *Plant Breed Biotechnol* 8 (4): 368-377 DOI: 10.9787/PBB.2020.8.4.368.
- Farid M, Nasaruddin YM, Ridwan I, Anshori MF. 2021. Effective screening of tropical wheat mutant lines under hydroponically induced drought stress using multivariate analysis approach. *Asian J Plant Sci* 20: 172-182. DOI: 10.3923/ajps.2021.172.182.
- Figueiredo UJ, Nunes JAR, Valle CB. 2012. Estimation of genetic parameters and selection of *Brachiaria humidicola* progenies using a selection index. *Crop Breed Appl Biotechnol* 12: 237-244. DOI: 10.1590/S1984-70332012000400002.
- Gazal A, Nehvi FA, Lone AA, Dar ZA, Wani MA. 2017. Smith hazel selection index for the improvement of maize inbred lines under water stress conditions. *Intl J Pure App Biosci* 5 (1): 72-81. DOI: 10.18782/2320-7051.2444.
- Hidayatullah A, Purwokol BS, Dewi IS, Suwarno WB. 2018. Agronomic performance and yield of doubled haploid rice lines in advanced yield trial. *SABRAO J Breed Genet* 50 (3): 242-253.
- Huang XQ, Coster H, Ganai MW, Roder MS. 2003. Advanced backcross QTL analysis for the identification of quantitative trait loci alleles from wild relatives of wheat (*Triticum aestivum* L.). *Theor Appl Genet* 106: 1379-1389. DOI: 10.18805/ijare.v0i0F.7634.
- Hussien A, Elahe, David SH, Maria MA, Gari JM, Laura R. 2014. Genetics of tillering in rice and barley. *Plant Genome* 7: 1-20. DOI: 10.3835/plantgenome2013.10.0032.
- Ibrahim AU. 2019. Genetic variability, correlation and path analysis for yield and yield components in F6 generation of wheat (*Triticum aestivum* Em. Thell.). *IOSR J Agric* 12: 17-23. DOI: 10.9790/2380-1201011723.
- Islam MR, Kayess MO, Hasanuzzaman M, Rahman MW, Uddin MJ, Zaman MR. 2017. Selection index for genetic improvement of wheat (*Triticum aestivum* L.). *J Chem Biol Phys Sci* 7 (1): 1-8.
- Kato K, Miura H, Sawada S. 2000. Mapping QTL controlling grain yield and its components on chromosome 5A of wheat. *Theor Appl Genet* 101: 1114-1121. DOI: 10.1007/s001220051587.
- Kumar N, Kulwal PL, Balyan HS, Gupta PK. 2007. QTL mapping for yield and yield contributing traits in two mapping populations of bread wheat. *Mol Breed* 19: 163-177.
- Liu H, Tessema BB, Jensen J, Cericola F, Andersen JR, Sørensen AC. 2019. ADAM-Plant: A software for stochastic simulations of plant breeding from molecular to phenotypic level and from simple selection to complex speed breeding programs. *Front Plant Sci* 9: 1926. DOI: 10.3389/fpls.2018.01926.
- Loz JN, Jayfred VG, Brian PW, Arron HC. 2019. Genomic prediction and indirect selection for grain yield in US Pacific northwest winter wheat using spectral reflectance Indices from high-throughout phenotyping. *Intl J Mol Sci* 21 (65): 1-18. DOI: 10.3390/ijms21010165.
- Jolliffe 2002. I.T. Jolliffe Principal Component Analysis (2nd Edition). Springer-Verlag, New York.
- Mattjik AA, Sumertajaya IM. 2011. Multivariate analysis using SAS. *Statistika F-MIPA IPB Press, Bogor*. [Indonesian]
- Nada RF, Hassan EE, Hassan AA, Attia ZMA. 2021. Analysis of genetic correlation between some traits in M2 wheat mutant as influenced by sodium azide under water stress. *J Product Dev* 26 (3): 463-478. DOI: 10.21608/jpd.2021.181117.
- Nasaruddin, Farid M, Musa Y, Iswoyo H. 2018. Assessment and selection of M3 generation of wheat mutants adaptive in lowland. *IOP Conf. Series: Earth and Environmental Science* 157 (1): 012051 DOI: 10.1088/1755-1315/157/1/012051.
- Nukasani V, Potdukhe NR, Bharad S, Deshmukh S, Shinde SM. 2013. Genetic variability, correlation and path analysis in wheat. *J Wheat Res* 5 (2): 48-51.
- Nur A, Syahrudin K, Azrai M, Farid M. 2018. Genetic by environment interaction and stability tropical wheat lines in Indonesia medium-plains. *IOP Conf. Series: Earth and Environmental Science* 157: 012049.
- Rafiuiddin, Amin AR, Nur, Farid M, Sari A. 2020. Evaluation on growth and production of convergent breeding wheat (*Triticum aestivum* L.) genotypes adaptive to lowland. *IOP Conf. Series: Earth and Environmental Science* 486: 012098 DOI: 10.1088/1755-1315/486/1/012098.
- Rahajeng W, Rahayuningsih SA. 2017. Evaluation for yield and yield contributing parameters of orange-fleshed sweetpotato genotypes in two environments. *Nusantara Biosci* 9: 275-281. DOI: 10.13057/nusbiosci/n090306.
- Ramos HCC, Pereira MG, Viana AP, da Luz LN, Cardoso DL, Ferregueti GA. 2014. Combined selection in backcross population of papaya (*Carica papaya* L.) by the mixed model methodology. *Amer J Plant Sci* 5: 2973-2983. DOI: 10.4236/ajps.2014.520314.
- Rajamani S, Sreekanth M, Naik V S, Ratnam M. 2016. Selection indices for yield attributing characters improvement in pigeon pea (*Cajanus cajan* L. Millspugh). *Intl J Life Sci Scient Res* 2 (2): 127-129.
- Sabouri H, Rabiei B, Fazlalipour M. 2008. Use of selection indices based on multivariate analysis for improving grain yield in rice. *Rice Sci* 15 (4): 303-310. DOI: 10.1016/S1672-6308(09)60008-1.
- Sari L, Purwito A, Sopandie D, Purnamaningsih R, Sudarmonowati E. 2016. Agronomic characterization of wheat mutants (*Triticum aestivum*) of M3 generation planted in Sukabumi. *Biosaintifika* 8 (3): 352-360. DOI: 10.15294/biosaintifika.v8i3.6612.
- Sidwell RJ, Smith EL, McNew RW. 1976. Inheritance and interrelationships of grain yield and selected yield-related traits in a hard red winter wheat cross. *Crop Sci* 16: 650-654. DOI: 10.2135/cropsci1976.0011183X001600050013x.
- Simeao R, Adriane S, Cacilda V, Marcos DR, Sergion M. 2015. Genetic evaluation and selection index in tetraploid *Brachiaria ruziziensis*. *Plant Breed* 135 (2): 1-8. DOI: 10.1111/pbr.12353.
- Singh RK, Chaudhary BD. 2007. *Biometrical Methods in Quantitative Genetic Analysis*. Kalyani Publisher, New Delhi.
- Sun JL, Du XM, Zhou ZL, Pan ZE, Pang BY. 2006. Law of genetic variation of the progeny of cotton (*Gossypium hirsutum* L.) irradiated by  $\gamma$  Ray. *Acta Agric Bor Sin* 21 (2): 27-33.
- Statistics Indonesia. 2021. *Statistics Indonesia 2021*, Jakarta: Statistics Indonesia. [Indonesian]
- Syuk M, Sujiprihati S, Yunianti R. 2015. *Plant Breeding Techniques (Revised Ed.)*. Penebar Swadaya, Depok. [Indonesian]
- Tharise N, Julianti E, Nurminah M. 2014. Evaluation of physico-chemical and functional properties of composite flour from cassava, rice, potato, soybean and xanthan gum as alternative of wheat flour. *Intl Food Res J* 21 (4): 1641-1649.
- Wening RH, Indrastuti AR, Bambang SP, Willy BS, Nurul K. 2019. Screening of submergence tolerant rice under artificial condition based on multiple selection indices. *Plant Breed Biotechnol* 7 (4): 360-374. DOI: 10.9787/PBB.2019.7.4.360.
- Xiang H, Chinyun Z, Huijun GYX, Linshu Z, Jiayu G, Shirong Z, Yuping D, Luxiang L. 2019. Transcriptome sequencing reveals hotspot mutation regions and dwarfing mechanisms in wheat mutants induced by  $\gamma$ -ray irradiation and EMS. *J Radiat Res* 61 (1): 44-57. DOI: 10.1093/jrr/rz075.

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

17%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	<a href="http://www.e-sc.org">www.e-sc.org</a> Internet Source	1%
2	<a href="http://repository.univ-tridianti.ac.id">repository.univ-tridianti.ac.id</a> Internet Source	1%
3	<a href="http://usnsj.com">usnsj.com</a> Internet Source	1%
4	Adinda Asri Laraswati, Rusnadi Padjung, Muh Farid, Nasaruddin Nasaruddin, Muhammad Fuad Anshori, Amin Nur, Andi Isti Sakinah. "Image Based-Phenotyping and Selection Index Based on Multivariate Analysis for Rice Hydroponic Screening under Drought Stress", Plant Breeding and Biotechnology, 2021 Publication	1%
5	Muhammad Fuad Anshori, Bambang Sapta Purwoko, Iswari Saraswati Dewi, Sintho Wahyuning Ardie, Willy Bayuardi Suwarno. "A New Approach to Select Doubled Haploid Rice Lines under Salinity Stress Using Indirect Selection Index", Rice Science, 2021 Publication	1%

6	<a href="http://www.scielo.cl">www.scielo.cl</a> Internet Source	1 %
7	<a href="http://oes.chileanjar.cl">oes.chileanjar.cl</a> Internet Source	1 %
8	<a href="http://garuda.ristekdikti.go.id">garuda.ristekdikti.go.id</a> Internet Source	1 %
9	Submitted to Higher Education Commission Pakistan Student Paper	1 %
10	M Farid, Nasaruddin, R M Amin, I Ridwan. " Tolerance of Wheat mutant ( ) genotypes in lowland against limited water availability ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1 %
11	<a href="http://mafiadoc.com">mafiadoc.com</a> Internet Source	1 %
12	<a href="http://om.ciheam.org">om.ciheam.org</a> Internet Source	<1 %
13	<a href="http://www.ijcmas.com">www.ijcmas.com</a> Internet Source	<1 %
14	A Yassi, K Mustari, A Guricci, E Syam'un, M Riadi, T Dariati, S N Adyla. " Growth and production of lowland rice ( L.) with water management systems on the application of various combination of fertilizers and planting	<1 %

systems ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

15

Karansher S. Sandhu, Lance F. Merrick, Sindhuja Sankaran, Zhiwu Zhang, Arron H. Carter. "Prospectus of Genomic Selection and Phenomics in Cereal, Legume and Oilseed Breeding Programs", Frontiers in Genetics, 2022

Publication

<1 %

16

[core.ac.uk](https://core.ac.uk)

Internet Source

<1 %

17

Rafiuddin, A R Amin, A Nur, M Farid, A T I Sari. "Evaluation on growth and production of convergent breeding wheat (Triticum aestivum L.) genotypes adaptive to lowland", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

18

[docsdrive.com](https://docsdrive.com)

Internet Source

<1 %

19

[mail.ccrjournal.com](https://mail.ccrjournal.com)

Internet Source

<1 %

20

K Syahrudin, A Nur, M Azrai, S H Larekeng. "Character evaluation and selection of tropical wheat lines in the long period of the dry season and high temperature", IOP

<1 %

# Conference Series: Earth and Environmental Science, 2019

Publication

---

21 M Farid BDR, Kahar Mustari, Risman Arisandi, Nasaruddin, Ambo Ala, Nurlina Kasim, Amirullah Dachlan, Ifayanti Ridwan. " Response of M5 wheat ( L.) mutant to low altitude growing condition ", IOP Conference Series: Earth and Environmental Science, 2020  
Publication

---

22 [jdmlm.ub.ac.id](http://jdmlm.ub.ac.id)  
Internet Source

---

23 M Farid BDR, Nasaruddin, H Iswoyo, I Ridwan, F Arsyad. "Analysis of heritability and correlation of agronomic character towards the yield of several m6 generation of wheat mutants (Triticum aestivum L.) in the lowlands", IOP Conference Series: Earth and Environmental Science, 2020  
Publication

---

24 [link.springer.com](http://link.springer.com)  
Internet Source

---

25 [j-tropical-crops.com](http://j-tropical-crops.com)  
Internet Source

---

26 [pr.hec.gov.pk](http://pr.hec.gov.pk)  
Internet Source

---

[textroad.com](http://textroad.com)

27

Internet Source

&lt;1 %

28

"New Horizons in Wheat and Barley Research", Springer Science and Business Media LLC, 2022

Publication

&lt;1 %

29

[iosrjournals.org](http://iosrjournals.org)

Internet Source

&lt;1 %

30

Hassan Auda Awaad. "Sustainable Agriculture in Egypt", Springer Science and Business Media LLC, 2022

Publication

&lt;1 %

31

M F Anshori, M Farid, Nasaruddin, Y Musa, H Iswoyo, A I Sakinah, M Arifuddin, A A Laraswati. "Development of image-based phenotyping for selection characters of rice adaptability on the seedling salinity screening", IOP Conference Series: Earth and Environmental Science, 2021

Publication

&lt;1 %

32

S. Pandiangan, S.T. Trina, S. Saragih. " Response of rice ( L.) on seedling age and number of seedlings per planting hole ", IOP Conference Series: Materials Science and Engineering, 2018

Publication

&lt;1 %

33

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

<1 %

34

Submitted to University of Westminster

Student Paper

<1 %

35

academic.oup.com

Internet Source

<1 %

36

assets.researchsquare.com

Internet Source

<1 %

37

ia804505.us.archive.org

Internet Source

<1 %

38

ijoeear.blogspot.com

Internet Source

<1 %

39

jurnal.uns.ac.id

Internet Source

<1 %

40

www.chemijournal.com

Internet Source

<1 %

41

Nasaruddin, M Farid, Y Musa, H Iswoyo.

"Assessment and selection of M3 generation of wheat mutants adaptive in lowland", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

42

disco.uni-muenster.de

Internet Source

<1 %

journal.ipb.ac.id

43

Internet Source

&lt;1 %

44

[krishikosh.egranth.ac.in](http://krishikosh.egranth.ac.in)

Internet Source

&lt;1 %

45

Ashish B. Rajurkar, C. Muthukumar, A. Bharathi, Helen Baby Thomas, R. Chandra Babu. "Saturation mapping of consistent QTLs for yield and days to flowering under drought using locally adapted landrace in rice (*Oryza sativa* L.)", NJAS - Wageningen Journal of Life Sciences, 2019

Publication

&lt;1 %

46

H. N. Shahi, M. S. Maskina, P. S. Gill. "Effect of different levels of gypsum application on soil characteristics and growth and yield of rice (*Oryza sativa* L.)", Plant and Soil, 1978

Publication

&lt;1 %

47

Saima Mir Arain, Mahboob Ali Sial, Karim Dino Jamali, Khalil Ahmed Laghari. "Grain yield performance, correlation, and cluster analysis in elite bread wheat (*Triticum aestivum* L.) lines", Acta Agrobotanica, 2018

Publication

&lt;1 %

48

[downloads.hindawi.com](http://downloads.hindawi.com)

Internet Source

&lt;1 %

49

[media.neliti.com](http://media.neliti.com)

Internet Source

&lt;1 %

50

[www.usnsj.com](http://www.usnsj.com)

Internet Source

&lt;1 %

51

Abeyot Tessema, Sentayehu Alamerew, Taye Kufa, Weyessa Garedew. "Variability and Association of Quality and Biochemical Attributes in Some Promising *Coffea arabica* Germplasm Collections in Southwestern Ethiopia", *International Journal of Plant Breeding and Genetics*, 2011

Publication

&lt;1 %

52

Huijun Guo, Zhihui Yan, Xiao Li, Yongdun Xie, Hongchun Xiong, Yunchuan Liu, Linshu Zhao, Jiayu Gu, Shirong Zhao, Luxiang Liu. "Development of a High-Efficient Mutation Resource with Phenotypic Variation in Hexaploid Winter Wheat and Identification of Novel Alleles in the TaAGP.L-B1 Gene", *Frontiers in Plant Science*, 2017

Publication

&lt;1 %

53

M Farid BDR, Nasaruddin, M F Anshori, A N J Chaerunnisa. " Evaluation on reproductive and productivity characters of wheat ( L.) genotypes grown in the lowlands ", *IOP Conference Series: Earth and Environmental Science*, 2020

Publication

&lt;1 %

54

R. C. Sharma, E. Duveiller. "Selection Index for Improving *Helminthosporium* Leaf Blight

&lt;1 %

# Resistance, Maturity, and Kernel Weight in Spring Wheat", Crop Science, 2003

Publication

55

[www.idosi.org](http://www.idosi.org)

Internet Source

<1 %

56

S N Hadi, Fatichin, A Fauzi, I Widiyawati, Y R Ahadiyat. "The role of phosphate solubilizing bacteria from Rhizosphere of upland rice in the growth and yield of upland rice on ultisol soil", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

57

[garuda.kemdikbud.go.id](http://garuda.kemdikbud.go.id)

Internet Source

<1 %

58

[perpus.univpancasila.ac.id](http://perpus.univpancasila.ac.id)

Internet Source

<1 %

59

[repository.ju.edu.et](http://repository.ju.edu.et)

Internet Source

<1 %

60

[www.chinaagrisci.com](http://www.chinaagrisci.com)

Internet Source

<1 %

61

A Octova, A S Muji, M Raeis, R R Putra. "Identification of Aquifer using Geoelectrical Resistivity Method with Schlumberger Array in Koto Panjang Area, Nagari Tigo Jangko, Lintau Buo Sub-District, Tanah Datar Regency", Journal of Physics: Conference Series, 2019

Publication

<1 %

62 M Farid, Nasaruddin, Y Musa, H Iswoyo, I Ridwan. "Test and Selection of Adaptive M6 Wheat Mutants Lines in the Jenepono Lowland", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

---

63 Muhammad Farid, Nasaruddin ., Yunus Musa, Ifayanti Ridwan, Muhammad Fuad Ansho. "Effective Screening of Tropical Wheat Mutant Lines under Hydroponically Induced Drought Stress Using Multivariate Analysis Approach", Asian Journal of Plant Sciences, 2020

Publication

<1 %

---

64 A Nur, M Riadi, A Yassi, M Farid, M F Anshori, A K Akfindarwan. "Selection and evaluation the corn lines from multiple-cross progeny based on targeted selection environment on acid soil", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

---

Exclude quotes  On

Exclude bibliography  On

Exclude matches

< 5 words