

[Skip to Main Content](#)

Viabilities of Odontoblast Cells Following Addition of Haruan Fish in Calcium Hydroxide

by Maria Tanumihardja

FILE	CELLS_FOLLOWING_ADDITION_OF_HARUAN_FISH_IN_CALCIIUM_HYDR OXIDE.PDF (478.91K)	WORD COUNT	4208
TIME SUBMITTED	12-JUN-2020 01:00PM (UTC+0700)	CHARACTER COUNT	22574
SUBMISSION ID	1342389686		



Viabilities of Odontoblast Cells Following Addition of Haruan Fish in Calcium Hydroxide

Maria Tanumihardja^{1*}, Sulistiyah Hastuti², Juni Jekti Nugroho³, Aries Chandra Trilaksana¹, Nurhayaty Natsir¹,
Christine Anastasia Rovani¹, Lukman Muslimin³

¹Department of Conservative Dentistry, Faculty of Dentistry, Hasanuddin University, Makassar 902424, Indonesia; ²Rumah Sakit Umum Daerah dr Abdul Rivai, Berau 77372, Indonesia; ³Department of Pharmaceutical Chemistry, Sekolah Tinggi Ilmu Farmasi Makassar, Makassar 90241, Indonesia

Abstract

BACKGROUND: Haruan fish (*Channa striata*) extract (HFE) contains all the essential amino acids and fatty acids that is believed to have therapeutic value, accelerate wound healing, and anti-inflammation.

AIM: This study was aimed to examine the viability of odontoblast MDPC-23 cell lines following the addition of HFE in toxin of *Lactobacillus* sp. and/or Ca(OH)₂.

MATERIALS AND METHODS: First, to find antiproliferative effective doses, MDPC-23 cells were treated with HFE. The cell viability was measured by 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide assay on 24 h after the last treatment, while cell death was induced by addition toxin and/or Ca(OH)₂ following adding antiproliferative effective doses of HFE (25.0, 50.0, and 100 µg/mL). Untreated cells were used as control.

RESULTS: Adding of HFE at range 25.0 until 100 µg/mL increased the MDPC-23 cells viability. MDPC-23 on toxin and/or Ca(OH)₂ reported decrease the viability of cells, while supplemented with HFE significantly increase in cell viability compared to untreated cell ($p < 0.05$).

CONCLUSION: HFE effectively increased the viability of odontoblast MDPC-23 cells and has the potency to be used together to avoid the negative side effect of Ca(OH)₂ as a capping agent.

Edited by: Filip Korinek
Citation: Tanumihardja M, Hastuti S, Nugroho JJ, Trilaksana AC, Natsir N, Rovani CA, Muslimin L. Viabilities of Odontoblast Cells Following Addition of Haruan Fish in Calcium Hydroxide. Open Access Maced J Med Sci. 2020 Apr 25; 6(1):58-63. https://doi.org/10.3889/oamjms.2020.4362
Keywords: Calcium hydroxide, Channa striata, Odontoblasts MDPC-23 cell line, Viability
Correspondence: Maria Tanumihardja, Department of Conservative Dentistry, Faculty of Dentistry, Hasanuddin University, Makassar 902424, Indonesia. E-mail: maria.tanumihardja@yahoo.com
Received: 29-Jan-2020
Revised: 07-Mar-2020
Accepted: 29-Mar-2020
Copyright: © 2020 Maria Tanumihardja, Sulistiyah Hastuti, Juni Jekti Nugroho, Aries Chandra Trilaksana, Nurhayaty Natsir, Christine Anastasia Rovani, Lukman Muslimin
Funding: This research did not receive any financial support
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution International License (CC BY-NC 4.0)

Introduction

Inflammation of the pulp tissue may be initiated by various toxins, necrotic cells, or stimulation of odontoblasts. Damage of the odontoblasts induces the release of bioactive molecules and initiates an inflammatory response in the pulp [1]. On occasions, inflammation ceases rapidly and enables complete pulp healing with the formation of a barrier of reactionary dentin by the original surviving odontoblasts and/or reparative dentin by newly differentiated odontoblast-like cells in animal models [2], [3].

Odontoblast survival is sensitive to the remaining dentin thickness (RDT). ARDT of 1 mm has been suggested to protect the pulp from the harmful constituents of dental materials [4]. When the pulp-dentin complex of RDT is <0.5 mm, it should be protected from any injury with pulp capping materials [5]. The primary goal of capping the pulp is to preserve the underlying pulp and maintain pulp vitality through the formation of reparative dentin, which in turn increases the survival of teeth life expectancy and improves the overall oral health [6].

Calcium hydroxide (Ca(OH)₂) is the most widely used capping agent [7]. It has an antimicrobial effect and dissolves calcium ions to create an alkaline environment that can stimulate reparative dentin [8]. However, Ca(OH)₂ dressings have some drawbacks, tunnel defects were reported as incomplete dentin bridge formation that compromises the integrity of reparative dentin [9]. It is assumed that the high pH of Ca(OH)₂ is toxic to odontoblast cells, which may cause severe inflammation and incomplete cell regeneration.

Several measures were proposed in maximizing reparative dentin formation, such as to optimize direct capping techniques, improve the biocompatibility of the materials, and enhance the biological responses of the pulp tissues [10]. In a previous study, pulp inflammation was reduced by applying anti-inflammatory substances, but the results were not satisfactory [11].

Haruan fish (HF), *Channa striata*, is a natural ingredient rich in albumin that has been widely studied to have the ability to regenerate cells. Some studies reported that HF plays a role in increasing the viability of fibroblast cells [12], decreases macrophages [13], and

lymphocyte count in inflammation [14]. The chemical composition of HF content of amino acid glycine and proline was collagen sources potential [15]. HF can increase the formation of new capillaries, which play an essential role in the healing process [16]. Palmitic acid which is isolated from *Gaultheria itoana* Hayata and *Sarcopyramis alensis* can suppress pro-inflammatory cytokines tumor necrosis factor- α , interleukin-6, cyclooxygenases-2, and nitric oxide in lipopolysaccharide-stimulated mouse peritoneal macrophages [17]. Therefore, we assumed that HF might be used in dental materials to improve the effectiveness of Ca(OH)_2 as a pulp capping agent since no studies have ever been performed in odontoblast cells.

Cell culture laboratory studies are beneficial for investigating some of the basic cell interaction/response requirements of the materials, although they have limitations because these cultures studies do not represent cell behavior in 3D structures [18]. This preliminary study is aimed to evaluate the viability of odontoblast MDPC-23 cells after the addition of Haruan fish (*Channa striatus*) extract (HFE) in toxin of *Lactobacillus* sp. and/or Ca(OH)_2 .

Materials and Methods

Materials

Materials that were used in this study include Dulbecco's modified eagle medium (DMEM), fetal bovine serum (FBS), penicillin-streptomycin 10000 U/mL, trypsin-EDT 0.25%, sodium dodecyl sulfate, and amphotericin B were purchased from Gibco, Grand Island, NY, while 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide (MTT) was purchased from Sigma-Aldrich, St. Louis, MO.

Research design

This was an experimental laboratory study with "post-test only control group design," conducted in the Laboratory of Hasanuddin University Hospital under the approval of the Ethics Commission of the Dental and Oral Hospital, Faculty of Dentistry, Hasanuddin University (0015/PL.09/KEPK FKG-RSGM/UNHAS/2018). The experiment design can be seen in Figure 1.

Sampling and extraction

The selected HF that is still alive came from the ponds of Tanjung Redeb, Berau, East Kalimantan. The fish used was firstly cleaned from the heads, gills, and scales. The flesh was extracted by steaming in a closed pan for ± 30 min until the pale




Figure 1: Scheme of the experiment protocol in the present study. Cell viability following application of HFE (A), induced by toxin (B), induced by toxin and Ca(OH)_2 .

yellowish liquid was obtained and separated. The liquid extract was further processed in powder using the freeze dryer (Büchi, German) and stored at the freezer, called HFE. Gently weighed 10 mg of HFE and dissolved in 10 mL DMEM medium (1000 $\mu\text{g/mL}$) as a stock solution, the diluted until reach 25.0; 50.0; and 100 $\mu\text{g/mL}$.

Cell lines culture

The following established *in vitro* cell lines were applied in odontoblast MDPC-23 cells (ATCC-CRL 2537). The MDPC-23 cells were cultured in DMEM medium supplemented 10% FBS, penicillin (100U/mL), streptomycin (100 $\mu\text{g/mL}$), and 0.5% fungizone in plastic flasks (Iwaki, Japan) at 37°C in a 5% CO_2 incubator. After reaching confluence, cells were harvested by 0.25% trypsin-EDTA.

Determination the effect of HFE on cell viability

The harvesting cells were seeded into well plate 96 (2×10^5 cell/mL). All the cells were treated with HFE and incubated for 24h. The medium discarded and washed with PBS. The cell viability was measured by MTT assay. About 100 μL MTT (0.5 mg/mL in DMEM) was added to each well, and the plate was incubated for 4 h at 37°C in a CO_2 incubator. Formed formazan crystals were solubilized by adding 100 μL of dimethyl sulfoxide per well. Finally, the intensity of the dissolved formazan crystals (purple color) was quantified using the microplate reader at 540 nm. The viability of the cell was calculated using the formula: Cell viability (%) = Mean OD/Control OD \times 100%.

Experimental models

Spread the harvested cells into well plate 96 (10^5 cell/mL). To induce injury and death, the cells were contacted with 0.04 μL toxin of *Lactobacillus* sp.

alone or with Ca(OH)₂ for 4 h then added with HFE. The number of cells was determined using the MTT method as described before (determination the effect of HEF 15 cell viability). The viability of the cell was calculated using the formula: Cell viability (%) = Mean OD/Control OD × 100%.

Statistical analysis

Data were collected in an Excel worksheet and analyzed by ANOVA. Data are presented as the mean ± standard deviation after deductible with baseline (untreated cell as 100%). Differences with p < 0.05 were considered significant. All of the experiments were evaluated in three replicates (n = 3).

Results

To select the optimal HFE concentrations for the cell culture experiments, we investigated the effect of HFE on cell viability after incubation for 24h by MTT reduction assay. Viability of MDPC-23 cells increased significantly following the addition of each concentration of HFE, as seen in Figure 2. It is meaning a safe dosage range to be used for further biological.

The effect of HFE on the viability of MDPC-23 cells induced *Lactobacillus* sp. toxin was illustrated in Figure 3. In general, despite the presence of the toxin, the viability of MDPC-23 cells increased significantly on each concentration of HFE. The toxin group indicates the decreasing of cell viability (-5.51 ± 7.00%) without the influence of the HFE. For the HFE treated cells with the lowest concentration (25.0 µg/mL) showed a significant increase in viability with a value of 9.5 ± 7.00%. When moderate and high concentration (50.0 and 100.0 µg/mL), they showed a relatively similar effect on the percentage of viability cells with a value of 26.38 ± 14.00 and 28.04 ± 10.07%.

The viability of MDPC-23 cells increased significantly, albeit Ca(OH)₂ and toxin of *Lactobacillus* sp. were added on each concentration of HFE as seen in Figure 4 at a dose-dependent manner. On the other hand, HFE 100 µg/mL showed the highest effect on the viability of the cell.

Discussion

The dental pulp is a connective tissue that responds to stimuli and insults. Maintenance of healthy pulp tissue is essential for the function and survival of




Fig 2. The effect of HFE on the viability of MDPC-23 cells. Cell viability following incubation with indicated concentrations of HFE for 24h was determined using the 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide assay. Cell viability is expressed as a percentage of the increasing cell compared to the untreated cells. *p < 0.05 (HFE 100 µg/mL compared to 25 and 50 µg/mL)




Fig 3. The effect of HFE on the alteration of cell viability of MDPC-23 cells was induced by 0.04 µL toxin of *Lactobacillus* sp. The viability of cells was determined after 24h of incubation by the 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide assay. Cell viability is expressed as a percentage of the increasing cell compared to the untreated cells. *p < 0.05 (All treatment group)




Fig 4. Viability of MDP C-23 cells was induced by 0.04 µL toxin of *Lactobacillus* sp. and Ca(OH)₂. The viability of cells was determined after 24 h of incubation by the 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazolium bromide assay. Cell viability is expressed as a percentage of the increasing cell compared to the untreated cells. *p < 0.05 (All treatment group compared to toxin and Ca(OH)₂.)

tooth [2]. Formation of secondary and tertiary dentin serves to protect the tooth from caries, infections, and traumatic dentin exposure. On the other hand, trauma, wear, or disease may impair the barrier and lead to pulpal necrosis by many insults during the lifetime of the tooth [19], [20]. Various toxins, necrotic cells, or stimulation of odontoblasts may all cause harmful effects on the pulp tissue with both acute and chronic responses depending on the magnitude and duration of the insult [21].

Cavity base materials are designed for the purpose of protecting the pulp from damage by toxins or material components. The potential ability of cavity base materials in inducing tertiary dentin formation was believed to be necessary for the function. This ability is mediated through the release of growth factors and other bioactive molecules [22]. Ca(OH)₂ is a cavity base material that induces tertiary dentin formation through the activation of blast cells in the pulp. These cells are recruited as new hard tissue-forming cells replacing the destroyed odontoblasts. Ca(OH)₂ produces a zone of superficial necrosis, which serves as a scaffold for an underlying hard tissue repair, forming a dentin bridge. This bridge is seldom continuous and complete, which compromises the barrier [9]. Several bioactive molecules have been experimentally tested for intended use as pulp capping and reparative dentin-forming materials and some products have shown some potential.

HFE is a natural ingredient that has been studied to have the ability to regenerate fibroblast cells [12]. This is in line with the results of this study that the viabilities of the odontoblast cell line significantly increase at each concentration of HFE added to the cells (Figure 2). The viability of these cells slightly decreased when a toxin was incorporated (Figure 3). This may be explained that the toxin used was an inactive toxin of *Lactobacillus* sp., with the purpose of injuring the cells that simulate the conditions of the oral cavity where bacterial toxin triggers injury and inflammation of the pulp cells [23]. In addition, bacterial toxins do not directly contribute to the cell's viability but act as an inflammatory agent. Lipoteichoic acid, the endotoxin of *Lactobacillus* sp. can provoke the immune response and activate odontoblast cells through TLR2, phosphorylate the intracellular pathways, activate the transcription factor nuclear factor-κB, and mitogen-activated protein kinase, which in turn synthesize various pro-inflammatory mediators and disrupt the normal functioning of the body [6], [24]. Therefore, the addition of HFE could increase the viability of cells in the presence of toxin (Figure 3).

Ca(OH)₂ has long been recommended for capping the exposed pulp. The main goal of a pulp capping treatment is to maintain pulp vitality of the affected tooth. A cascade of physiologic, biochemical events occurs, starting from a slight inflammatory reaction to generating chemical signals that will lead dental pulp progenitor cells to migrate to the affected area. These cells have been shown to proliferate and differentiate to form a mineralized bridge that will separate the pulp tissue from the noxious stimuli to avoid further damage and eventually to enable the pulp to heal [18], [25]. Contrastly higher dose, Ca(OH)₂ is toxic to normal cells, which may cause severe inflammation and incomplete cell regeneration [26].

In this study, Ca(OH)₂ and toxin significantly decreased the viabilities of cells to 24.7% (Figure 4). Ca(OH)₂ is known to have a high pH that has a degenerative effect on cells, although it is generally accepted as the material of choice for capping the exposed pulp. It was reported that in some cases, the pulp remains chronically inflamed, and the tissue degenerates. The hydroxyl ions released are free radical and highly reactive. It causes protein denaturation, proliferation inhibition, and ATP depletion resulting in necrosis and cell lysis [11], [27]. This may partly explain the decreased cells viability. However, the addition of HFE significantly reincreased the viability of the cells and was in accordance with a dose-dependent manner (Figure 4). These results are similar to the previous study, who reported the highest percentage increase in the number of fibroblast cells at a concentration of 100%. This might be attributed to the high albumin content in HFE which is a crucial nutrient for recovery of damaged cells [12], [14], [16], [28], [29]. Albumin also plays an important role in maintaining the capillary

osmotic pressure that can prevent excessive edema and avoid cell death [30].

Albumin of bovine serum has ever been studied and developed as a biodegradable cavity liner, a hydrogel with the potency as a capping agent. Hydrogel constitutes injectable scaffolds, cells, and bioactive molecules carriers for tissue engineering. Using the rat model, hydrogel showed homogenous dentinal bridge formation compared to Dycal capping with tunnel disturbances after 3 weeks. The inflammation was weak or barely detectable after hydrogel capping. In contrast, 3ycal capping has a continuously high pH which causes superficial burn covering a scar at the pulp surface with the consequence of inflammation in a limited area. This inflammation is still observed after 4 weeks and a mild necrotic area was noticed between the reparative dentinal bridge and vital pulp [31]. Investigations of HFE are still in their infancy, and whether it has the potency as albumin of bovine serum needs further studies.

Under the limitations of this study, HFE has shown its effectiveness to increase the viability of odontoblast MDPC-23 cells despite the presence of toxin and $\text{Ca}(\text{OH})_2$. Further studies should be carried out using an animal model to evaluate pulp tissue reactions when exposed to these bioactive substances that might be utilized in dentistry.

Conclusion

The HFE enhanced the proliferation of odontoblast cells on MDPC-23 cells induced toxin and $\text{Ca}(\text{OH})_2$. It suggests that HFE has the potency to be used together with $\text{Ca}(\text{OH})_2$ as a capping agent to induce the formation of reparative dentin.

References

- Schuh CMAP, Benso B, Aguayo S. Potential novel strategies for the treatment of dental pulp-derived pain. *Pharmacological approaches and beyond Front Pharmacol*. 2019;10:1068. <https://doi.org/10.3389/fphar.2019.01068>. PMID:31620000
- Farges JC, Alliot-Licht B, Renard E, Ducret M, Gaudin A, Smith AJ, et al. Dental pulp defence and repair mechanisms in dental caries. *Mediators Inflamm*. 2015;2015:230251 <https://doi.org/10.1155/2015/230251>. PMID:26538821
- Ricucci D, Lughin S, Lin LM, Spångberg LS, Tay FR. Is hard tissue formation in the dental pulp after the death of the primary odontoblasts a regenerative or a reparative process? *J Dent*. 2014;42(9):1156-70. <https://doi.org/10.1016/j.jdent.2014.06.012>. PMID:25008021
- Arandi NZ. Calcium hydroxide liners: A literature review. *Clin Cosmet Investig Dent*. 2017;9:67-72. PMID:28761378
- Berbari R, Khairallah A, Kazan HF, Ezzedine M, Bandon D, Sfeir E. Measurement reliability of the remaining dentin thickness below deep carious lesions in primary molars. *Int J Clin Pediatr Dent*. 2018;11(1):23-8. <https://doi.org/10.5005/jp-journals-10005-1478>. PMID:29805230
- Yumoto H, Hirao K, Hosokawa Y, Kuramoto H, Takegawa D, Nakanishi T, et al. The roles of odontoblasts in dental pulp innate immunity. *Jpn Dent Sci Rev*. 2018;54(3):105-17. <https://doi.org/10.1016/j.jdsr.2018.03.001>. PMID:30128058
- Al-Saudi KW, Nabih SM, Farghaly AM, AboHager EA. Pulpal repair after direct pulp capping with new bioceramic materials: A comparative histological study. *Saudi Dent J*. 2019;31(4):469-75. <https://doi.org/10.1016/j.sdentj.2019.05.003>. PMID:31695296
- Kiranmayi G, Hussainy N, Lavanya A, Swapna S. Clinical performance of mineral trioxide aggregate versus calcium hydroxide as indirect pulp-capping agents in permanent teeth: A systematic review and meta-analysis. *J Int Oral Health*. 2019;11(5):235-43. https://doi.org/10.4103/jioh.jioh_122_19
- Song M, Yu B, Kim S, Hayashi M, Smith C, Sohn S, et al. Clinical and molecular perspectives of reparative dentin formation: Lessons learned from pulp-capping materials and the emerging roles of calcium. *Dent Clin North Am*. 2017;61(1):93-110. <https://doi.org/10.1016/j.cden.2016.08.008>. PMID:27912821
- Schmalz G. Strategies to improve biocompatibility of dental materials. *Curr Oral Health Rep*. 2014;1(4):222-31. <https://doi.org/10.1007/s40496-014-0028-5>
- Jaian AL, Warhadpande MM, Dakshindas DM. A comparison of human dental pulp response to calcium hydroxide and biodentine as direct pulp-capping agents. *J Conserv Dent*. 2017;20(2):129-33. <https://doi.org/10.4103/0972-0707.212247>. PMID:28855762
- Siswanto A, Dewi N, Hayatie L. Effect of haruan (*Channa striata*) extract on fibroblast cells count in wound healing. *J Dentomaxillofac Sci*. 2016;1(2):89-94. <https://doi.org/10.15562/jdmfs.v1i2.3>
- Hartini P, Dewi N, Hayatie. Extract of haruan (*Channa striata*) decreases macrophages count in inflammation phase of wound healing process. *J Dentomaxillofac Sci*. 2015;14(1):6-10. <https://doi.org/10.15562/jdmfs.v14i1.417>
- Izzaty A, Dewi N, Indah D. Extract of haruan (*Channa Striata*) decrease lymphocyte count in inflammatory phase of wound healing process effectively. *J Dentomaxillofac Sci*. 2014;13(3):176-81. <https://doi.org/10.15562/jdmfs.v13i3.411>
- Rosmawati, Abustam E, Tawali AB, Said MI. Chemical composition, amino acid and collagen content of snakehead (*Channa striata*) fish skin and bone. *Scie Res J*. 2018;6(1):1-4. <https://doi.org/10.1007/s12562-018-1248-8>
- Reza M, Dewi N, Kaustiyah I. Extract of haruan (*Channa Striata*) increase neocapillaries count in wound healing process. *J Dentomaxillofac Sci*. 2015;14(1):1-5. <https://doi.org/10.22208/jdmfs.v14i1.1>
- Lee JY, Lee HJ, Jeong JA, Jung JW. Palmitic acid inhibits inflammatory responses in lipopolysaccharide-stimulated mouse peritoneal macrophages. *Orient Pharm Exp Med*. 2010;10(1):37-43. <https://doi.org/10.3742/opem.2010.10.1.037>
- Pedano MS, Li X, Jeanneau C, Ghosh M, Yoshihara K, Van Landuyt K, et al. Survival of human dental pulp cells after 4-week culture in human tooth model. *J Dent*. 2019;86:33-40. PMID:31121243

19. Bjørndal L, Simon S, Tomson PL, Duncan HF. Management of deep caries and the exposed pulp. *Int Endod J*. 2019;52(7):949-73. <https://doi.org/10.1111/iej.13128>
20. Andreasen FM, Kahler B. Pulpal response after acute dental injury in the permanent dentition: Clinical implications-a review. *J Endod*. 2015;41(3):299-308. <https://doi.org/10.1016/j.joen.2014.11.015>
PMid:25601716
21. Yu CY, Abbott PV. Responses of the pulp, periradicular and soft tissues following trauma to the permanent teeth. *Aust Dent J*. 2016;61(Suppl 1):39-58. <https://doi.org/10.1111/adj.12397>
PMid:26923447
22. Hosoya A, Nakamura H. Ability of stem and progenitor cells in the dental pulp to form hard tissue. *Jpn Dent Sci Rev*. 2015;51(3):75-83. <https://doi.org/10.1016/j.jdsr.2015.03.002>
23. Berbari R, Fayyad-Kazan H, Ezzedine M, Fayyad-Kazan M, Bandon D, Sfeir E. Relationship between the remaining dentin thickness and coronal pulp status of decayed primary molars. *J Int Soc Prev Community Dent*. 2017;7(5):272-8. <https://doi.org/10.5005/jp-journals-10005-1478>
PMid:29026700
24. Widjiastuti I, Inatari N, Rukmo M. Stimulation of propolis extract in odontoblast like cells induced by *Lactobacillus acidophilus* inactivity to TLR2 and TNF α expression. *Odonto Dental J*. 2017;4(2):85-93. <https://doi.org/10.30659/odj.4.2.85-93>
25. Qureshi A, Soujanya E, Nandakumar, Pratapkumar, Sambashivarao. Recent advances in pulp capping materials: An overview. *J Clin Diagn Res* 2014;8(1):316-21.
PMid:24596805
26. Mostafa N, Moussa S. Mineral trioxide aggregate (MTA) vs calcium hydroxide in direct pulp capping-literature review. *Online J Den Oral Health*. 2018;1(2):1-6. <https://doi.org/10.33552/ojdoh.2018.01.000508>
27. Qadiri SY, Mustafa S. Role of calcium hydroxide in root canal therapy: A comprehensive review. *J Adv Med Dent Sci Res*. 2019;7(7):1-3.
28. Andrie M, Sihombing D. The wound healing effect of cork (*Channa striata*) extract in ointment on acute stage II opened wound in in male Wistar strain rats. *Pharma Sci Res*. 2017;4(2):88-101.
29. Laila L, Febriyenti F, Salhimi SM, Baie S. Wound healing effect of Haruan (*Channa striatus*) spray. *Int Wound J*. 2011;8(5):484-91. <https://doi.org/10.1111/j.1742-481x.2011.00820.x>
PMid:21722317
30. Michelis R, Sela S, Zeitun T, Geron R, Kristal B. Unexpected normal colloid osmotic pressure in clinical states with low serum albumin. *PLoS One*. 2016;11(7):e0159839. <https://doi.org/10.1371/journal.pone.0159839>
PMid:27453993
31. Njeh A, Uzunoglu E, Ardila-Osorio H, Simon S, Berdal A, Kellermann O, et al. Reactionary and reparative dentin formation after pulp capping: Hydrogel vs. Dycal. *Evid Based Endod*. 2016;1:e3. <https://doi.org/10.1186/s41121-016-0003-9>

Viabilities of Odontoblast Cells Following Addition of Haruan Fish in Calcium Hydroxide

ORIGINALITY REPORT

% **20**

SIMILARITY INDEX

% **14**

INTERNET SOURCES

% **13**

PUBLICATIONS

% **13**

STUDENT PAPERS

PRIMARY SOURCES

1	www.tandfonline.com Internet Source	% 2
2	www.mdpi.com Internet Source	% 1
3	evidencebasedendodontics.springeropen.com Internet Source	% 1
4	www.hindawi.com Internet Source	% 1
5	Submitted to University of Birmingham Student Paper	% 1
6	Minju Song, Bo Yu, Sol Kim, Marc Hayashi et al. "Clinical and Molecular Perspectives of Reparative Dentin Formation", Dental Clinics of North America, 2017 Publication	% 1
7	www.scielo.br Internet Source	% 1

Submitted to University of Ulster

8

Student Paper

%1

9

www.science.gov

Internet Source

%1

10

Ju-Young Lee. "Palmitic acid inhibits inflammatory responses in lipopolysaccharide-stimulated mouse peritoneal macrophages", Oriental Pharmacy and Experimental Medicine, 03/31/2010

Publication

%1

11

Submitted to Universitas Hasanuddin

Student Paper

%1

12

toxsci.oxfordjournals.org

Internet Source

%1

13

www.rjpharmacognosy.ir

Internet Source

%1

14

www.freedomwares.ca

Internet Source

%1

15

link.springer.com

Internet Source

<%1

16

Min Yang, Shengjun Wang, Suoying Wang, Jie Ma et al. "Tat-mediated Intracellular Delivery of T-bet Protein into THP-1 Cells can Induce Th1-Type Response", Immunological Investigations, 2009

<%1

17

eprint.ncl.ac.uk

Internet Source

<% 1

18

www.spandidos-publications.com

Internet Source

<% 1

19

Submitted to Oregon Health and Sciences University

Student Paper

<% 1

20

www.allbusiness.com

Internet Source

<% 1

21

univmed.org

Internet Source

<% 1

22

Submitted to Cardiff University

Student Paper

<% 1

23

diposit.ub.edu

Internet Source

<% 1

24

www.physiology.org

Internet Source

<% 1

25

Maharani Laillyza A, Dewi Puspitasar. "Effect of Channa micropeltes for Increasing Lymphocyte and Fibroblast Cells in Diabetic Wound Healing", Journal of Medical Sciences, 2018

Publication

<% 1

26

portal.koreascience.or.kr

Internet Source

<% 1

-
- 27 Submitted to Royal College of Surgeons <% 1
Student Paper
-
- 28 Submitted to University of Alabama at <% 1
Birmingham
Student Paper
-
- 29 Submitted to Chester College of Higher <% 1
Education
Student Paper
-
- 30 www.labome.org <% 1
Internet Source
-
- 31 marmaraeczacilikdergisi.com <% 1
Internet Source
-
- 32 Submitted to University of Nottingham <% 1
Student Paper
-
- 33 www.wjgnet.com <% 1
Internet Source
-
- 34 bioz.com <% 1
Internet Source
-
- 35 www.oncotarget.com <% 1
Internet Source
-
- 36 Myrna Nurlatifah Zakaria, Nisa Fitriani Nurul <% 1
Pauziah, Indah Puti Sabirin, Arief Cahyanto.
"Evaluation of Carbonate Apatite Cement in
Inducing Formation of Reparative Dentin in

Exposed Dental Pulp", Key Engineering
Materials, 2017

Publication

37 Submitted to Higher Education Commission
Pakistan <% 1
Student Paper

38 Lee, Jienny, Myung-Sunny Kim, Channy Park,
Eun-Bae Jung, Duk-Hwa Choi, Tae-Yo Kim,
Sung-Keun Moon, and Raekil Park. "Morphine
Prevents Glutamate-Induced Death of Primary
Rat Neonatal Astrocytes Through Modulation of
Intracellular Redox", Immunopharmacology and
Immunotoxicology, 2004. <% 1
Publication

39 Khaled Wagih Al-Saudi, Sameh Mahmoud
Nabih, Ahmed Mustafa Farghaly, Eman Abdul-
Aziz AboHager. "Pulpal repair after direct pulp
capping with new bioceramic materials: A
comparative histological study", The Saudi
Dental Journal, 2019 <% 1
Publication

40 Submitted to Universiti Malaysia Sarawak <% 1
Student Paper

41 A Njeh, E Uzunoğlu, H Ardila-Osorio, S Simon,
A Berdal, O Kellermann, M Goldberg. <% 1
"Reactionary and reparative dentin formation
after pulp capping: Hydrogel vs. Dycal",

Evidence-Based Endodontics, 2016

Publication

EXCLUDE QUOTES ON

EXCLUDE ON

BIBLIOGRAPHY

EXCLUDE MATCHES

< 5

WORDS