

Molecular mechanisms of lidocaine

by Agussalim Bukhari

Submission date: 10-Sep-2021 08:47AM (UTC+0700)

Submission ID: 1644922136

File name: Agussalim_Bukhari-Resiana2.pdf (1.95M)

Word count: 8716

Character count: 50217



Review

Molecular mechanisms of lidocaine

Resiana Karnina^{a,b,*}, Syafri Kamsul Arif^c, Mochammad Hatta^d, Agussalim Bukhari^e

^a Doctoral Program of Biomedical Sciences, Faculty of Medicine, Hasanuddin University, Makassar, Sulawesi Selatan, Indonesia

^b Faculty of Medicine, Muhammadiyah University of Jakarta, Banten, Indonesia

^c Department of Anesthesiology, Faculty of Medicine, Hasanuddin University, Sulawesi Selatan, Indonesia

^d Department of Microbiology, Faculty of Medicine, Hasanuddin University, Makassar, Sulawesi Selatan, Indonesia

^e Department of Nutritional Sciences, Faculty of Medicine, Hasanuddin University, Sulawesi Selatan, Indonesia

ARTICLE INFO

Keywords:

Pain
Analgesia
Inflammation
Lidocaine
Local anesthesia
Neurotransmitter

ABSTRACT

Lidocaine is an amide-class local anesthetic used clinically to inhibit pain sensations. Systemic administration of lidocaine has antinociceptive, antiarrhythmic, anti-inflammatory, and antithrombotic effects. Lidocaine exerts these effects under both acute and chronic pain conditions and acute respiratory distress syndrome through mechanisms that can be independent of its primary mechanism of action, sodium channel inhibition. Here we review the pathophysiological underpinnings of lidocaine's role as an anti-nociceptive, anti-inflammatory mediated by toll-like receptor (TLR) and nuclear factor kappa- β (NF- κ B) signalling pathways and downstream cytokine effectors high mobility group box 1 (HMGB1) and tumour necrosis factor- α (TNF- α).

1. Introduction

Post-surgery inflammation is characterized by increased blood flow and vascular permeability, the upregulation of inflammatory mediators, and leukocyte accumulation [1]. Cytokines are key modulators of inflammation and can play both pro- and anti-inflammatory roles. A dynamic balance exists between pro- and anti-inflammatory cytokines that affects organ dysfunction, immunity, infection, wound healing, and pain – all of which are associated with surgery [1,2]. Surgical injury induces endogenous mediators and activates hemodynamic, metabolic, and immune responses [1]. This immune response initiates immediately after a surgical injury. Polymorphonuclear leukocytes (PMNs), endothelial cells, macrophages, and lymphocytes are activated by the secretion of pro-inflammatory mediators including cytokines, chemokines, and other molecules including but not limited to reactive oxygen species, nitric oxide, and platelet-activating factor [3]. While essential, when unchecked, excessive inflammation can disrupt the body's immune system, potentially leading to certain inflammation-related conditions and even organ failure [4,5].

Lidocaine was first synthesized by Nils Lofgren in 1935 in the in the Stockholm laboratory of Professor Hans von Euler where Lofgren began tasting the compounds he and his colleagues had synthesized [6]. In

1943, Lofgren found that the 57th compound he tested rapidly numbed his tongue. The patent for Xylocaine® was approved in Sweden on May 11, 1948, based on Goldberg's (toxicology) and Gordh's (clinical results) papers, suggesting that lidocaine had a strong and unexpected anesthetic effect. The Food and Drug Administration approved Xylocaine® for usage in the United States in November 1948 [6,7]. Torsten Gordh's clinical testing revealed that lidocaine represented a significant improvement over procaine, the gold standard for managing surgical pain at the time [8,9]. This amide-class anesthetic is still used widely to ease the pain associated with surgery, provide neuropathic pain relief, and treat ventricular arrhythmias [10,11].

In addition to its use as a local anesthetic and anti-arrhythmic agent, lidocaine has analgesic properties for various pain conditions. The nociceptive agonist effects of intravenous lidocaine have been well established in a variety of acute and chronic pain conditions [7,12]. Moreover, preclinical and clinical data evidence the antihyperalgesic effects of parenteral lidocaine [7]. The recommended initial dose is 1–2 mg/kg administered intravenously followed by a continuous infusion of 2–4 mg/kg/h, resulting in steady plasma concentrations of 1–3 mg/ml [7,13].

The mechanism of action of lidocaine as a local anesthetic is through a blockade of voltage-gated sodium channels (VGSCs) leading to a

* Corresponding author. Faculty of Medicine, Muhammadiyah University of Jakarta, Jl. Kh. Achmad Dahlan, RW.2, Cireundeu, Kec. Ciputat Tim, Kota Tangerang, Selatan, Banten, 15419, Indonesia.

E-mail addresses: sublime_dr@yahoo.com (R. Karnina), syafrikarif@yahoo.com (S.K. Arif), hattaram@yahoo.com (M. Hatta), agussalim.bukhari@med.unhas.ac.id (A. Bukhari).

<https://doi.org/10.1016/j.amsu.2021.102733>

Received 10 July 2021; Received in revised form 13 August 2021; Accepted 15 August 2021

Available online 17 August 2021

2049-0801/© 2021 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

reversible block of action potential propagation [7]. Lidocaine affects inflammatory cells in vitro, for example by inhibiting the priming of human peripheral PMNCs and neutrophils [7,14]. Lidocaine can further reduce the release of pro-inflammatory mediators such as IL-4, IL-6, and tumour necrosis factor-alpha (TNF- α) [7].

It is well established in the literature that lidocaine exerts its anti-inflammatory effects by inhibiting the expression of pro-inflammatory cytokines, the metabolic activity of leukocytes, and the release of histamine [10,15]. The effects of lidocaine are achieved by preventing NF- κ B activation and its downstream cytokine storm [16]. Specifically, lidocaine significantly reduces TNF- α levels relative to vehicle-treated controls [17]. Therefore, lidocaine's anti-inflammatory mechanism of action at the level of receptor engagement and pro-inflammatory cytokine release in the context of post-surgical injury is the entry point for the present review.

2. Systemic effect of lidocaine in molecular biology

Lidocaine is an amide class local anesthetics used in medicine to inhibit pain sensations [17,18]. It consists of lipophilic and hydrophilic subunits connected by hydrocarbon chains. The hydrophilic portion is composed of tertiary amines (e.g., diethylamine) while the lipophilic portion is composed of unsaturated aromatic rings [e.g., para-aminobenzoic acid (PABA)] [19,20]. Based on this structure, these local anesthetics can be classified into amino-esters and amino-amides. The lipophilic portion determines the anesthetic activity of the local anesthetic drug [9]. Local anesthetics act on sodium ion channels to reduce the permeability of cell membranes, thereby blocking depolarization and preventing the conduction of the electrical impulse through which pain occurs [21].

Chemically, lidocaine [2-(diethylamino)-N-(2,6-dimethylphenyl)acetamide] contains three basic components: hydrophilic amine groups, aromatic residues, and intermediary groups that connect these two (Fig. 1) [22,23]. The amine group is a tertiary or secondary amine, between an aromatic residue group and an intermediate group connected by an amide bond. Lidocaine is a weak alkaline with pKa of 8, protein binding of 64%, and fat solubility of 1%. Lidocaine remains the drug of choice for a variety of medical procedures due to its strong anesthetic potential, fast onset of action, and wide safety limits [24,25]. Moreover, lidocaine can be administered via many routes, including topical (i.e., skin and airway), subcutaneous, intravenous, perineural, epidural, and intrathecal [22,26,27]. After intravenous administration, peak plasma levels are achieved within 3–5 min with a half-life of 30–120 min [28].

In the liver, lidocaine is dealkylated by dual-function oxidizing enzymes to the pharmacologically active metabolite, monoethylglycinexylidide (MEGX) and then metabolized by the P450 3A4 isoenzyme into N-ethylglycine (NEG) and glycinexylidide (GX). MEGX is 80% as potent as the parent drug, whilst GX is nearly ineffective [29].

In clinical practice, lidocaine is also used as a class IB antiarrhythmic




Fig. 1. Chemical structure of lidocaine [21].

drug (sodium channel blocker). Sodium channels have three basic states: (1) resting (phase 1), while they await the arrival of an action potential; (2) open/active (phase 0), during which the channel is activated and conducts a sodium current; and (3) inactivated/refractory (phase 2), after the channel has conducted a sodium current but has not yet returned to its resting state. During this refractory period, the sodium channel cannot yet be activated again. Lidocaine occupies receptors on the sodium channel in its open/active (phase 0) and inactivated/refractory (phase 2) states, which lidocaine has a high affinity for its receptors during both phases [30].

Lidocaine's effects on the central nervous system include inhibiting nicotinic and acetylcholine receptors, inhibiting presynaptic calcium channels in the dorsal root ganglion, inhibiting opioid receptors, inhibiting of neurite growth, inhibiting muscarinic cholinergic receptors, and preventing substance P from binding to natural killer (NK) cell receptors [7,31,32].

The anti-inflammatory effect of lidocaine occurs at lower concentrations than required to block sodium channels [33,34]. The effect of lidocaine on inflammation, particularly against inflammatory polymorphonuclear granulocytes (PMNs), macrophages, and monocytes is not due to blocked sodium channels [35,36].

Priming can be described as a process that gives a resting neutrophil a functional response that can be greatly amplified upon exposure to another stimulus [37,38]. The second stimulus is usually considered an activating agent or agonist. The enhanced functional response keeps the neutrophils in an active state. Thus, full neutrophil activation is a two-step process, starting with initial exposure to primary agents such as cytokines (e.g., IL-1 α , GM-CSF, and TNF- α) and antigens (e.g., pathogenic endotoxin). Priming and activation change neutrophils from a resting state to an active state, thus enabling them to perform antibacterial, pro- and anti-inflammatory functions [39]. PMN priming regulates the function of PMNs and is implicated in cases of excessive inflammatory responses that cause tissue damage [39,40]. Some potential mechanisms include local anesthetics inhibiting G-protein-coupled receptors (GPCRs) signals that mediate inflammatory responses such as lysophosphatidic acid and thromboxane A2 as well as the M1 muscarinic acetylcholine receptor. GPCRs consist of muscarinic acetylcholine receptors M1-M5, which regulate several functions of the nervous system (Fig. 2) [23]. Moreover, M1 and M4 receptors are the treatment sites for several central nervous system disorders such as Alzheimer's disease, schizophrenia, and drug addiction [23,41].

3. Lidocaine as an anti-inflammatory

Lidocaine is a potent anti-inflammatory agent, whose properties are often compared with steroids and non-steroidal anti-inflammatory drugs (NSAIDs) [42]. The definite anti-inflammatory mechanism of lidocaine remains vague; however, it is presumed that the drug affects a multitude of inflammatory processes such as phagocytosis, migration, exocytosis, and cellular metabolism. In vitro experiments on human polymorphonuclear granulocytes suggest that lidocaine the membrane-ion transporters, thereby, dysregulating cellular pH levels and eventually depressing cytokine release [43].

There is still no universal reference for lidocaine administration as an anti-inflammatory agent. Ortiz et al. conducted a double-blind, randomized trial studying the effect of endovenous lidocaine on serum inflammatory cytokine levels using bolus lidocaine of 1.5 mg/kg at the start of the procedure with a maintenance dose of 3 mg/kg until 1 h after the end of the surgery [44]. A significant reduction in serum levels of pro-inflammatory markers (IL-2, IL-6, TNF- α , and IFN- γ) was observed in the IV lidocaine group in comparison with a control group [44,45]. In this same study, there were no statistically significant differences regarding the postoperative pain intensity, morphine consumption, ileus, and hospital stay compared with the control group, meaning there were no proven secondary effects at this dose [44].

Multiple, complex mechanisms likely underlie lidocaine's anti-




Fig. 2. The mechanism of action of local anesthetics on inflammation [21].

inflammatory effects in a synergy that involves numerous pathways, receptors, cells, and mediators. More specifically, the modulation of high mobility group box 1 (HMGB1), toll-like receptor (TLR), nuclear factor kappa-β (NF-κβ), and TNF-α have been implicated in previous studies.

40

3.1. High Mobility Group Box 1 (HMGB1) cytokine activity

HMGB1 functions as a pathogenetic cytokine regulator under these conditions. HMGB1 is actively secreted from various cell types including macrophages, NK cells, dendritic cells (DCs), blood vessel endothelial, and platelets [46,47]. HMGB1 is also passively released from necrotic cells, damaged cells, and after extracellular injury [48]. Cells undergoing apoptosis release less HMGB1 than necrotic cells, yet macrophages covered with apoptotic cells can still stimulate the discharge of HMGB1 from WEHI-231, Jurkat, and HL-60 cells. Pyroptosis and caspase-1-associated necrosis are responsible for the continuous discharge of HMGB1 controlled by dsRNA-dependent protein kinase (PKR) and the inflammasome. Pyroptosis follows the activation of inflammasomes, leading to the expression of caspase-1 and its downstream effects, including generation of the cytokines IL-1β and IL-18 by the cleavage of their precursors [49].

Release mechanisms such as necrosis, macrophage activation, pyroptosis, and apoptosis discharge HMGB1 in various redox forms. Necrotic and pyroptotic cells produce thiol in its reduced form, which then binds to the chemokine CXCL12 and the CXCR4 receptor to stimulate the process of chemotaxis. Pyroptosis and TLR4 stimulation also release reduced HMGB1, disulfide bond HMGB1, C23, C45, and C106, which are all in thiol form. This form of HMGB1 then stimulates cytokine production via TLR4 signaling. Activated macrophages also release the cytokine-inducing form of HMGB1 upon TLR4 activation. Apoptotic cells release HMGB1 that is partially oxidized or completely oxidized at the critical cysteine residues. Completely oxidized HMGB1, with cysteines in the form of sulfonates, is unable to stimulate cytokines or induce chemotaxis; apoptotic cells expressing this form of oxidized HMGB1 can induce tolerance [49].

3.2. Cellular inflammatory response and HMGB1 receptors

HMGB1 is a classic pro-inflammatory mediator because [4]:

- a. Injuries and infections stimulate its release.

- b. It causes immuno-competent cells to release TNF-α, IL 1, and other related substances.
- c. It reduces symptoms of pyrexia, and the sickness syndrome in vivo;
- d. It is activated by exogenous TLR agonists and other cytokines that stimulate inflammation.
- e. It can be specifically targeted to therapeutic advantage in sterile and infectious disease syndromes associated with elevated HMGB1 levels.

18

One difference between HMGB1 and conventional PCs (e.g., TNF-α, IL-1) is it stimulates systemic inflammatory responses through receptors that report and interact with foreign substances [50]. Unlike TNF-α and IL-1, the allied plasma membrane receptor family interacts with HMGB1 and initiates signal transduction through endogenous (RAGE) and exogenous ligands (TLR2, TLR4, and TLR9) [51,52]. These processes show that HMGB1 elicits various inflammatory responses in a diversity of infections and injuries. HMGB1's ability to modulate the magnitude of the inflammatory response in clinical syndromes associated with injury is discussed below. These mechanisms have been explored in loss-of-function-type studies based on HMGB1 antagonists and/or the deletion of receptors via genetic clustering techniques [53].

HMGB1 binding to TLR4-MD2 acts as a measure of surface plasmin resonance and signal transducers that stimulate macrophages to release [4]. These processes require redox-sensitive cysteine 106, which prevents HMGB1 from adhering to TLR4, the endogenous HMGB1 receptor responsible for regulating macrophage activation, cytokine release, and tissue injury repair by activating IκB kinase (IKK)-β, IKK-α (active endotoxin only IKK-β), and active nuclear translocation NF-κβ [54] [56]. One difference between HMGB1- and endotoxin-mediated signaling, is that the former does not bind to TLR4 as readily as LPS. Another difference is the pattern of gene expression induced upon activation. While both HMGB1 and LPS significantly increase NF-κβ nuclear translocation and Akt/p38 MAPK phosphorylation, LPS increases the production of NF-κβ and TNF more than HMGB1. Furthermore, HMGB1-induced secretion of TNF exhibits a biphasic kinetic profile, while the endotoxin LPS stimulates monophasic TNF release [56].

Animal studies have shown that HMGB1 levels increase during injury due to a lack of oxygen [4]. HMGB1 protein levels rise within 1 h of reperfusion and remain elevated for up to 24 h. Treating wild-type (C3H/HeOuj) mice with anti-HMGB1 antibodies helps to protect them from liver injury, while the TLR4 deficient (C3H/HeJ) mice garner no

benefit from these antibodies [4,16,57]. HMGB1 signaling via TLR4 is an effective target for solid tumors treatment strategies such as antigen cross-presentation or chemotherapy. Renal tubular TLR4 expression in donor kidneys is indicated by HMGB1 immunoreactivity, this reveals its importance in developing kidney graft inflammation and sterile injury [4].

The impact of HMGB1 protein binding on the release of cytokines (e.g., CXCL12, TLR9, thrombospondin, syndecan, TLR2, MAC1, TREM1) in the pathogenesis of sterile infection and injury remains unknown [4]. Furthermore, HMGB1 regulates the body's inflammatory response to sterile threats and infections through TLR4 receptor-mediated signaling [58,59]. HMGB1 then reacts with CD24, a plasma protein working with Siglec-10 to suppress nuclear translocation. This process is controlled by HMGB1 and mediated TLR4 activation, not pathogen-mediated TLR activation, indicating that the outcome of HMGB1 via TLR4 signaling is different from CD24-siglec-10 when it comes to sterile damage [56].

3.3. HMGB1 pathway to NF- κ B

Receptors involved in HMGB1 binding include the receptor for advanced glycation end products (RAGE), a transmembrane, cell surface, multi-ligand member of the large immunoglobulin family (Fig. 3) [4]. Consequently, RAGE-mediated HMGB1 signaling stimulates chemotaxis, cell growth, differentiation, and the migration of immune/smooth muscle cells by engaging with cell-surface moieties such as RAGE/TLR4. Although HMGB1 and RAGE bind to each other, TLR4 still controls the secretion of HMGB1 from macrophages due to the inhibition of RAGE macrophages and the inactivation of TNF-producing macrophages by TLR2. However, TLR4 does not mediate macrophage inactivation [53].

4. TOLL-LIKE receptors (TLRs)

4.1. TLR signaling pathway

The TLR pathways and signals are initiated by a ligand binding to TLRs expressed on the plasma membrane, endoplasmic reticulum, or endosome, which induces the dimerization of the TLRs proteins. Ligand-induced dimerization of TLRs likely transposes the TIR from the cytoplasmic tail of each adjacent protein, followed by a TIR-containing adapter protein that recruits and activates protein kinases to stimulate transcription factors.

Interferon response factor 3 (IRF3), NF- κ B, activation protein 1 (AP-1), and IRF1 are the transcription factors stimulated by TLR signaling pathways upon PAMPs and DAMPs recognition, TLRs recruit TIR domain-containing adaptor proteins such as MyD88 and TRIF [60]. Furthermore, NF- κ B and AP-1 are essential proteins that mediate the expression of molecules involved in the inflammatory response such as chemokines (e.g., CCL2, CXCL), cytokines (e.g., TNF, IL-1), and endothelial adhesion molecules (e.g., N-cadherin, E-selectin). Different combinations of ligands and intermediate messengers dictate the shared and unique effects of TLRs. Upon engagement at the cell's surface, dimerized TLRs bind to the MyD88 adapter, which activates NF- κ B, while the signal adapter TRIF (TIR domain-containing IFN- β inducing adapter) activates IRF3. For all TLRs except TLR3, signaling via MyD88 activates NF- κ B and initiates a pro-inflammatory response. Via TRIF, TLR3 activates IRF3 and induces type I interferon expression. Regardless of adapter (i.e., MyD88 or TRIF), TLR-ligand interactions lead to a pro-inflammatory downstream response [61]. TLR7 and TLR9 are abundant within the cytoplasm of dendritic cells and use MyD88-dependent, TRIF-independent pathways to activate NF- κ B and IRFs. TLR7, TLR9, and TLR4 all stimulate reactions against bacterial, single-stranded (ss)RNA from viruses, viral DNA, and bacterial lipopolysaccharide (LPS) [60,62].

There is some evidence of an interaction between lidocaine and TLRs. Lidocaine (50 mM) inhibits the activation of TLR4 and subsequently also NF- κ B and mitogen-activated protein kinases (MAPKs) in LPS-stimulated murine macrophages [7].

5. Nuclear factor kappa- β (NF- κ B)

5.1. NF- κ B in inflammation

Inflammation and the actions of the adaptive and innate branches of the immune system are regulated by NF- κ B [63]. An inflammatory response comprises several activated signaling pathways that modulate the activation of pro- and anti-inflammatory regulators in resident tissue cells and blood-derived leukocytes [64]. Information on inflammation and its underlying signaling pathways is mostly obtained through studies on the IL-1, TNF, and IL-17 protein families aimed at understanding how microbes in the IL-17 family operate. IL-1 and TNF- α represent a baseline pattern of pro-inflammatory cytokines released rapidly in response to tissue injury and infection. TLR recognizes




Fig. 3. The relationship between HMGB1, TLR4, and RAGE. TLR4 binding induces cytokine secretion from macrophages and monocytes (left), meanwhile, RAGE modulates endothelial and tumor cell function (right) [3].

microbial molecular patterns and, as such, is known as a pattern recognition receptor (PRR). TLRs represent germline-encoded, non-self-recognition systems poised to induce inflammation [65]. However, endogenous ligands also activate TLR during particular forms of damage, which causes inflammation in the absence of infection. IKK kinase (IKK) and NF- κ B make use of similar signal-transduction mechanisms even though their structures differ [65,66].

6. Tumor necrosis factor- α (TNF- α)

6.1. Systemic effects of TNF in the inflammatory response

TNF receptors are expressed differently in a variety of cells and tissues; however, the pro-inflammatory effects of TNF occur based on interactions between blood vessel cells and leukocytes. These cells then initiate inflammatory processes with different temporal, spatial, and anatomical presentations using different leukocyte adhesion molecules, including E-selectin, intercellular adhesion molecule-1 (ICAM-1), and vascular cell adhesion molecule-1 (VCAM-1) [67].

This response, when combined with chemokine release (e.g., IL-8, MCP-1, IP-10) leads to the recruitment of several leukocytes independent of antigen recognition [68,69]. Furthermore, the local effects of TNF on endothelial cells lead to many classic inflammatory features. For example, TNF-induced cyclo-oxygenase 2 expression can increase the EC production of PGI₂, which results in vasodilation, and 'rubor' and 'heat' through increased local blood flow. Swelling is created by edema caused by the accumulation of endothelial fluid and other macromolecules [70]. Furthermore, TNF-induced pro-coagulant protein expression (including tissue-specific factors), anti-coagulant protein regulation, and thrombo modulation can cause intravascular thrombosis [67].

6.2. TNF to NF- κ B family signaling

The NF- κ B pathway is activated by various immunologically-relevant ligands and receptors such as the TNF receptor (TNFR), TLR, IL-1 receptor (IL-1R), and other antigen receptor groups [66,71]. Furthermore, these receptors lack any enzymatic activity, while the results of ligand binding are always relayed through the lipid bilayer and converted into kinases via adapter proteins [66].

The process that holds the ligand bonds also fosters the creation of a receptor oligomer/ligand complex [66]. TNF family signaling via NF- κ B causes several adapter proteins to bind to the ligand-receptor complex, thereby activating IKK [72]. This adapter protein belongs to that protein is well described as the domain of protein interactions participating in assembly of very complex oligomeric proteins that join ligand-receptor complexes to IKK [66]. Adapter protein interactions important for TNF-NF- κ B family signaling include death fold domain (DD), RIP (receptor-interacting protein), homotypic interaction motifs (RHIM), and TRAF (TNF receptor-associated factor) domains [66, 73].

This adapter protein directs signaling to various end pathways. In this case, the functions of transcription factors such as NF- κ B in signaling are discussed. Furthermore, changes in receptor conformation localization and increased avidity are caused by the ligand binding of adapter proteins on cytoplasmic receptor tails [66,74]. This leads to the mobilization of members of the TRAF and RIP families into a complex protein structure. Consequently, the oligomerization of these complexes is sustained by cross-receptors, multimer ligands, and the ability of proteins to form high-order trimers and oligomers [66]. Upstream kinases that control the activation of IKK and NF- κ B signaling through the TNF cytokine family include NIK (NF- κ B inducing kinase) and TAK1 (transforming growth factor- β activated kinase; MAP3K7) [66,71]. Furthermore, NIK controls the stimulation of IKK α and non-canonical pathways. The function of TAK1 in canonical pathways is NEMO-dependent; however, the requirements of TAK1 in all forms of TNF and NF- κ B signaling are not fully understood. After activation, IKK kinase (IKK-K)

triggers a traditional kinase cascade within and outside of the NF- κ B pathway [66].

7. Lidocaine in acute and chronic pain

Lidocaine exerts additional analgesic effects through other pathways. Various studies show the analgesic benefit of systemic lidocaine during surgical procedures, especially during laparoscopic abdominal surgery [75] [-] [80]. Usually, lidocaine is administered initially at doses of 1.5–2 mg/kg BW and maintained by 1.5–3 mg/BW/hr [36–3 mg/min. This results in serum levels of lidocaine that range from 0.5 to 5 μ g/ml (~2–21 μ M), which is similar to after epidural administration [75,77,78,80,81]. The benefits of lidocaine include a quick recovery, reduced time in the hospital, unaffected bowel movements, and less post-operative pain (Table 1) [7,82].

7.1. Opioid receptors

One study describes the absence of any cellular interaction between lidocaine and recombinant μ -, κ -, and δ -opioid receptors [83,84]. This is because all opioid receptors belong to a family of GPCRs lacking Gaq units, and lidocaine only influences receptors with G α q. Meanwhile, another study found that when opioids and lidocaine are administered simultaneously, they synergistically potentiate anti-nociception. However, the mechanism underlying this effect, for example, whether occurs at the receptor level (e.g., via TLR4 signaling, Kir, or Ca²⁺ channels) or involves regional/systemic interplay, has yet to be elucidated [7,83].

7.2. Toll-like receptors (TLRs)

Another study describes lidocaine's negligible inhibitory effect on TLRs. However, lidocaine (50 μ M) inhibits the activation of TLR4, along with NF- κ B and MAPKs, in LPS-stimulated murine macrophages. This mechanism is mediated by VGSCs [1].

Intravenous administration of lidocaine in rats with LPS-induced sepsis reduces organ failure significantly compared with control. It was also shown to protect against organ dysfunction through the downregulation of TLR4. The expression of TLR4, NF- κ B, and interleukin (IL)-6 was reduced in the lidocaine group by this mechanism [85]. Nebulized lidocaine prevents the respiratory system of mice from becoming inflamed by reducing TLR2 expression [86]. Further investigation is needed to determine whether these processes can be applied for chronic pain management (Table 2).

8. Lidocaine as an anti arrhythmic agent

In the 1950s, the first case of cardiac arrhythmia treated successfully with intravenous lidocaine was reported: ventricular fibrillation that occurred during cardiac catheterization was successfully reverted to a regular sinus rhythm with electric shock and lidocaine administration [87]. The dose of lidocaine used to treat ventricular arrhythmias

Table 1
The biological effects of lidocaine on various forms of chronic pain [6,81].

Source	Spontaneous pain	Hyperalgesia	Allodynia
Neuropathic pain	Reduced		
Diabetic neuropathy	Reduced		
Peripheral nerve injury	Reduced	Reduced	
Post herpetic neuralgia	Reduced	Reduced	Reduced
Chronic regional pain syndrome	Reduced at high (3 μ g/ml) plasma concentration	Reduced for cold threshold only	Reduced for cold and mechanical thresholds
Central pain	Reduced	Reduced	Reduced

Table 2

Presents the results of studies on the relationship between lidocaine and acute/chronic post-operative pain. Recent reviews highlight differences in the results obtained by these studies [6,7,6,86].

Perioperative antinociceptive effects	Pain at rest <24 h	Pain during activity	Opioid requirement	PONV
Abdominal surgery	Reduced	Reduced	Reduced	Reduced
CPSP	Odds ratio for incidence of CPSP (4–6 months)		Pain intensity (McGill pain questionnaire score)	
Overall	Reduced		Unchanged (P = 0.06)	
Breast surgery	Reduced			
Non-breast surgery	Reduced			

Abbreviations: CPSP: Chronic post-surgical pain, PONV: postoperative nausea and vomiting.

(tachycardia) is 1–1.5 mg/kg BW administered intravenously followed by 5 infusion of 1–4 mg/kg BW/hr [28,88].

The mechanism underlying the antiarrhythmic effect appears to be similar to that of procaine. Ventricular excitability is depressed with increasing doses, conduction time is slowed, and the refractory period is prolonged. Cardiac contractility is not diminished in therapeutic dosages [87]. In contrast to procaine and procaine amide, doses of lidocaine that produce comparable increases in the diastolic stimulation threshold cause no fall in blood pressure or decreased myocardial contractility.

Before amiodarone was approved, lidocaine was listed as the primary drug of choice to treat ventricular fibrillation (VF) or Ventricular tachycardia (VT) in the Advanced cardiac life support (ACLS) algorithm. Even though lidocaine had been used to treat VF/VT for years, there was a lack of evidence for its efficacy over other drugs. While lidocaine was since removed from the ACLS Algorithm, it is still considered a suitable alternative if amiodarone is ineffective or unavailable in cardiac arrest from VT/VF [89].

9. Lidocaine and covid-19-associated acute respiratory distress syndrome (ARDS)

9.1. Pathophysiology of COVID-19

COVID-19 is a disease that results from infection with the newly identified SARS-CoV2 virus. Very little has been published to date regarding its pathology and mechanism. Initial pathological changes experienced at the tissue and cellular level include death of alveolar epithelial cells, intraluminal edema, fibrin exudation, hyaline membrane formation, hemorrhage, infiltration of inflammatory cells (e.g., monocytes, macrophages, neutrophils, and lymphocytes) into the alveolar wall and lumina, and elevated levels of serum pro-inflammatory cytokines and chemokines [13].

Other studies show high levels of cytokines and chemokines in blood samples collected during SARS-CoV infection. Kong et al. reported an increase in circulating cytokines, (TNF- α , CXCL-10 [interferon gamma inducible protein 10-stimulated leukocyte activator], IL-6 and -8) [91]. Other studies confirm these pro-inflammatory cytokine findings [92, 93]. After an initial disease phase, the epithelial lining experiences less damage, reduced interstitial and alveolar fibrosis, bronchiolitis obliterans pneumonia, regeneration of type II pneumocyte hyperplasia (e.g., IL1, IL-6, IL-12, IFN- γ , TGF- β), and chemokines. During this fibrotic phase, interstitial thickening is caused by moderate fibrosis and few inflammatory cells [94].

9.2. Lidocaine, acute lung injury, and ARDS

COVID-19 can progress to acute respiratory distress syndrome

(ARDS), a serious outcome of pulmonary viral infection [95,96]. The primary underlying pathophysiology of ARDS is vascular lung injury [97]. However, local anesthesia addresses the underlying etiology by inhibiting TNF α , which signals to endothelial cells to induce neutrophil binding and increases endothelial permeability [98].

Recently, preliminary studies have explored the anti-inflammatory effects of lidocaine in the context of acute lung injury [43]. The predominant mechanism of action in the specific clinical application remains unclear; however, lidocaine might exert its anti-inflammatory effects by regulating cellular metabolic activity, migration, exocytosis, and phagocytosis by reversibly interacting with membrane proteins and lipids thereby attenuating the inflammatory response by decreasing PMN granulocyte accumulation in the lung [43]. These promising preliminary findings have encouraged pulmonary physicians to introduce lidocaine into their clinical practice.

Nebulized lidocaine appears to be a novel potential treatment for improving COVID-19-related lung injury by reducing the cytokine storms associated with the disease, which would theoretically reverse ARDS. Such, more pre-clinical research and clinical trials are warranted to further define the efficacy and safety of lidocaine for treating patients with severe ARDS due to COVID-19.

10. Conclusions

The analgesic and anti-hyperalgesic effects of systemic lidocaine have served the clinical practice of medicine for 75 years. Prospective clinical investigations have confirmed that lidocaine is a highly safe and effective therapeutic option in cases of acute and chronic pain. The positive effects of incorporating lidocaine into multimodal pain therapy also include acute lung injury applications, based on evidence from experimental animal models.

Provenance and peer review

Not commissioned, externally peer-reviewed.

Please state any conflicts of interest

The authors declare that they have no conflict of interests.

Please state any sources of funding for your research

No funding or sponsorship.

Ethical approval

Review article applicable for exemption by our Institutional review board.

Consent

This manuscript does not involve human participants, human data, or human tissue.

Author contribution

Syafri Kamsul Arif, Mochammad Hatta, Agussalim Buchari: Design, editing and writing of the manuscript, supervision of the paper, and approved the final manuscript. Resiana Kamina: Editing, final review and approved the final manuscript.

Registration of research studies

Not applicable as this is a review article.

Guarantor

Resiana Kamina.

Acknowledgement

We acknowledge Muhammad Faruk, M.D, for his help in providing us with the linguistic assistance for this article review.

References

- [1] C.H. Hsing, J.J. Wang, Clinical implication of perioperative inflammatory cytokine alteration, *Acta Anaesthesiol. Taiwanica* 53 (2015) 23–28, <https://doi.org/10.1016/j.aat.2015.03.002>.
- [2] J.M. Cicchese, S. Evans, C. Hult, L.R. Joslyn, T. Wessler, J.A. Millar, S. Marino, N. A. Cilfone, J.T. Mattila, J.J. Linderman, D.E. Kirschner, Dynamic balance of pro- and anti-inflammatory signals controls disease and limits pathology, *Immunol. Rev.* 285 (2018) 147–167, <https://doi.org/10.1111/immr.12671>.
- [3] A. Albayyan, S. McSorley, C. Roxburgh, R. Kearns, P. Horgan, D. McMillan, The effect of anesthesia on the postoperative systemic inflammatory response in patients undergoing surgery: a systematic review and meta-analysis, *Surg. Open Sci.* 2 (2020) 1–21, <https://doi.org/10.1016/j.sopen.2019.06.001>.
- [4] U. Andersson, K.J. Tracey, HMGB1 is a therapeutic target for sterile inflammation and infection, *Annu. Rev. Immunol.* 29 (2011) 139–162, <https://doi.org/10.1146/annurev-immunol-030409-101323>.
- [5] F. Josh, T.H. Soekamto, J.R. Adriani, B. Jonatan, H. Mizuno, M. Faruk, The combination of stromal vascular fraction cells and platelet-rich plasma reduces malondialdehyde and nitric oxide levels in deep dermal burn injury, *J. Inflamm. Res.* 14 (2021) 3049–3061, <https://doi.org/10.2147/JIR.S318055>.
- [6] T. Gordh, T.E. Gordh, K. Lindqvist, D.S. Warner, Lidocaine, The origin of a modern local anesthetic, *Anesthesiology* 113 (2010) 1433–1437, <https://doi.org/10.1097/ALN.0b013e3181fce4f8>.
- [7] H. Hermanns, M.W. Hollmann, M.F. Stevens, P. Lirk, T. Brandenburger, T. Piegeler, R. Werdehausen, Molecular mechanisms of action of systemic lidocaine in acute and chronic pain: a narrative review, *Br. J. Anaesth.* 123 (2019) 335–349, <https://doi.org/10.1016/j.bja.2019.06.014>.
- [8] M.H. Holmdahl, Xylocain (lidocaine, lignocaine), its discovery and Gordh's contribution to its clinical use, *Acta Anaesthesiol. Scand. Suppl.* 113 (1998) 8–12, <https://doi.org/10.1111/j.1399-6576.1998.tb04979.x>.
- [9] A.J. Johansson, Inga fischer-hjalmars (1918–2008): Swedish pharmacist, humanist, and pioneer quantum chemist, *J. Chem. Educ.* 89 (2012) 1274–1279, <https://doi.org/10.1021/ed300024g>.
- [10] C. Berger, J. Rossaint, H. Van Aken, M. Westphal, K. Hahnenkamp, A. Zarbock, Lidocaine reduces neutrophil recruitment by abolishing chemokine-induced arrest and transendothelial migration in septic patients, *J. Immunol.* 192 (2014) 367–376, <https://doi.org/10.4049/jimmunol.1301363>.
- [11] Y. Zhang, G.-J. Tao, L. Hu, J. Qu, Y. Han, G. Zhang, Y. Qian, C.-Y. Jiang, W.-T. Liu, Lidocaine alleviates morphine tolerance via AMPK-SOCS3-dependent neuroinflammation suppression in the spinal cord, *J. Neuroinflammation* 14 (2017) 211, <https://doi.org/10.1186/s12974-017-0983-6>.
- [12] F.W. Bach, T.S. Jensen, J. Kastrop, B. Stigsby, A. Dejgård, The effect of intravenous lidocaine on nociceptive processing in diabetic neuropathy, *Pain* 40 (1990) 29–34, [https://doi.org/10.1016/0304-3959\(90\)91047-M](https://doi.org/10.1016/0304-3959(90)91047-M).
- [13] W. Koppert, V. Brueckl, C. Weidner, M. Schmelz, Mechanically induced axon reflex and hyperalgesia in human UV-B burn are reduced by systemic lidocaine, *Eur. J. Pain* 8 (2004) 237–244, <https://doi.org/10.1016/j.ejpain.2003.08.007>.
- [14] A. Ploppa, R.T. Kiefer, D.M. Haverstick, D.S. Groves, K.E. Unertl, M.E. Durieux, Local anesthetic effects on human neutrophil priming and activation, *Reg. Anesth. Pain Med.* 35 (2010) 45–50, <https://doi.org/10.1097/AAP.0b013e3181c75199>.
- [15] F.F. Cruz, P.R.M. Rocco, P. Pelosi, Anti-inflammatory properties of anesthetic agents, *Crit. Care* 21 (2017) 67, <https://doi.org/10.1186/s13054-017-1645-x>.
- [16] H.-L. Wang, Y.-Q. Xing, Y.-X. Xu, F. Rong, W.-F. Lei, W.-H. Zhang, The protective effect of lidocaine on septic rats via the inhibition of high mobility group box 1 expression and NF- κ B activation, *Mediat. Inflamm.* (2013) 1–9, <https://doi.org/10.1155/2013/570370>, 2013.
- [17] I. Garutti, L. Rancan, C. Simón, G. Cusati, G. Sanchez-Pedrosa, F. Moraga, L. Olmedilla, M.T. Lopez-Gil, E. Vara, Intravenous lidocaine decreases tumor necrosis factor Alpha expression both locally and systemically in pigs undergoing lung resection surgery, *Anesth. Analg.* 119 (2014) 815–828, <https://doi.org/10.1213/ANE.0000000000000360>.
- [18] F. Wen, Y. Liu, H. Wang, W. Tang, Y.D. Hou, H.L. Wang, Lidocaine inhibits the production of IL-1 β from macrophages RAW264.7 induced with lipopolysaccharide, *Int. J. Clin. Exp. Pathol.* 10 (2017) 6582–6588.
- [19] R.S. Vardanyan, V.J. Hruby, Local anesthetics, *Synth. Essent. Drugs* (2006) 9–18, <https://doi.org/10.1016/B978-0-44452166-8/50002-9>, Elsevier.
- [20] B. Testa, Principles of drug metabolism 2: hydrolysis and conjugation reactions, *Compr. Med. Chem. II*, Elsevier, 2007, pp. 133–166, <https://doi.org/10.1016/B0-08-04504-X/00120-6>.
- [21] H.A. Fozzard, M.F. Sheets, D.A. Hanck, The sodium channel as a target for local anesthetic drugs, *Front. Pharmacol.* 2 (2011), <https://doi.org/10.3389/fphar.2011.00068>.
- [22] R.S. Vardanyan, V.J. Hruby, Antiarrhythmic drugs, *Synth. Essent. Drugs* (2006) 245–255, <https://doi.org/10.1016/B978-0-44452166-8/50018-2>, Elsevier.
- [23] M.W. Hollmann, M.E. Durieux, D.M. Fisher, Local anesthetics and the inflammatory response, *Anesthesiology* 93 (2000) 858–875, <https://doi.org/10.1097/0000542-200009000-00038>.
- [24] J.M. Mumba, F.K. Kabambi, C.T. Ngaka, Pharmacology of local anaesthetics and commonly used recipes in clinical practice, *Curr. Top. Anesthesiol., Intech*, 2017, <https://doi.org/10.5772/67048>.
- [25] D.E. Becker, K.L. Reed, Local anesthetics: review of pharmacological considerations, *Anesth. Prog.* 59 (2012) 90–101, <https://doi.org/10.2344/0003-3006-59.2.90>, quiz 102–3.
- [26] A. Taylor, G. McLeod, Basic pharmacology of local anaesthetics, *BJA Educ* 20 (2020) 34–41, <https://doi.org/10.1016/j.bjae.2019.10.002>.
- [27] A. Albayati, A. Krauthamer, C. Baaron, Pharmacologic agents used in image-guided interventions, *Image-Guided Interv.*, Elsevier, 2019, pp. 35–41, <https://doi.org/10.1016/B978-0-323-61204-3.00006-3>.
- [28] D. Darbar, Standard antiarrhythmic drugs, *Card. Electrophysiol. From Cell to Bedside*, Elsevier, 2018, pp. 1062–1075, <https://doi.org/10.1016/B978-0-323-44733-1.00112-7>.
- [29] H. Hermanns, M.W. Hollmann, M.F. Stevens, P. Lirk, T. Brandenburger, T. Piegeler, R. Werdehausen, Molecular mechanisms of action of systemic lidocaine in acute and chronic pain: a narrative review, *Br. J. Anaesth.* 123 (2019) 335–349, <https://doi.org/10.1016/j.bja.2019.06.014>.
- [30] D.B. Tikhonov, B.S. Zhovov, Mechanism of sodium channel block by local anesthetics, antiarrhythmics, and anticonvulsants, *J. Gen. Physiol.* 149 (2017) 465–481, <https://doi.org/10.1085/jgp.201611668>.
- [31] K.S.P. Abelson, A.U. Höglund, Intravenously administered lidocaine in therapeutic doses increases the intraspinal release of acetylcholine in rats, *Neurosci. Lett.* 317 (2002) 93–96, [https://doi.org/10.1016/S0304-3940\(01\)02440-5](https://doi.org/10.1016/S0304-3940(01)02440-5).
- [32] H. Hiruma, K. Shimizu, T. Takenami, H. Sugie, T. Kawakami, Effects of clonidine on lidocaine-induced inhibition of axonal transport in cultured mouse dorsal root ganglion neurones, *Br. J. Anaesth.* 101 (2008) 659–665, <https://doi.org/10.1093/bja/aen265>.
- [33] L. Weinberg, Pharmacokinetics and pharmacodynamics of lignocaine: a review, *World J. Anesthesiol.* 4 (2015) 17, <https://doi.org/10.5313/wja.v4.i2.17>.
- [34] G.R. Lauretti, Mecanismo envolvidos na analgesia da lidocaína por via venosa, *Rev. Bras. Anestesiol.* 58 (2008) 280–286, <https://doi.org/10.1590/S0034-70942008000300011>.
- [35] N. Chiang, J.M. Schwab, G. Fredman, K. Kasuga, S. Gelman, C.N. Serhan, Anesthetics impact the resolution of inflammation, *PLoS One* 3 (2008), e1879, <https://doi.org/10.1371/journal.pone.0001879>.
- [36] H.C.P.M. Caracas, J.V.B. Maciel, P.M.R. e S. Martins, M.M.G. de Souza, L.C. Maia, The use of lidocaine as an anti-inflammatory substance: a systematic review, *J. Dent.* 37 (2009) 93–97, <https://doi.org/10.1016/j.jdent.2008.10.005>.
- [37] A.S. Cowbum, A.M. Condliffe, N. Farahi, C. Summers, E.R. Chilvers, Advances in Neutrophil Biology, *Chest* 134 (2008) 606–612, <https://doi.org/10.1378/chest.08-0422>.
- [38] A.M. Condliffe, E. Kitchen, E.R. Chilvers, Neutrophil priming: pathophysiological consequences and underlying mechanisms, *Clin. Sci.* 94 (1998) 461–471, <https://doi.org/10.1042/cs0940461>.
- [39] A.E. Ekpenyong, N. Toepfner, E.R. Chilvers, J. Guck, Mechanotransduction in neutrophil activation and deactivation, *Biochim. Biophys. Acta Mol. Cell Res.* 1853 (2015) 3105–3116, <https://doi.org/10.1016/j.bbamer.2015.07.015>.
- [40] K.L. Vogt, C. Summers, E.R. Chilvers, A.M. Condliffe, Priming and de-priming of neutrophil responses in vitro and in vivo, *Eur. J. Clin. Invest.* 48 (2018), e12967, <https://doi.org/10.1111/eji.12967>.
- [41] J. Svoboda, A. Popelikova, A. Stuchlik, Drugs interfering with muscarinic acetylcholine receptors and their effects on place navigation, *Front. Psychiatr.* 8 (2017), <https://doi.org/10.3389/fpsy.2017.00215>.
- [42] A.A.S. Maab, H. F. Mustafa, Anti-inflammatory aspects of Lidocaine: a neglected therapeutic stance for COVID-19, *Hear, Lung* 49 (2020) 877–878, <https://doi.org/10.1016/j.hrling.2020.09.001>.
- [43] Z.A. Ali, R.S. El-Mallakh, Nebulized lidocaine in COVID-19, an hypothesis, *Med. Hypotheses* 144 (2020) 109947, <https://doi.org/10.1016/j.mehy.2020.109947>.
- [44] M.P. Ortiz, M.C. de M. Godoy, R.S. Schlosser, R.P. Ortiz, J.P.M. Godoy, E. S. Santiago, F.K. Rigo, V. Beck, T. Duarte, M.M.M.F. Duarte, M.S. Menezes, Effect of endovenous lidocaine on analgesia and serum cytokines: double-blinded and randomized trial, *J. Clin. Anesth.* 35 (2016) 70–77, <https://doi.org/10.1016/j.jclinane.2016.07.021>.
- [45] R. Kamina, S.K. Arif, M. Hatta, A. Bukhari, R. Natzir, Hisbullah, I. Patellongi, C. Kaelan, Systemic lidocaine administration influences NF- κ B gene expression, NF- κ B and TNF- α protein levels on BALB/c mice with musculoskeletal injury, *Ann. Med. Surg.* (2021) 102660, <https://doi.org/10.1016/j.amjs.2021.102660>.
- [46] G. Li, X. Liang, M.T. Lotze, HMGB1: the central cytokine for all lymphoid cells, *front. Immunol.* 4 (2013), <https://doi.org/10.3389/fimmu.2013.00068>.
- [47] L. Huang, Y. Yao, Z. Sheng, Novel insights for high mobility group box 1 protein-mediated cellular immune response in sepsis: A systemic review, *World J. Emerg. Med.* 3 (2012) 165, <https://doi.org/10.5847/wjem.j.issn.1920-8642.2012.03.001>.
- [48] P. Scaffidi, T. Misteli, M.E. Bianchi, Release of chromatin protein HMGB1 by necrotic cells triggers inflammation, *Nature* 418 (2002) 191–195, <https://doi.org/10.1038/nature00858>.
- [49] M. Magna, D.S. Pisetsky, The role of HMGB1 in the pathogenesis of inflammatory and autoimmune diseases, *Mol. Med.* 20 (2014) 138–146, <https://doi.org/10.2119/molmed.2013.00164>.
- [50] J.S. Park, F. Gamboni-Robertson, Q. He, D. Svetkauskaite, J.-Y. Kim, D. Strassheim, J.-W. Sohn, S. Yamada, I. Maruyama, A. Banerjee, A. Ishizaka, E. Abraham, High mobility group box 1 protein interacts with multiple Toll-like receptors, *Am. J.*

- Physiol. Physiol. 290 (2006) C917–C924, <https://doi.org/10.1152/ajpcell.00401.2005>.
- [51] W.S. Yang, N.J. Han, J.J. Kim, M.J. Lee, S.-K. Park, TNF- α activates high-mobility group box 1 - toll-like receptor 4 signaling pathway in human aortic endothelial cells, *Cell. Physiol. Biochem.* 38 (2016) 2139–2151, <https://doi.org/10.1159/000445570>.
- [52] H. Wähämaa, H. Schierbeck, H.S. Hreggvidsdottir, K. Palmblad, A.-C. Aveberger, U. Andersson, H. Harris, High mobility group box protein 1 in complex with lipopolysaccharide or IL-1 promotes an increased inflammatory phenotype in synovial fibroblasts, *Arthritis Res. Ther.* 13 (2011) R136, <https://doi.org/10.1186/ar3450>.
- [53] M. Stros, HMGB proteins: interactions with DNA and chromatin, *Biochim. Biophys. Acta - Gene Regul. Mech.* 1799 (2010) 101–113, <https://doi.org/10.1016/j.bbagen.2009.09.008>.
- [54] S. Kim, S.Y. Kim, J.P. Pribis, M. Lotze, K.P. Mollen, R. Shapiro, P. Loughran, M. J. Scott, T.R. Billiar, Signaling of high mobility group box 1 (HMGB1) through toll-like receptor 4 in macrophages requires CD14, *Mol. Med.* 19 (2013) 88–98, <https://doi.org/10.2119/molmed.2012.00306>.
- [55] A. Friggeri, Y. Yang, S. Banerjee, Y.-J. Park, G. Liu, E. Abraham, HMGB1 inhibits macrophage activity in efferocytosis through binding to the α v β 3-integrin, *Am. J. Physiol. Physiol.* 299 (2010) C1267–C1276, <https://doi.org/10.1152/ajpcell.00152.2010>.
- [56] H.E. Harris, U. Andersson, D.S. Pisetsky, HMGB1: a multifunctional alarmin driving autoimmune and inflammatory disease, *Nat. Rev. Rheumatol.* 8 (2012) 195–202, <https://doi.org/10.1038/nrrheum.2011.222>.
- [57] H. Yang, D.J. Antoine, U. Andersson, K.J. Tracey, The many faces of HMGB1: molecular structure-functional activity in inflammation, apoptosis, and chemotaxis, *J. Leukoc. Biol.* 93 (2013) 865–873, <https://doi.org/10.1189/jlb.1212662>.
- [58] D. Bertheloot, E. Latz, HMGB1, IL-1 α , IL-33 and S100 proteins: dual-function alarmins, *Cell. Mol. Immunol.* 14 (2017) 43–64, <https://doi.org/10.1038/cmi.2016.34>.
- [59] R. Kang, R. Chen, Q. Zhang, W. Hou, S. Wu, L. Cao, J. Huang, Y. Yu, X. Fan, Z. Yan, X. Sun, H. Wang, Q. Wang, A. Tsung, T.R. Billiar, H.J. Zeh, M.T. Lotze, D. Tang, HMGB1 in health and disease, *Mol. Aspect. Med.* 40 (2014) 1–116, <https://doi.org/10.1016/j.mam.2014.05.001>.
- [60] T. Kawasaki, T. Kawai, Toll-like receptor signaling pathways, *Front. Immunol.* 5 (2014), <https://doi.org/10.3389/fimmu.2014.00461>.
- [61] L.A. Couture, W. Piao, L.W. Ru, S.N. Vogel, V.Y. Toshchakov, Targeting toll-like receptor (TLR) signaling by toll/interleukin-1 receptor (TIR) domain-containing adapter protein/MyD88 adapter-like (TRAP/Mal)-derived decoy peptides, *J. Biol. Chem.* 287 (2012) 24641–24648, <https://doi.org/10.1074/jbc.M112.360925>.
- [62] M. Miettinen, T. Saraneva, I. Julkunen, S. Matikainen, IFNs activate toll-like receptor gene expression in viral infections, *Gene Immun.* 2 (2001) 349–355, <https://doi.org/10.1038/sj.gene.6363791>.
- [63] M.S. Hayden, S. Ghosh, NF- κ B in immunobiology, *Cell Res.* 21 (2011) 223–244, <https://doi.org/10.1038/cr.2011.13>.
- [64] T.-h. Lin, J. Pajarinen, L. Lu, A. Nabeshima, L.A. Cordova, Z. Yao, S.B. Goodman, NF- κ B as a therapeutic target in inflammatory-associated bone diseases, in: R. Donev (Ed.), *Adv. Protein Chem. Struct. Biol.*, Elsevier Inc, 2017, pp. 117–154, <https://doi.org/10.1016/b.sapsb.2016.11.002>.
- [65] B. Banoth, B. Chatterjee, B. Vijayaragavan, M. Prasad, P. Roy, S. Basak, Stimulus-selective cross talk via the NF- κ B signaling system reinforces innate immune response to alleviate gut infection, *Elife* 4 (2015), <https://doi.org/10.7554/eLife.05648>.
- [66] M.S. Hayden, S. Ghosh, Regulation of NF- κ B by TNF family cytokines, *Semin. Immunol.* 26 (2014) 253–266, <https://doi.org/10.1016/j.smim.2014.05.004>.
- [67] N. Parameswaran, S. Patil, Tumor necrosis factor- α signaling in macrophages, *Crit. Rev. Eukaryot. Gene Expr.* 20 (2010) 87–103, <https://doi.org/10.1615/CritRevEukaryotGeneExpr.v20.i2.10>.
- [68] M.D. Turner, B. Nedjai, T. Hurst, D.J. Pennington, Cytokines and chemokines: at the crossroads of cell signalling and inflammatory disease, *Biochim. Biophys. Acta Mol. Cell Res.* 1843 (2014) 2563–2582, <https://doi.org/10.1016/j.bbamcr.2014.05.014>.
- [69] S. Kany, J.T. Vollrath, B. Relja, Cytokines in inflammatory disease, *Int. J. Mol. Sci.* 20 (2019) 6008, <https://doi.org/10.3390/ijms20236008>.
- [70] E. Ricciotti, G.A. FitzGerald, Prostaglandins and inflammation, *Arterioscler. Thromb. Vasc. Biol.* 31 (2011) 986–1000, <https://doi.org/10.1161/ATVBAHA.110.207449>.
- [71] T. Liu, L. Zhang, D. Joo, S.-C. Sun, NF- κ B signaling in inflammation, *Signal Transduct. Target. Ther.* 2 (2017) 17023, <https://doi.org/10.1038/sigtrans.2017.23>.
- [72] J.-H. Shi, S.-C. Sun, Tumor necrosis factor receptor-associated factor regulation of nuclear factor κ B and mitogen-activated protein kinase pathways, *Front. Immunol.* 9 (2018), <https://doi.org/10.3389/fimmu.2018.01849>.
- [73] Y.L. Pobezinskaya, Z. Liu, The role of TRADD in death receptor signaling, *Cell Cycle* 11 (2012) 871–876, <https://doi.org/10.4161/cc.11.5.19300>.
- [74] J.R. Bradley, J.S. Pober, Tumor necrosis factor receptor-associated factors (TRAFs), *Oncogene* 20 (2001) 6482–6491, <https://doi.org/10.1038/sj.onc.1204788>.
- [75] L. Vigneault, A.F. Turgeon, D. Côté, F. Lauzier, R. Zarychanski, L. Moore, L. A. McIntyre, P.C. Nicole, D.A. Fergusson, Perioperative intravenous lidocaine infusion for postoperative pain control: a meta-analysis of randomized controlled trials, *Can. J. Anesth. Can. d'anesthésie.* 58 (2011) 22–37, <https://doi.org/10.1007/s12630-010-9407-0>.
- [76] S.-C. Sun, Non-canonical NF- κ B signaling pathway, *Cell Res.* 21 (2011) 71–85, <https://doi.org/10.1038/cr.2010.177>.
- [77] P. Kranke, J. Jokinen, N.L. Pace, A. Schnabel, M.W. Hollmann, K. Hahnenkamp, L. H. Eberhart, D.M. Poepping, S. Weibel, Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery, *Cochrane Database Syst. Rev.* (2015), <https://doi.org/10.1002/14651858.CD009642.pub2>.
- [78] S. Weibel, Y. Jeltting, N.L. Pace, A. Helf, L.H. Eberhart, K. Hahnenkamp, M. W. Hollmann, D.M. Poepping, A. Schnabel, P. Kranke, Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery in adults, *Cochrane Database Syst. Rev.* (2018), <https://doi.org/10.1002/14651858.CD009642.pub3>.
- [79] M. Bailey, T. Corcoran, S. Schug, A. Toner, Perioperative Lidocaine Infusions for the Prevention of Chronic Postsurgical Pain, 2018, p. 1, <https://doi.org/10.1097/j.pain.0000000000001273>. Pain.
- [80] N.T. Ventham, E.D. Kennedy, R.R. Brady, H.M. Paterson, D. Speake, I. Foo, K.C. H. Fearon, Efficacy of intravenous lidocaine for postoperative analgesia following laparoscopic surgery: a meta-analysis, *World J. Surg.* 39 (2015) 2220–2234, <https://doi.org/10.1007/s00268-015-3105-6>.
- [81] Y. Sun, T. Li, N. Wang, Y. Yun, T.J. Gan, Perioperative systemic lidocaine for postoperative analgesia and recovery after abdominal surgery, *Dis. Colon Rectum* 55 (2012) 1183–1194, <https://doi.org/10.1097/DCR.0b013e318259bcd8>.
- [82] G.C. McCarthy, S.A. Megalla, A.S. Habib, Impact of intravenous lidocaine infusion on postoperative analgesia and recovery from surgery, *Drugs* 70 (2010) 1149–1163, <https://doi.org/10.2165/00000000000000000000>.
- [83] K. Hirota, H. Okawa, B.L. Appadu, D.K. Grandy, D.G. Lambert, Interaction of local anaesthetics with recombinant μ , κ , and δ -opioid receptors expressed in Chinese hamster ovary cells, *Br. J. Anaesth.* 85 (2000) 740–746, <https://doi.org/10.1093/bja/85.5.740>.
- [84] G.W. Pasternak, Y.-X. Pan, Mu opioids and their receptors: evolution of a concept, *Pharmacol. Rev.* 65 (2013) 1257–1317, <https://doi.org/10.1124/pr.112.007138>.
- [85] J. Liu, H. Zhang, Z. Qi, X. Zheng, Lidocaine protects against renal and hepatic dysfunction in septic rats via downregulation of Toll-like receptor 4, *Mol. Med. Rep.* 9 (2014) 118–124, <https://doi.org/10.3892/mmr.2013.1799>.
- [86] L. Wang, M. Wang, S. Li, H. Wu, Q. Shen, S. Zhang, L. Fang, R. Liu, Nebulized lidocaine ameliorates allergic airway inflammation via downregulation of TLR2, *Mol. Immunol.* 97 (2018) 94–100, <https://doi.org/10.1016/j.molimm.2018.03.010>.
- [87] J. Frieden, Antiarrhythmic drugs. Part VII. Lidocaine as an antiarrhythmic agent, *Am. Heart J.* 70 (1965) 713–715, [https://doi.org/10.1016/0002-8703\(65\)90399-6](https://doi.org/10.1016/0002-8703(65)90399-6).
- [88] F.J. Dowd, Antiarrhythmic drugs. Pharmacol. Ther. 39:3, Elsevier, 2017, pp. 287–300, <https://doi.org/10.1016/b978-0-323-39007-2.00019-9>.
- [89] A. Mizzi, T. Tran, D. Mangar, E.M. Camporesi, Amiodarone supplants lidocaine in ACLS and CPR protocols, *Anesthesiol. Clin.* 29 (2011) 535–545, <https://doi.org/10.1016/j.anclin.2011.05.001>.
- [90] Y. Chen, Q. Liu, D. Guo, Emerging coronaviruses: genome structure, replication, and pathogenesis, *J. Med. Virol.* 92 (2020) 418–423, <https://doi.org/10.1002/jmv.25681>.
- [91] S.L. Kong, P. Chui, B. Lim, M. Salto-Tellez, Elucidating the molecular physiopathology of acute respiratory distress syndrome in severe acute respiratory syndrome patients, *Virus Res.* 145 (2009) 260–269, <https://doi.org/10.1016/j.virusres.2009.07.014>.
- [92] V.J. Costela-Ruiz, R. Illescas-Montes, J.M. Puerta-Puerta, C. Ruiz, L. Melguizo-Rodríguez, SARS-CoV-2 infection: the role of cytokines in COVID-19 disease, *Cytokine Growth Factor Rev.* <https://doi.org/10.1016/j.cytogfr.2020.06.001>, 2020, 54, 62, 75.
- [93] D. Darif, I. Hammi, A. Kihel, I. El Idrissi Saik, F. Guessous, K. Akarid, The pro-inflammatory cytokines in COVID-19 pathogenesis: what goes wrong? *Microb. Pathog.* 153 (2021) 104799, <https://doi.org/10.1016/j.micpath.2021.104799>.
- [94] P.G.S.G.S. Papathanasiou, Local anesthetics and Covid19 associated acute respiratory distress syndrome: A New therapeutic indication? *Clin. Res. Open Access.* 6 (2020) <https://doi.org/10.16966/2469-6714.156>.
- [95] Y.-R. Guo, Q.-D. Cao, Z.-S. Hong, Y.-Y. Tan, S.-D. Chen, H.-J. Jin, K.-S. Tan, D.-Y. Wang, Y. Yan, The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak – an update on the status, *Mil. Med. Res.* 7 (2020) 11, <https://doi.org/10.1186/s40779-020-00240-0>.
- [96] D. Wang, B. Hu, C. Hu, F. Zhu, X. Liu, J. Zhang, B. Wang, H. Xiang, Z. Cheng, Y. Xiong, Y. Zhao, Y. Li, X. Wang, Z. Peng, Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in wuhan, China, *J. Am. Med. Assoc.* 323 (2020) 1061, <https://doi.org/10.1001/jama.2020.1585>.
- [97] M.A. Matthay, R.L. Zemans, The acute respiratory distress syndrome: pathogenesis and treatment, *Annu. Rev. Pathol.* 6 (2011) 147–163, <https://doi.org/10.1146/annurev-pathol-011110-130158>.
- [98] T. Piegeler, E.G. Votta-Velis, F.R. Bakhshi, M. Mao, G. Carnegie, M.G. Bonini, D. E. Schwartz, A. Borgeat, B. Beck-Schimmer, R.D. Minshall, Endothelial barrier protection by local anesthetics, *Anesthesiology* 120 (2014) 1414–1428, <https://doi.org/10.1097/ALN.000000000000174>.

Molecular mechanisms of lidocaine

ORIGINALITY REPORT

35%

SIMILARITY INDEX

22%

INTERNET SOURCES

28%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Henning Hermanns, Markus W. Hollmann, Markus F. Stevens, Philipp Lirk, Timo Brandenburger, Tobias Piegeler, Robert Werdehausen. "Molecular mechanisms of action of systemic lidocaine in acute and chronic pain: a narrative review", *British Journal of Anaesthesia*, 2019
Publication 5%
- 2 neuro.unboundmedicine.com
Internet Source 2%
- 3 www.annualreviews.org
Internet Source 2%
- 4 molmed.biomedcentral.com
Internet Source 2%
- 5 Julian Frieden. "Antiarrhythmic drugs. Part VII. Lidocaine as an antiarrhythmic agent", *American Heart Journal*, 1965
Publication 2%
- 6 Andrew E. Ekpenyong, Nicole Toepfner, Edwin R. Chilvers, Jochen Guck. 2%

"Mechanotransduction in neutrophil activation and deactivation", Biochimica et Biophysica Acta (BBA) - Molecular Cell Research, 2015

Publication

7

Chung-Hsi Hsing, Jhi-Joung Wang. "Clinical implication of perioperative inflammatory cytokine alteration", Acta Anaesthesiologica Taiwanica, 2015

Publication

1 %

8

ecommons.aku.edu

Internet Source

1 %

9

Hira Maab, Faryal Mustafa, Shajeea Arshad Ali. "Anti-inflammatory aspects of Lidocaine: a neglected therapeutic stance for COVID-19", Heart & Lung, 2020

Publication

1 %

10

coek.info

Internet Source

1 %

11

sciforschenonline.org

Internet Source

1 %

12

Ziad A. Ali, Rif S. El-Mallakh. "Nebulized Lidocaine in COVID-19, An Hypothesis", Medical Hypotheses, 2020

Publication

1 %

13

www.ncbi.nlm.nih.gov

Internet Source

1 %

14	derangedphysiology.com Internet Source	1 %
15	Submitted to University of Arizona Student Paper	1 %
16	link.springer.com Internet Source	1 %
17	Submitted to National University Student Paper	1 %
18	Andersson, Ulf, and Kevin J. Tracey. "HMGB1 Is a Therapeutic Target for Sterile Inflammation and Infection", Annual Review of Immunology, 2011. Publication	1 %
19	pubs.asahq.org Internet Source	1 %
20	Submitted to University of Technology, Sydney Student Paper	<1 %
21	Jia Chen, Lanwen Hu, Rui Jiang. "A commentary on "Updated evaluation of endoscopic submucosal dissection versus surgery for early gastric cancer: A systematic review and meta-analysis" (Int J Surg 2020; 73:28-41)", International Journal of Surgery, 2021 Publication	<1 %

22 Ortiz, Michele Purper, Maria Celoni de Mello Godoy, Rochelle Silveira Schlosser, Rafael Purper Ortiz, João Pedro Mello Godoy, Eduardo Sagrillo Santiago, Flávia Karine Rigo, Verônica Beck, Thiago Duarte, Marta Maria Medeiros Frescura Duarte, and Miriam Seligman Menezes. "Effect of endovenous lidocaine on analgesia and serum cytokines: double-blinded and randomized trial", Journal of Clinical Anesthesia, 2016.
Publication

23 Submitted to Texas State Technical College
Student Paper

24 Submitted to Herzing University
Student Paper

25 journals.plos.org
Internet Source

26 www.hindawi.com
Internet Source

27 academy.esraeurope.org
Internet Source

28 doku.pub
Internet Source

29 worldwidescience.org
Internet Source

30	Submitted to Associatie K.U.Leuven Student Paper	<1 %
31	www.frontiersin.org Internet Source	<1 %
32	Hayden, Matthew S., and Sankar Ghosh. "Regulation of NF-κB by TNF family cytokines", Seminars in Immunology, 2014. Publication	<1 %
33	Yang, H.. "Targeting HMGB1 in inflammation", BBA - Gene Regulatory Mechanisms, 201001/02 Publication	<1 %
34	bmchealthservres.biomedcentral.com Internet Source	<1 %
35	koreascience.or.kr Internet Source	<1 %
36	onlinelibrary.wiley.com Internet Source	<1 %
37	courses.lumenlearning.com Internet Source	<1 %
38	A. Taylor, G. McLeod. "Basic pharmacology of local anaesthetics", BJA Education, 2020 Publication	<1 %
39	Huiling Wu, Renske Steenstra, Elianne C.S. de Boer, Cathy Y. Zhao, Jin Ma, Jorieke M. van der	<1 %

Stelt, Steven J. Chadban. "Preconditioning with recombinant high-mobility group box 1 protein protects the kidney against ischemia-reperfusion injury in mice", *Kidney International*, 2014

Publication

40

academic.oup.com

Internet Source

<1 %

41

annalsofintensivecare.springeropen.com

Internet Source

<1 %

42

dirjournal.org

Internet Source

<1 %

43

Chandra Mohan, Chaim Putterman. "Genetics and pathogenesis of systemic lupus erythematosus and lupus nephritis", *Nature Reviews Nephrology*, 2015

Publication

<1 %

44

Zhou, Guang-Yao, Yong-Xiang Yi, Ling-Xiang Jin, Wei Lin, Pei-Pei Fang, Xiu-Zheng Lin, Yi Zheng, and Chen-Wei Pan. "The protective effect of juglanin on fructose-induced hepatitis by inhibiting inflammation and apoptosis through TLR4 and JAK2/STAT3 signaling pathways in fructose-fed rats", *Biomedicine & Pharmacotherapy*, 2016.

Publication

<1 %

45

aepi.biomedcentral.com

Internet Source

<1 %

46

downloads.hindawi.com

Internet Source

<1 %

47

journals.lww.com

Internet Source

<1 %

48

publications.aston.ac.uk

Internet Source

<1 %

49

repository.ubn.ru.nl

Internet Source

<1 %

50

www.biotek.es

Internet Source

<1 %

51

www.freepatentsonline.com

Internet Source

<1 %

52

www.karger.com

Internet Source

<1 %

53

www.scielo.br

Internet Source

<1 %

54

www.tandfonline.com

Internet Source

<1 %

55

Bernhard Englinger, Christine Pirker, Petra Heffeter, Alessio Terenzi, Christian R. Kowol, Bernhard K. Keppler, Walter Berger. "Metal

<1 %

Drugs and the Anticancer Immune Response", Chemical Reviews, 2018

Publication

56

Ray, Rashmi, Judyta K. Juranek, and Vivek Rai.
"RAGE axis in neuroinflammation,
neurodegeneration and its emerging role in
the pathogenesis of amyotrophic lateral
sclerosis", Neuroscience & Biobehavioral
Reviews, 2016.

Publication

57

lawancovid19.tangerangselatankota.go.id

Internet Source

58

Junying Ding, Xuran Cui, Qingquan Liu.
"Emerging role of HMGB1 in lung diseases:
friend or foe", Journal of Cellular and
Molecular Medicine, 2017

Publication

59

Mirhan N. Makled, Mohammed S. El-Awady,
Rania R. Abdelaziz, Nadia Atwan et al.
"Pomegranate protects liver against cecal
ligation and puncture-induced oxidative stress
and inflammation in rats through TLR 4 /NF -
κB pathway inhibition", Environmental
Toxicology and Pharmacology, 2016

Publication

<1 %

<1 %

<1 %

<1 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 5 words