

Dextrose 10% drink is superior to sodium-dextrose drink in increasing blood glucose and sprint speed in soccer players: A double-blinded randomized crossover trial study

by

Submission date: 13-Jan-2022 03:50AM (UTC+0700)

Submission ID: 1740787454

File name: 10._Agussalim_Bukhari-Chris.pdf (936.85K)

Word count: 7657

Character count: 39460

MOTS CLÉS

Effet ergogénique ;
Aide ergogénique ;
Footballeurs masculins ;
Glycémie ;
Sprint vitesse ;
VO₂max ;
Dextrose

Results. – The mean BG level was higher in D compared with Sodium-D group [136 ± 22.9 mg/dl vs. 118 ± 21.5 mg/dl] ($P=0.009$). Mean differences of Δ BG of D (mean before and mean after) compared with Δ BG Sodium-D (mean before and mean after) is 16.8 mg/dl ($P=0.44$). The mean of sprint speed is faster in the D group [15.2 ± 1.25 sec vs. 15.9 ± 1.61 sec] ($P=0.019$), but we found no significant differences in VO₂max between both groups [42.1 ± 3.44 ml/min/kg vs. 42.3 ± 3.98 ml/min/kg] ($P=0.834$).

Conclusion. – The present study demonstrated that dextrose drink without added sodium has a large influence on increasing blood glucose level and sprint speed.

© 2021 Elsevier Masson SAS. All rights reserved.

Résumé

Objectifs. – Cette étude visait à examiner l'effet d'une boisson à 10 % de dextrose sur la glycémie, la VO₂max et la vitesse de sprint par rapport à une boisson Sodium-Dextrose.

Équipement et méthodes: Nous avons mené une étude croisée en double insu sur 22 jeunes footballeurs de sexe masculin ($19 \pm 1,1$ ans), consommant une boisson au Dextrose ou une boisson Sodium-Dextrose. Nous avons comparé les données pré- et post- sur la glycémie, la VO₂max et la vitesse de sprint entre les groupes de boissons Dextrose et Sodium-Dextrose. Le critère de jugement principal était les différences de glycémie Δ entre les groupes Dextrose et Sodium-Dextrose. Les critères de jugement secondaires étaient des différences de vitesse Δ VO₂max et Δ sprint entre les groupes Dextrose et Sodium-Dextrose.

Résultats. – Le niveau moyen de glycémie était plus élevé dans D par rapport au groupe Sodium Dextrose [$136 \pm 22,9$ mg/dl vs. $118 \pm 21,5$ mg/dl] ($p=0,009$). Les différences moyennes de Δ BG de Dextrose 10 % (moyenne avant et moyenne après) par rapport à Δ BG Sodium-D (moyenne avant et moyenne après) sont de $16,8$ mg/dl ($p=0,001$). La vitesse moyen sprint est plus rapide dans le groupe D [$15,2 \pm 1,25$ s vs. $15,9 \pm 1,61$ s] ($p=0,019$), mais nous n'avons trouvé aucune différence significative de VO₂max entre les deux groupes [$42,1 \pm 3,44$ ml/min/43 kg vs. $42,3 \pm 3,98$ ml/min/kg] ($p=0,834$).

Conclusion. – La présente étude a démontré que la boisson de dextrose sans sodium ajouté a un effet portant sur l'augmentation du taux de glycémie et de la vitesse de sprint.

© 2021 Elsevier Masson SAS. Tous droits réservés.

1. Introduction

Carbohydrate (CHO) ingestion has been known to improve skilled and cognitive performance, and rapid reduction in Blood Glucose (BG) could harmfully affect the skilled actions in a soccer match [1]. BG decreased sharply with the onset of soccer [2] due to the augmentation of glucose uptake via translocation of glucose transporters type 4 (GLUT4) to the active site of muscle membrane [3].

Two groups of glucose transporters are responsible for facilitating glucose passage into and out of the cells: the sodium-glucose co-transporters (SGLTs) and the facilitative glucose transporters (GLUTs) [4]. SGLT1 and GLUT5 are localized in the brush border membrane (BBM) of enterocytes; GLUT2 in the basolateral membrane (BLM) at low luminal dextrose concentration, but in the BBM as BLM at high dextrose concentrations. Hence, dextrose absorption at low luminal concentration is mediated via SGLT1 in the BBM and GLUT2 in the BLM, but in high concentration, it is also enhanced by GLUT2 in the BBM [5].

Enterocyte has a limited capacity to absorb glucose [6]. Exercise studies have shown that CHO's delivery is limited by the transport capacity of SGLT-1 [7]. High intestinal luminal glucose concentrations ≥ 15 mM, or more modest loads, induces expression of GLUT-2 in BBM [8,9]. Glucose absorption can be amplified by BBM GLUT-2 when the enterocyte and submucosal glucose concentration are lower than in the

lumen [10]. The time required to reach steady-state glucose accumulation within the enterocytes in vitro following exposure is ≤ 2 min [11,12] and within 5 to 10 min [13], and a soccer player must achieve it as soon as possible. A lower concentration of luminal glucose than the submucosal has been reported [14], and it will hinder apical GLUT-2 for passive glucose absorption rather than assisted net absorption [10,15].

Outfield soccer players covered 9-12 km of distance during a match; 12% are sprinting [16]. Hence, soccer players require rapid energy turnover from both aerobic and anaerobic metabolisms [17]. The ergogenic effects of CHO ingestion were associated with the maintenance of glycogen levels because of a 30–50 times increment of muscle glucose uptake in exercising muscle [18]. As a result, CHO's availability was shown to influence soccer performance by enhancing the capacity to perform the intermittent activity and may become a limiting performance factor when it is depleted [19].

Sports drinks are designed to replenish CHO, electrolytes, and fluids to the body, which required fast absorption in the BBM [20]. Generally, sports drinks contain either CHO and or sodium-CHO, but to know which one is better in terms of ergogenic effect (BG concentration, VO₂ max, and sprint speed), there is still no trial that has been established. Therefore, we designed a randomized, double-blind trial in male soccer players to evaluate the effect of dextrose

drink with no sodium versus the dextrose drink with 20 mM sodium in blood glucose, VO_2 max, and sprint speed. The primary outcome was a mean difference in BG level between Dextrose 10% (D) and Sodium-Dextrose 10% (Na-D) intervention. Secondary outcomes were mean differences in VO_2 max and sprint speed. We hypothesized that D supplementation gives a superior effect than Na-D supplementation in blood glucose, VO_2 max, and sprint speed.

39 2. Methods

2.1. Study design and procedure

25
This study was a randomized crossover (double-blind) design. Participants and researchers were blinded from the administered fluid type. Participants were randomly assigned to receive either dextrose (D) solution or sodium dextrose (Na-D) solution. Randomization took place immediately after the participants had been confirmed to be eligible to enter the study. The drink's randomization code was not made available to anyone involved in conducting or evaluating the study and was released after the blind review and the freezing of the final database. After randomizing, the study was divided into three phases, as shown in Fig. 1. Assessment of health status, dietary intake, body composition measurement, baseline blood glucose, baseline VO_2 max, and baseline sprint speed was evaluated in phase I. We measured the blood glucose before the participants drink the dextrose or sodium dextrose supplementation in phase II. We asked the participants to sprint before received the solution, then we recorded the time and calculated the VO_2 max. After that, we gave one type of solution (dextrose or sodium dextrose). In order to have a comparison of the post supplementation effect, the participants repeated the previously mentioned procedures in phase II. Phase III was similar to phase II except for the nutritional supplementation. The participants could receive either dextrose or sodium dextrose in phase II and the opposite alternative in phase III. This trial was conducted at UNM Banta-Bantaeng, Makassar, Indonesia, from April 2019 to May 2019.

2.2. Supplement preparation

The supplements consisted of dextrose (D) or sodium dextrose (Na-D). Glucose is a monosaccharide, which acts as an essential energy source through aerobic or anaerobic metabolism. Dextrose or D-glucose is an aldohexose stereoisomer of glucose. It is the most commonly occurring isomer of glucose in nature. Our pharmacist used 15 g of dextrose anhydrous to make a 150 ml dextrose 10%, with pH 5.27. Another 150 ml Sodium dextrose solution consisted of 10% dextrose, added with 20 mM of sodium. The material is taken from Japan (Otsuka Holdings Co., Ltd.), and Akademis hospital's licensed pharmacist supervised the entire mixing process.

2.3. Participants

Sample sizes for this research are twenty one, and it was calculated by equation $n = \left[\frac{(Z_{\alpha} + Z_{\beta}) \times Sd}{d} \right]^2$, with

$Z_{\alpha} = 1.96$; $Z_{\beta} = 0.842$; $Sd = 28.2$; $d = 10$. Random allocation was performed using a random electronic generator via <https://www.random.org/lists/>. Our pharmacist staffs were responsible for generating the random allocation sequence, enrolled participants, and assigned participants to interventions. After the assignment of interventions, all of the participants and the researchers were blinded. A total of 50 male outfield academy soccer players were screened for the study. Thirty-two participants entered into the inclusion criteria. All were healthy and free of injury in the three months preceding the study. Three participants refused to join, and 6 were dropped out. Twenty-two participants participated (age, 19.6 ± 1.1 years; height, 165.7 ± 5.25 cm; weight, 52.5 ± 8.47 kg; body fat, $12.5 \pm 4.68\%$; muscle mass, $82 \pm 4.7\%$; BMR, 1346 ± 147.0 kcal) in the study and were successfully analyzed until the end. Before enrolling in the study, each participants' physical examination and health history were taken. Eligibility criteria were healthy men soccer players between 18 and 23 years of age, who exercised on average between 8 and 12 h per week the last month before inclusion. All participants were within the last meal four hours before the test. Exclusion criteria were the use of amylase supplement, suffering from fever and diarrhea, using laxative agents within 24h, consuming CHO absorption inhibitors, caffeine, creatinine, beta-alanine, sodium bicarbonate supplement within 24h, mean arterial pressure <65 mmHg, knee or muscle injuries, history of diabetes mellitus and heart disease, going through the ketogenic diet program, history of gastrointestinal surgery, and total body fat percentages > 30%. Participants who were currently taking any other dietary supplement, sports drink, or functional food intended to enhance performance or muscle mass or had taken any of these in the previous week were also excluded.

The study was approved by the Faculty of Medicine Hasanuddin University Research Ethics Committee with reference number 214/UN4.6.4.5.31/PP36/2019. Written informed consent was obtained from all participants before inclusion. The study was registered at ClinicalTrials.gov, with reference number NCT04206579.

2.4. Intervention and procedures

The participants were instructed to abstain from exercise 24h before the testing in phase I, II, and III, and they arrived at the field track by car, motorcycle, or by public transportation. They were recommended to maintain the usual eating pattern 48h before the test and had the last meal 4hours before phase I. Besides, they were told to refrain from coffee, and sports drink 48h before the test.

Dietary intake was collected using two days of a food recall. BG was measured from capillary blood glucose from the fingertip and immediately analyzed (Aviva; Accucheck, Roche Diagnostics, Indiana, USA), blood pressure was measured using an aneroid sphygmomanometer (R1 shock-proof; Riester, Jungingen, Germany), heart rate was measured with wrist band pulse monitor (Bluetooth 4.0 wireless sport heart rate monitor WP290; Egoman, Shenzhen, China), body weight, muscle, fat, water, metabolic rate was measured using the body composition analyzer (BC-545N; Tanita,




Figure 1 Flowchart of the study.

Tokyo, Japan), body height was measured using a stadiometer (HR-200, Tanita, Tokyo, Japan), sprint speed was measured using a digital stopwatch (sec23589 sec23589P1; Seiko, Tokyo, Japan).

Participants were instructed to refrain from strenuous physical activity in the 2 days preceding trial sessions and recorded all food consumed in the 2 days before the trial. Food records subsequently were analyzed using professional German nutrition software (EBISpro, Nutrisurvey 2007). On arrival at the field, pre supplementation capillary blood samples were collected, and then all players run for 2×100 m and calculated the VO_2 max using Uth-Sorenen-Overgaard-Pedersen formula, and sprint speed was recorded. After doing the baseline measurement, each player waited for 15 minutes to consume either dextrose or sodium dextrose solution and then waited for 15 minutes to have another subsequent capillary blood sample measurement. After that, the players run for 2×100 m, recorded the VO_2 max and sprint speed. The players remained in a rested state for 120 minutes as a crossover washed-out period and then did the same protocol with a different solution.

2.5. Statistics

Descriptive statistics were used to characterize the participants (mean, standard deviation [SD]), with a 95% confidence interval, and significance was accepted at $P < 0.05$. Data were checked for normality, as indicated by the Shapiro–Wilk test. Paired samples *t*-tests were used for comparison before and after the condition in blood sugar, VO_2 max, and sprint speed. All reported *P*-values were based on two-sided tests. Exact values of 0.05 were considered statistically significant. The data were analyzed using IBM SPSS Statistics 25 for windows (SPSS Inc., Chicago, Illinois, USA). The sample size was selected based on a priori power calculation. Accordingly, with a type I error of 0.05, power analysis revealed that for a 2-tailed paired data Student *t*-test, a sample size of 21 individuals is adequate to achieve a power of > 80%. To interpret the magnitude of the effect, Cohen's *d* effect sizes (\pm 95% confidence limits) were estimated using purpose-built spreadsheet, with effect size thresholds set at < 0.20, > 0.50, and > 0.80 for small, moderate, large, effects respectively.

29

Table 1 Baseline Characteristics of the study participants ($n=22$).

Characteristic	Mean \pm SD
Age (years)	19.6 \pm 1.12
Height (cm)	165.7 \pm 5.25
Weight (kg)	52.7 \pm 8.47
Systolic blood pressure (mmHg)	109 \pm 5.4
Diastolic blood pressure (mmHg)	79 \pm 9.5
Body fat (%)	12.5 \pm 4.68
Muscle mass (%)	82 \pm 4.7
Bone mass (kg)	2.5 \pm 0.31
BMR (kcal)	1346 \pm 147.0
Visceral fat level	2.9 \pm 1.0
Total body water (%)	60.7 \pm 4.91
Blood Glucose (mg/dl)	93 \pm 2.5
VO ₂ max (ml/min/kg)	41.0 \pm 3.21
Sprint speed (sec)	16.13 \pm 1.333
Heart rate (times/minute)	60 \pm 4.1
Protein Intake (g) ^a	98.4 \pm 16.23
Carbohydrate Intake(g) ^a	327.4 \pm 41.28
Fat intake (g) ^a	48.1 \pm 14.68
Energy intake (kcal) ^a	2137 \pm 295.8

Data are presented as mean \pm standard deviation (SD) unless otherwise stated.

^a Average of 48h food recall.

3. Results

The baseline characteristic for all 22 participants was shown in Table 1. All of the variables are within a normal distribution. All participants ($n=22$) completed the test without any injuries, and none of the participants in both groups reported any side effects. Due to the little differences in the muscle ($82 \pm 4.7\%$) and total body water percentages ($60.7 \pm 4.91\%$) of all participants, we used similar dextrose and sodium dextrose doses.

The effect of dextrose or sodium dextrose supplementation on blood glucose, VO₂max, and sprint are presented in Figs. 2–4, respectively. The mean of blood glucose before and after dextrose supplementation is 91 ± 11.6 mg/dl and 136 ± 22.9 mg/dl. The mean of blood glucose before and after sodium dextrose supplementation is 91 ± 9.1 mg/dl and 118 ± 21.5 mg/dl. Dextrose supplementation had a higher blood glucose level than sodium dextrose (136 mg/dl versus 118 mg/dl). This data was also supported by BG's mean differences before dextrose supplementation compared with before sodium dextrose was 0.4 ± 5.86 mg/dl, $p=0.747$ (95% CI $-2.19, 3.01$), which means their differences were not statistically significant. The data between the dextrose and sodium dextrose group was shown in Tables 2 and 3. Another corroboration of the data was found that BG's mean differences after dextrose supplementation compared with after sodium dextrose was 17 ± 28.2 mg/dl, $p=0.009$ (95% CI $4.7-29.8$). Cohen's d effect's size = 0.60. However, to sum it all, the primary endpoint is to know the mean differences of Δ BG in the dextrose group (before and after supplementation) compared with a Δ BG sodium dextrose group (before and after supplementation) as shown in




Figure 2 Blood glucose differences. Blood glucose differences before and after dextrose supplementation ($P=0.001$). Blood glucose differences before and after sodium dextrose supplementation ($P=0.001$). Shading bar: blood glucose before supplementation; white bar: blood glucose after supplementation




Figure 3 VO₂max differences. VO₂max differences before and after dextrose supplementation ($P=0.282$). VO₂max differences before and after sodium dextrose supplementation ($P=0.219$). Shading bar: VO₂max before supplementation; white bar: VO₂max after supplementation

Table 3, which has the P -value of 0.011 and effect's sizes 0.6.

The mean of VO₂max before and after dextrose supplementation is 42.7 ± 3.69 ml/min/kg and 42.1 ± 3.44 ml/min/kg. The mean of VO₂max before and after sodium dextrose supplementation are 42.5 ± 3.98 ml/min/kg and 42.3 ± 3.99 ml/min/kg. The mean differences of VO₂max before dextrose supplementation compared with before sodium dextrose was 0.2 ± 4.57 ml/min/kg, $p=0.810$ (95% CI $-1.79, 2.26$). The mean differences of VO₂max after dextrose supplementation compared with after sodium dextrose was -0.16 ± 3.65 ml/min/kg, $p=0.834$ (95% CI $-1.78, 1.45$). Cohen's d effect's size = 0.04. The secondary endpoint is to know the mean differences of the Δ VO₂max dextrose

Table 2 Before and after administration of dextrose and sodium dextrose.

	Dextrose				Sodium dextrose			
	n = 22				n = 22			
	Before	After	P-value	Cohen's d effect sizes	Before	After	P-value	Cohen's d effect sizes
Mean \pm SD	Mean \pm SD	Mean \pm SD			Mean \pm SD			
BG (mg/dl)	91.4 \pm 11.62	136.0 \pm 22.91	0.001	3.02	91.0 \pm 9.18	118.7 \pm 21.53	0.001	1.29
Sprint (sec)	14.69 \pm 1.330	15.26 \pm 1.258	0.012	0.59	14.61 \pm 1.468	15.90 \pm 1.612	0.001	1.23
VO ₂ max	42.7 \pm 3.69	42.1 \pm 3.44	0.282	0.23	42.5 \pm 3.98	42.3 \pm 3.99	0.219	0.27
HR (bpm) ^a	148 \pm 3.04	149 \pm 3.26	0.204	0.27	148 \pm 2.78	148 \pm 3.41	0.590	0.11

Data are presented as mean values \pm standard deviations (SD). The P-value represents the significance level of the difference between the before dextrose or sodium dextrose after the dextrose or sodium dextrose supplementation. Cohen's d represents the effect sizes of the p-value. BG, blood glucose; VO₂max, maximal oxygen uptake; HR, heart rate; BPM, beat per minute.

^a Measured at the end of the sprint sessions.

Table 3 Differences of blood glucose, sprint and VO₂max before and after dextrose or sodium dextrose supplementation.

	Dextrose	Sodium Dextrose	Diff. Dextrose versus Sodium Dextrose			
	n = 22	n = 22				
	Mean diff \pm SD	Mean diff \pm SD	Mean diff	95% CI	P-value	Cohen's d effect sizes
BG (mg/dl)	-44.5 \pm 14.76	-27.7 \pm 21.33	16.8	4.27, 29.44	0.011	0.6
Sprint (sec)	-0.57 \pm 0.966	-1.28 \pm 1.035	-0.71	-1.24, -0.18	0.011	0.6
VO ₂ max	0.5 \pm 2.46	0.1 \pm 0.65	-0.40	-1.64, 0.83	0.507	0.1

Data are presented as mean values, standard deviations (SD), 95% confidence interval (CI), and P-value. Diff. dextrose versus sodium dextrose: differences between before and after supplementation of dextrose versus sodium dextrose. BG: blood glucose; VO₂max: maximal oxygen uptake.

group (before and after supplementation) compared with a Δ VO₂max sodium dextrose group (before and after supplementation), as shown in Table 3, which has a P-value of 0.507 and effect's sizes 0.1, as shown in Table 3.

Before and after dextrose supplementation, the mean sprint speed is 14.6 \pm 1.33sec and 15.2 \pm 1.25sec. The mean sprint speed before and after sodium dextrose supplementation is 14.6 \pm 1.46sec and 15.9 \pm 1.61sec. The mean differences of sprint speed before dextrose supplementation compared with before sodium dextrose were 0.07 \pm 0.724sec, $p=0.640$ (95% CI -0.24, 0.39). The mean differences of sprint speed after dextrose supplementation compared with after sodium dextrose was -0.64 \pm 1.184sec, $p=0.019$ (95% CI -1.16, -0.11). Cohen's d effect's size = 0.55. The mean sprint speed is faster in a dextrose supplementation group than the sodium dextrose group (15.2sec versus 15.9sec). The secondary endpoint of this study supported this value. The mean differences of Δ sprint speed in the dextrose group (before and after

supplementation) compared with the Δ sprint speed in sodium dextrose group (before and after supplementation) as shown in Table 3, which has a P-value of 0.011 and effect's sizes 0.6, as shown in Table 3.

4. Discussion

The mean differences of BG after dextrose versus after sodium dextrose group were 136 \pm 22.9mg/dl and 118 \pm 21.5mg/dl. This positive finding may suggest the mechanism that solely dextrose supplementation via a GLUT-2 was able to transport CHO into the plasma faster than SGLT-1. CHO can be moved through the BBM via a GLUT-2 pathway [15], which may occur if the CHO supplementation is given without sodium as a co-transporter, evading the SGLT-1 pathway [21].

The total CHO concentration in sports drinks is 4–8%, and the sodium content is 10–25mM [22]. In this study, we were




Figure 4 Sprint differences. Sprint differences before and after dextrose supplementation ($P=0.012$). Sprint differences before and after sodium dextrose supplementation ($P=0.001$). Shading bar: sprint speed before supplementation; white bar: sprint speed after supplementation

using a higher CHO concentration of 10% with 20mM sodium. Our purpose with using the higher CHO concentration was to induce the GLUT2 absorption pathway in BBM. The dextrose group has a higher mean compared with the sodium dextrose group, and this condition may indicate that at a higher CHO concentration, it trigger absorption through the GLUT 2 pathway because GLUT2 is rapidly and transiently recruited to the brush border membrane of the enterocyte, leading to a three-fold enhancement of glucose transport [23]. It is also known that the apical GLUT-2 absorption component is greater than the SGLT-1 at high luminal glucose concentrations [24,25].

Traditionally, SGLT-1 is the main route of CHO absorption in the villous BBM, in which absorption of glucose is coupled with the presence of sodium through the Na/K-ATPase activity [26,27], but GLUT-2, another transporter, can also be recruited to the BBM when high concentrations of glucose are present [28]. Expression of BBM GLUT-2 by the high intestinal luminal glucose concentrations is stimulated by enterocyte AMP kinase (AMPK), triggered by the opening of Cav 1.3 Ca^{2+} channels following SGLT-1-dependent depolarization of the apical membrane potential [6].

Apical GLUT-2 is tightly regulated by long, and short-term CHO supply and by local and endocrine hormones, disease (diabetes mellitus), cellular energy level, and the regulation occurs through intracellular signalling pathways [29]. Thus, apical GLUT-2 can act as the main pathway for sugar absorption. Apical GLUT-2 in the disease model acts as a guard mechanism by preventing high concentrations of CHO escape into the large intestine [30].

At ingestion rates over 60–70 g of CHO per h, exogenous CHO oxidation peaks around 50-60 g/h, but at 180 g/h, did not increase exogenous CHO oxidation rates much above 60 g/h [26]. Because this limitation was not caused by gastric emptying, liver glycogen storage, and muscle glucose

uptake, it was deduced that absorption had to be limiting [28]. When fructose was ingested in addition to larger amounts of glucose, CHO oxidation rates were elevated above 60 g/h [7], suggested that glucose transport across the epithelial cell was the limiting factor and that the maximal transport capacity of SGLT-1 was reached [31].

Evidence shows that the diffusion part is mediated by the translocation of the GLUT-2 from the intracellular enterocyte vesicles into the apical membrane, especially after high luminal glucose, allowing bulk absorption of glucose [32,33]. The apical GLUT-2 pathway from luminal sugar uptake could be inserted within minutes into the apical of enterocytes in response to high glucose levels during food assimilation [8]. As in physiologic condition, after a meal, the glucose level in the gut lumen rises, which could saturate the SGLT-1. The GLUT-2 pathway in the apical membrane then facilitates CHO diffusion down the concentration gradient. This study used 150 ml of dextrose 10%, equivalent to 15 grams of dextrose.

The slower mean increment of BG in the administration of sodium dextrose solution can be partly explained by the fact that SGLT-1 is a rate-limiting pathway for enterocyte glucose absorption. SGLT-1 uses a transmembrane sodium gradient to carry out the cellular uptake [34]. Such a situation may occur slower, considering the condition of soccer players is not in the actual match condition [35], suggesting lesser sodium loss through sweat perspiration that may result in a weaker sodium transmembrane gradient [36,37]. One main rule when using sports drinks that contain dextrose and sodium is that they should not be diluted. This would change the concentrations of CHO and sodium. Both will affect the speed at which the drink empties from the stomach, so the whole energy and fluid delivery mechanism will be delayed. The benefits of using sports drinks are proven only for adults involved in rigorous exercise and doing training in the hot condition and high humidity [20].

Soccer is an intermittent, high-intensity sport. Carbohydrates in the formed BG and muscle glycogen are the major substrate to replenish the adenosine triphosphate (ATP) in the contracting skeletal muscles during high and moderate-intensity work-out [38]. ATP is the form of the primary energy system, but it is only stored in small amounts (8 mmol/kg wet weight of muscle), and the phosphocreatine (PCr) system will resynthesize the ATP again from adenosine diphosphate [39]. The stored amounts of ATP and PCr are used up within 10 seconds of an intense exercise or an explosive movement like a sprint because of more than a 1,000-fold increase in ATP demand [39]. In order to deplete the ATP-PCr system, we designed the 2 × 100 m of a sprint (the first 100 m is to reduce the ATP-PCr system) with a 30 sec of resting time in between and followed the second 100 m (because to recover the PCr stores in muscle, it may take time around from 5 minutes) [39]. Once the storage is depleted, the body will use the BG in the glycolytic system to maintain a continuous ATP supply. As a metabolic fuel, BG, and muscle glycogen used to drop to levels that may damage cognitive function [40]. After dextrose supplementation, skeletal muscle is responsible for taking up 70–90% of the blood glucose [41]. Hence, soccer players are often advised to consume the CHO solution during competitions to supply muscle glycogen reservations and maintain BG concentrations during the match [42] [43]. Consumption of

CHO is recommended to maintain endogenous energy stores and prevent excessive physical and cognitive performance decline throughout the match [42].

Consumption of CHO has been proven to improve the performance index in playing soccer, where the reduction in BG levels can negatively influence the skilled actions involved [44]. Extended CHO release from a low glycemic index food also supports the benefits to fuel the bodily glucose reserve for endurance athletes [45]. Considering the performance benefits of CHO supplementation, which can reduce glycogen "depletion" levels throughout the game, it is recommended that soccer players consume 30–60 g/h of exogenous CHO supplementation during competitions to maintain the BG level and glycogen reserves [46]. However, the superior effect on the rate of increase in BG of pure ergogenic dextrose supplementation compared with sodium dextrose has not yet been known, therefore started from that point, this study was conducted.

In soccer players, an increase in BG levels can be observed from 75 min onwards when a 10% CHO-electrolyte solution is consumed before and during soccer matches, compared to a placebo-fluid-electrolyte control [47]. 6–7% of CHO solutions have succeeded in increasing the sprint speed [43]. Glycogen breakdown declines with a simultaneous elevation in extra-muscular BG levels to maintain normal BG during the match [19]. CHO supplementation improved intermittent running, soccer skill performance, the distance covered during the game and was correlated with the maintenance of glycogen levels compared to a placebo beverage [19]. In our trial, the faster sprint speed in the dextrose group might be explained by the effect of the faster increment in BG mean in the dextrose group, which served as a readily available substrate to fuel the muscle glycogen. CHO concentration in the sports drinks range between 4–8%, and the sodium content is between 10–25 mM [22], but in our trial, we used a higher CHO concentration reached 10% with 20 mM sodium. This fact may explain that it will trigger absorption through the GLUT 2 BBM pathway at higher CHO concentrations. Half-time for apical GLUT 2 membrane insertion is around 3.5 min (hence we used a 15 minutes period before tested the effect, but still below 20 minutes since it will provoke the insulin release) and correlated with activation of the protein kinase C β II isoform, which mark the physiological importance of fast response to increasing BG levels [48]. The transient presence of GLUT 2 in the apical enterocyte has been proved using antibodies to epitopes within N-terminal regions [49]. Cell immunofluorescence imaging's location in the apical membrane has been confirmed using extracellular loop antibodies [50]. GLUT 2 can transport glucose and has a very high K_t (transport constant) for glucose, indicate that apical GLUT 2 has the role of the diffusive component of glucose absorption [5], which at high glucose levels, the GLUT2 component accounted for \pm 75% of the absorption [8]. When the glucose concentration is higher in the intestinal gut lumen than in plasma, rapid trafficking of GLUT2 to the BBM accounts for the glucose absorption [51]. With BG level rising faster in the dextrose group, it will provide the added benefit of a faster sprint speed, as the muscle can readily uptake the surge elevation of BG. One of the possible underlying mechanisms was the increased availability of substrates, which can be used as a rapid fuel for muscle contraction [52], because during the exercise,

the amount of muscle glucose uptake increases, which it is leading to a decrease in blood glucose levels [53].

VO_{2max} differences between groups given dextrose compared with sodium dextrose were not significant. Soccer is a predominantly intermittent aerobic sport interspersed with periods of high-intensity anaerobic, which may perform different exercise intensities and use the aerobic and anaerobic energy systems. The total duration of active play in soccer is \pm 95 min, suggested the main energy source during the match is supplied through aerobic glycolysis. The average VO_{2max} during match play was reported to be around 75–80% of maximum oxygen uptake [54]. The mean and peak heart rate of players was estimated to be around 85 and 98%, respectively [55]. Nonetheless, VO_{2max} values of soccer players were shown to vary noticeably, with current high-level soccer players are suggested to acquire maximal oxygen uptakes between 60–70 ml/kg/min, with an estimated minimum of 65 ml/kg/min in elite soccer players [19].

People with high fitness (as measured by VO_{2max}) are more physically active in their daily life than those with low fitness. VO_{2max} is a modifiable variable, and it can be increased through exercise. High-intensity interval training was found to improve VO_{2max} more effectively than moderate continuous sessions [56]. We found no differences between the two groups because it has been known that VO_{2max} is not only influenced by BG levels but also by other factors, such as cardiorespiratory fitness [57]. However, to determine if a dose-response relationship exists between exercise intensity and training-induced increases in VO_{2max} , a result from meta-analysis suggests that exercise training intensity does not affect the elevation of training-induced VO_{2max} . However, similar adaptations can be achieved in low training doses at higher exercise intensities than higher training doses of lower intensity [58]. VO_{2max} is a measurement of the maximum capacity to utilize oxygen in multilevel exercise tests and acts as a measure of soccer players' cardiorespiratory fitness system. The cardiovascular and thermoregulatory systems are first affected by low hydration, especially when exercise is performed in high temperature or high humidity conditions [20], but in this experiment design, the players were not performed in the heat environment. Nevertheless, our study's limitation is that the research was not performed under a real soccer match.

5. Conclusions

Our study indicated that solely dextrose supplementation effectively supports blood sugar and sprint speed in soccer players. A solution of 150 ml dextrose 10% elevated blood glucose faster than sodium dextrose solution. Based on our results, we conclude that concerning the beginning of the soccer match, a 10% of dextrose solution ingested 15 minutes before the match could elevate BG faster and may have a good impact on sprint velocity, as at the beginning of a match, to kick off and dominate it. These findings suggest a future suggestion to re-evaluate the energy drink, which nearly always has a mixture of CHO and electrolyte.

Disclosure of interest

10

The authors would like to thank the participants for their effort and cooperation.

The authors declare that they have no competing of interest.

References

- [1] Russell M, Benton D, Kingsley M. Carbohydrate ingestion before and during soccer match play and blood glucose and lactate concentrations. *J Athl Train* 2014;49(4):447–53.
- [2] Russell M, Rees G, Benton D, Kingsley M. An exercise protocol that replicates soccer match-play. *Int J Sports Med* 2011;32(7):511–8.
- [3] Bangsbo J, Iaia FM, Krstrup P. Metabolic Response Fatigue. *Int J Sports Physiol Perform* 2007;2:111–27.
- [4] Navale AM, Paranjape AN. Glucose transporters: physiological and pathological roles. *Biophys Rev* 2016;8(1):5–9.
- [5] Koepsell H. Glucose transporters in the small intestine in health and disease. *Pflugers Arch Eur J Physiol* 2020;472(9):1207–48.
- [6] Jeukendrup AE. Training the Gut for Athletes. *Sport Med* 2017;47(s1):101–10.
- [7] Jeukendrup A. A step towards personalized sports nutrition: Carbohydrate intake during exercise. *Sport Med* 2014;44(Suppl. 1).
- [8] Kellett GL, Brot-Laroche E, Mace OJ, Leturque A. Sugar absorption in the intestine: the role of GLUT2. *Annu Rev Nutr* 2008;28(1):35–54.
- [9] Mace OJ, Affleck J, Patel N, Kellett GL. Sweet taste receptors in rat small intestine stimulate glucose absorption through apical GLUT2. *J Physiol* 2007;582(1):379–92.
- [10] Naftalin RJ. A computer model simulating human glucose absorption and metabolism in health and metabolic disease states. *F1000Research* 2016;5(0):1–37.
- [11] Kimmich GA, Randles J. Sodium-sugar coupling stoichiometry in chick intestinal cells. *Am J Physiol* 1984;247(1 Pt 1):74–82.
- [12] MacLeod RJ, Hamilton JR. Volume regulation initiated by Na⁺ - nutrient cotransport in isolated mammalian villus enterocytes. *Am J Physiol - Gastrointest Liver Physiol* 1991;260(1 23-1).
- [13] Boyd CA, Parsons DS. Movements of monosaccharides between blood and tissues of vascularly perfused small intestine. *J Physiol* 1979;287(1):371–91.
- [14] Esposito BP, Breuer W, Sirankapracha P, Pootrakul P, Hershko C, Cabantchik ZI. Labile plasma iron in iron overload: Redox activity and susceptibility to chelation. *Blood* 2003;102(7):2670–7.
- [15] Naftalin RJ. Does apical membrane GLUT2 have a role in intestinal glucose uptake? *F1000Research* 2014;3:1–33.
- [16] Gabbett TJ, Mulvey MJ. Time-motion analysis of small-sided training games and competition in elite women soccer players. *J Strength Cond Res* 2008;22(2):543–52.
- [17] Hargreaves M, Spriet LL. Skeletal muscle energy metabolism during exercise. *Nat Metab* 2020;2(9):817–28.
- [18] Swanwick E. Energy Systems: A new look at aerobic metabolism in stressful exercise. *MOJ Sport Med* 2018;2(1):15–22.
- [19] Alghannam FA. Physiology of soccer: the role of nutrition in performance. *J Nov Physiother* 2013.
- [20] Simulescu V, Ilia G, Macarie L, Merghes P. Sport and energy drinks consumption before, during and after training. *Sci Sport* 2019;34(1):3–9.
- [21] Afshar N, Safaei S, Nickerson DP, Hunter PJ, Suresh V. Computational modeling of glucose uptake in the enterocyte. *Front Physiol* 2019;10:1–12.
- [22] Burke LM, Meyer NL, Pearce J. National nutritional programs for the 2012 London Olympic Games: a systematic approach by three different countries. *Nestle Nutr Inst Workshop Ser* 2013;76:103–20.
- [23] Merino B, Fernández-Díaz CM, Cózar-Castellano I, Perdomo G. Intestinal fructose and glucose metabolism in health and disease. *Nutrients* 2020;12(1):1–35.
- [24] Schreck K, Melzig MF. Intestinal saturated long-chain fatty acid, glucose and fructose transporters and their inhibition by natural plant extracts in caco-2 cells. *Molecules* 2018;23(10):1–30.
- [25] Röder PV, Geillinger KE, Zietek TS, Thorens B, Koepsell H, Daniel H. The role of SGLT1 and GLUT2 in intestinal glucose transport and sensing. *PLoS One* 2014;9(2):20–2.
- [26] Dobbins RL, Greenway FL, Chen L, Liu Y, Breed SL, Andrews SM, et al. Selective sodium-dependent glucose transporter 1 inhibitors block glucose absorption and impair glucose-dependent insulinotropic peptide release. *Am J Physiol - Gastrointest Liver Physiol* 2015;308(11):G946–54.
- [27] Erokhova L, Horner A, Ollinger N, Siligan C, Pohl P. The sodium glucose co-transporter SGLT1 is an extremely efficient facilitator of passive water transport. *J Biol Chem* 2016;291(18):9712–20.
- [28] Kellett GL. The facilitated component of intestinal glucose absorption. *J Physiol* 2001;531(3):585–95.
- [29] Kellett GL, Brot-Laroche E. Apical GLUT2: a major pathway of intestinal sugar absorption. *Diabetes* 2005;54(10):3056–62.
- [30] Kellett GL. Perspectives in diabetes. *J Natl Med Assoc* 2005;54:3056–62.
- [31] Stellingwerff T, Pfeiffer B, Stellingwerff T, Zaltas E, Jeukendrup AE. Oxidation of solid versus liquid CHO sources during exercise: 2015. p. 2030–8.
- [32] Chaudhry RM, Scow JS, Madhavan S, Duenes JA, Sarr MG. Acute enterocyte adaptation to luminal glucose: a posttranslational mechanism for rapid apical recruitment of the transporter GLUT2. *J Gastrointest Surg* 2012;16(2):312–9.
- [33] Scow JS, Tavakkolizadeh A, Zheng Y, Sarr MG. Acute "adaptation" by the small intestinal enterocyte: a post-transcriptional mechanism involving apical translocation of nutrient transporters. *Surgery* 2011;149(5):601–5.
- [34] Seidelmann SB, Feofanova E, Yu B, Franceschini N, Claggett B, Kuokkanen M, et al. Genetic variants in SGLT1, glucose tolerance, and cardiometabolic risk. *J Am Coll Cardiol* 2018;72(15):1763–73.
- [35] Godek SF, Peduzzi C, Burkholder R, Condon S, Dorshimer G, Bartolozzi AR. Sweat rates, sweat sodium concentrations, and sodium losses in 3 groups of professional football players. *J Athl Train* 2010;45(4):364–71.
- [36] Baker LB. Sweating rate and sweat sodium concentration in athletes: a review of methodology and intra/interindividual variability. *Sport Med* 2017;47(s1):111–28.
- [37] Thorsen K, Drengstig T, Ruoff P. Transepithelial glucose transport and Na⁺/K⁺ homeostasis in enterocytes: an integrative model. *Am J Physiol - Cell Physiol* 2014;307(4):320–37.
- [38] Mul JD, Stanford KI, Hirshman MF, Goodyear LJ. Exercise and regulation of carbohydrate metabolism. *Prog Mol Biol Transl Sci* 2015;135:17–37.
- [39] Baker JS, McCormick MC, Robergs RA. Interaction among skeletal muscle metabolic energy systems during intense exercise. *J Nutr Metab* 2010;2010 (Figure 1).
- [40] Russell M, Kingsley M. The efficacy of acute nutritional interventions on soccer skill performance. *Sport Med* 2014;44(7):957–70.
- [41] Evans PL, McMillin SL, Weyrauch LA, Witczak CA. Regulation of skeletal muscle glucose transport and glucose metabolism by exercise training. *Nutrients* 2019;11(10):1–24.
- [42] Hills SP, Russell M. Carbohydrates for soccer: a focus on skilled actions and half-time practices. *Nutrients* 2018;10(1).

- [43] Baker LB, Rollo I, Stein KW, Jeukendrup AE. Acute effects of carbohydrate supplementation on intermittent sports performance. *Nutrients* 2015;7:5733–63.
- [44] Russell M, Benton D, Kingsley M. Influence of carbohydrate supplementation on skill performance during a soccer match simulation. *J Sci Med Sport* 2012;15(4):348–54.
- [45] Gao R, Shaw K, Kaviani M, Gordon J, Zello GA, Chilibeck PD. Glycemic index of lentil- and cherry-based sport nutrition products for endurance athletes. *Sci Sport* 2020:1–6.
- [46] Thomas DT, Erdman KA, Burke LM. American college of sports medicine joint position statement. nutrition and athletic performance. *Med Sci Sport Exerc* 2016;48(3):543–68.
- [47] Kingsley M, Penas-Ruiz C, Terry C, Russell M. Effects of carbohydrate-hydration strategies on glucose metabolism, sprint performance and hydration during a soccer match simulation in recreational players. *J Sci Med Sport* 2014;17(2):239–43.
- [48] Cohen M, Kitsberg D, Tsytkin S, Shulman M, Aroeti B, Nahmias Y. Live imaging of GLUT2 glucosdependent trafficking and its inhibition in polarized epithelial cysts. *Open Biol* 2014;4.
- [49] Long W, Cheeseman CI. Structure of, and functional insight into the GLUT family of membrane transporters. *Cell Health Cytoskelet* 2015;7:167–83.
- [50] Ait-Omar A, Monteiro-Sepulveda M, Poitou C, Le Gall M, Cotillard A, Gilet J, et al. GLUT2 accumulation in enterocyte apical and intracellular membranes: a study in morbidly obese human subjects and ob/ob and high fat-fed mice. *Diabetes* 2011;60(10):2598–607.
- [51] Thazhath SS, Wu T, Young RL, Horowitz M, Rayner CK. Glucose absorption in small intestinal diseases. *Expert Rev Gastroenterol Hepatol* 2014;8(3):301–12.
- [52] Jentjens RPLG, Achten J, Jeukendrup AE. High oxidation rates from combined carbohydrates ingested during exercise. *Med Sci Sports Exerc* 2004;36(9):1551–8.
- [53] Gholizadeh M, Rahmaninia F, Siahkohian M. Effects of high and low glycemic isoenergetic breakfast on glucose homeostasis and substrate oxidation during high intensity intermittent exercise. *Sci Sport* 2020:1–10.
- [54] Slimani M, Znazen H, Miarka B, Bragazzi NL. Maximum oxygen uptake of male soccer players according to their competitive level. playing position and age group: implication from a network meta-analysis. *J Hum Kinet* 2019;66(1):233–45.
- [55] Bangsbo J, Mohr M, Krstrup P. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci* 2006;24(7):665–74.
- [56] Zisko N, Stensvold D, Hordnes-Slagsvold K, Rognmo Ø, Nauman J, Wisløff U, et al. Effect of change in $\dot{V}O_2\max$ on daily total energy expenditure in a cohort of norwegian men: a randomized pilot study. *Open Cardiovasc Med J* 2015;9(1):50–7.
- [57] Amman M. V AMC. Pulmonary system limitations to endurance exercise performance in humans. *Exp Physiol* 2012;97(3):311–8.
- [58] Trisha S. School of K. the effect of training intensity on $\dot{V}O_2\max$ in young healthy adults: a meta-regression and meta-analysis. *Int J Exerc Sci* 2016;9(2):230–47.

Dextrose 10% drink is superior to sodium-dextrose drink in increasing blood glucose and sprint speed in soccer players: A double-blinded randomized crossover trial study

ORIGINALITY REPORT

15%

SIMILARITY INDEX

10%

INTERNET SOURCES

7%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universiti Teknologi MARA Student Paper	2%
2	meridian.allenpress.com Internet Source	1%
3	opencardiovascularmedicinejournal.com Internet Source	1%
4	Hermann Koepsell. "Glucose transporters in the small intestine in health and disease", Pflügers Archiv - European Journal of Physiology, 2020 Publication	1%
5	jissn.biomedcentral.com Internet Source	1%
6	Submitted to University of Brighton Student Paper	1%
7	eprints.bournemouth.ac.uk Internet Source	<1%

8	Submitted to Flinders University Student Paper	<1 %
9	Submitted to Mary Baldwin University Student Paper	<1 %
10	onlinelibrary.wiley.com Internet Source	<1 %
11	hdl.handle.net Internet Source	<1 %
12	journals.plos.org Internet Source	<1 %
13	Submitted to SCL Education Student Paper	<1 %
14	www.karger.com Internet Source	<1 %
15	M. Gholizadeh, F. Rahmaninia, M. Siahkohian. "Effects of high and low glycemic isoenergetic breakfast on glucose homeostasis and substrate oxidation during high intensity intermittent exercise", Science & Sports, 2020 Publication	<1 %
16	iranarze.ir Internet Source	<1 %
17	qspace.library.queensu.ca Internet Source	<1 %

18

Rajia Rajia, Muh Nasrum Massi, Mardiana Ahmad, Sharvianty Arifuddin, Upik Anderiani Miskad. "Parity, exposure to cigarette smoke and the presence of bacteria *Stenotrophomonas maltophilia* are related to preterm labor incidence", *Enfermería Clínica*, 2020

Publication

<1 %

19

digitalcommons.wku.edu

Internet Source

<1 %

20

Submitted to Universitas Hasanuddin

Student Paper

<1 %

21

Drug Discovery and Evaluation
Pharmacological Assays, 2016.

Publication

<1 %

22

www.tandfonline.com

Internet Source

<1 %

23

Submitted to Leeds Metropolitan University

Student Paper

<1 %

24

bmcinfctdis.biomedcentral.com

Internet Source

<1 %

25

Submitted to Napier University

Student Paper

<1 %

26

Submitted to Nottingham Trent University

Student Paper

<1 %

27	Submitted to University of Fort Hare Student Paper	<1 %
28	ir.canterbury.ac.nz Internet Source	<1 %
29	jyx.jyu.fi Internet Source	<1 %
30	www.jove.com Internet Source	<1 %
31	www.wjgnet.com Internet Source	<1 %
32	ejnmires.springeropen.com Internet Source	<1 %
33	koreascience.or.kr Internet Source	<1 %
34	link.springer.com Internet Source	<1 %
35	www.e-pan.org Internet Source	<1 %
36	Chooi Yeng Lee. "Chronic restraint stress induces intestinal inflammation and alters the expression of hexose and lipid transporters", Clinical and Experimental Pharmacology and Physiology, 2013 Publication	<1 %

37 Hee Kyong Na, Kee Don Choi, Young Soo Park, Hwa Jung Kim et al. "Endoscopic Scoring System for Gastric Atrophy and Intestinal Metaplasia: Correlation with OLGA and OLGIM Staging: A Single Center Prospective Pilot Study", Research Square Platform LLC, 2021

Publication

<1 %

38 Thorsen, K., T. Drengstig, and P. Ruoff. "Transepithelial glucose transport and Na⁺/K⁺ homeostasis in enterocytes: an integrative model", AJP Cell Physiology, 2014.

Publication

<1 %

39 escholarship.umassmed.edu

Internet Source

<1 %

40 www.spandidos-publications.com

Internet Source

<1 %

41 "Abstract Issue", Planta Medica, 2019

Publication

<1 %

42 Adina Braha, Alin Albai, Romulus Timar, Laura Diaconu, Lucian Vasiluță, Daniela Cipu, Bogdan Timar, Alexandra Sima. "Factors Associated with the Remission of Type 1 Diastolic Dysfunction after Dapagliflozin Treatment in Patients with Type 2 Diabetes", Journal of Clinical Medicine, 2020

Publication

<1 %

43 ccforum.biomedcentral.com <1 %
Internet Source

44 eprints.soton.ac.uk <1 %
Internet Source

45 umexpert.um.edu.my <1 %
Internet Source

46 www.medrxiv.org <1 %
Internet Source

47 G. L. Kellett, E. Brot-Laroche. "Apical GLUT2: A Major Pathway of Intestinal Sugar Absorption", Diabetes, 2005 <1 %
Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On