

Changes of Anthocyanin Contents, Radical Scavenging Activities and Total Microbial Counts of Buni Fruit (*Antidesmabunius*) Juice during Different Storage Times

by Agussalim Bukhari

Submission date: 25-Apr-2021 09:37AM (UTC+0700)

Submission ID: 1568832945

File name: Agussalim_Bukhari-Suryani_tawali-2018.pdf (822.36K)

Word count: 3497

Character count: 19005

Research Article

Changes of Anthocyanin Contents, Radical Scavenging Activities and Total Microbial Counts of Buni Fruit (*Antidesma bunius*) Juice during Different Storage Times

Suryani Tawali^{1*}, Agussalim Bukhari², Mulyati M. Tahir³,

Nur Faidah Rahman³, Suryani As'ad², Mochammad Hatta⁴ and Indra Yuliana³

¹Department of Public Health and Preventive Medicine, Faculty of Medicine

²Department of Nutrition, Faculty of Medicine

³Department of Food Technology, Faculty of Agriculture

⁴Immunology and Biomolecular Laboratory/ Department of Microbiology, Faculty of Medicine

^{1,2,3,4}Hasanuddin University, Makassar, Indonesia

*Corresponding author : email address : suryani_tawali@med.unhas.ac.id suryani_tawali@yahoo.com
 telephone no. : +62411586010 : +62411586297

ABSTRACT

Background: Cardiovascular disease is still the leading cause of death in the world. Damage to vascular endothels resulted from oxidative reactions caused by excessive exposure to free radicals due to changes in lifestyle and environment are the main cause of the problem. Consumption of foods with high antioxidants content is believed can prevent and ceased the damaging process in the blood vessels. A local Indonesian fruit that has a potentially high antioxidant content is buni fruit (*Antidesma bunius*). The aim of this study was to investigate the changes of anthocyanin content, antioxidant capacity, total microbial counts of buni fruit juice during storage time. **Results:** The total anthocyanin content was 302.03 - 496.26 mg/ 100 g fresh. The content decreased to 155-166.48 mg/ 100 g after one week storage. Then further decreased to 94.61-136.46; 64.59-100.37; 59.44-81.57 mg/ 100 g after 2, 3, and 4 weeks storage, respectively. The highest antioxidant capacity is in the fresh juice: 66-70%, and it reduced significantly to 2-12% in the 4th week storage time. Total microbial count increased from 0 in fresh juice to 2- 5 x 10⁵ CFU / ml in week 3. **Conclusion :** Buni fruit (*Antidesma bunius*) has a relatively high anthocyanin content and antioxidant capacity, however, during storage, the bioactive contents and activities reduced and contamination of microbes increased. Therefore, further research is required to maintain the quality of buni fruit juice during storage.

Keywords : Bunifruit; *Antidesma bunius*; anthocyanins; antioxidant activities; antioxidant capacity; total microbial counts.

BACKGROUND

Cardiovascular disease is still the major cause of death in the world, more than 50% of deaths are caused by non-infectious diseases, and 48% are caused by the diseases of the heart and blood vessels. Changes in lifestyle, high exposure to chemicals either from food or from air pollution are believed to be the risk factors for the onset of the disease. This is attributed to the increasing number of free radicals in the body that trigger

physiological changes in the level of cells that eventually leads to such diseases [1]. Free radicals are naturally produced in the physiological processes in the body. The number of free radicals increases if the body is working beyond its capacity. Besides these free radicals can also come from unhealthy foods, chemicals, exposure to ultraviolet rays, cigarette smoke and air pollution. In normal circumstances free

radicals are neutralized by antioxidants produced by the body's own (endogenous antioxidant). However, if the number of excessive free radicals in the body, the body can not neutralize all free radicals. And this is what causes damage to cells and tissues of the body, accumulation of fat in the arteries, hardening of the arteries that causes heart disease and hypertension [2]. Researchers have found that many plants contain substances that can neutralize free radicals, which can prevent and treat various diseases and cancer. Substances in plants that are believed to neutralize free radical oxidation reaction by most of the class of polyphenolic compounds. The large part of natural polyphenol compounds are flavonoids. Flavonoids are found in fruits and vegetables that are colorful. Many studies have proven the benefit of flavonoids in preventing a variety of chronic diseases, especially cardiovascular disease and cancer [3,4]. Anthocyanins as one group of flavonoids are abundant in colorful fruits and vegetables. Berries groups are known as sources of anthocyanins such as blueberry, raspberry, blackberry, gooseberry, etc. Many studies have revealed the benefits of fruit and vegetables anthocyanin for health [3,4,5, 6,7,8,9,10]. One of the local Indonesian plant that has similar physical appearance with berry group particularly cranberry is Buni fruit (Antidesmabunius). In ripe stage it has dark black-red color [11]. It has been used as traditional remedy for some diseases and consumed raw or processed in relatively small amount particularly in rural places. However it becomes less popular since people tend to consume more "modern, western" fruit such as grapes and apples. One reason why people is becoming less likely to eat this fruit because its taste which is sour, and also the benefit of this fruit has not been explored and promoted. In Thailand, the fruit is known as Mao Luang. Samappito and Butkhup has found that buni fruit in Northeast Thailand contained high amount of flavonoids, they are catechin, procyanidin B1, procyanidin B2 and several organic acids [12,13]. In South Sulawesi, Indonesia, the fruits are harvested from the end of February to April, and they can be found in traditional

markets during harvest seasons, however because of the low demand for consumption the amount of buni fruit marketed decreased in the recent decade.

The aim of this study was to investigate the anthocyanin content of buni fruit juice and its antioxidant capacity and the total microbial colony.

MATERIALS AND METHODS

Materials and Sample Preparation

Buni fruits (Antidesmabunius) were purchased from local markets (Makassar, Indonesia). Only the ripe fruits which were completely reddish and red-black in color were chosen. The collected berries were washed, drained, then weighted. Five hundred grams fruit was extracted using water with ratio 1 : 1, then filtered. The extraction was repeated four times until the anthocyanin content could not be detected anymore. Twenty a 150 ml juice samples were treated differently, 5 samples without added fructose, 5 with fructose 5%, 5 with fructose 10%, and the rest 5 samples with added fructose 15%. The juice samples then bottled in the dark bottles, pasteurized, and sealed then stored at 10°C until further analysis.

Chemical Analysis

Analysis was conducted for anthocyanin, vitamin C, and TSS content, total microbial count, pH, antioxidant activities, and organoleptic test. All analysis was conducted every week (week 0, week 1, week 2, and week 3) except for organoleptic test which was conducted only in week 0. For each analysis consisted of two replications.

Anthocyanins

The total anthocyanins were estimated by a pH differential method [14,15,16]. From each sample, 0.05 ml was dissolved with buffer solution of potassium chloride 0.025 M (4.95 ml) pH 1.0 and another 0.05 ml with solution of sodium acetate buffer 0.4 M (4.95 ml). Using UV-Visible spectrophotometer, the absorbance for each sample was then read versus the buffer solution pH 1.0 and buffer solution pH 4.5 as the blank at $\lambda = 510 \text{ nm}$ (for the cyanidin 3-glucoside) and $\lambda = 700 \text{ nm}$ (for correction

factor). The absorbance values were calculated using the equation:

$$A = \{(A516 - A700) \text{ pH } 1 - (A516 - A700) \text{ pH } 4.5\}$$

Anthocyanin concentration is calculated as cyanidin-3-glycosides using a molar extinction coefficient of $29,600 \text{ L mol}^{-1} \text{ cm}^{-1}$ (ϵ); a molecular weight of 449.2 g mol^{-1} (MW) using the formula:

$$\text{TAC} = \frac{A}{C \times L} \times \text{MW} \times \text{DF} \times \frac{V}{\text{Wt}} \times 100\%$$

Where TAC is total anthocyanin content ($\text{mg} \cdot 100 \text{ g}^{-1}$); A is absorbance value; ϵ is molar extinction coefficient; L is width of cuvette; DF is dilution factor; V is final volume after dilution; and Wt is weight of original extract.

Antioxidant capacity

Antioxidant capacity was measured by the ability of the sample to scavenge the free radicals [15,17]. The free radical used was 1,1-diphenyl-2-picrylhydrazyl (DPPH). The DPPH analysis was carried out using method from Brand-Williams et. al. [18] with modification. To make DPPH solution, 0,0157 gram DPPH dissolved in 100 ml of absolute ethanol in a measured flask. The radical scavenging activity of the buni fruit juice was measured by adding 1 ml DPPH solution into 100 ml juice sample and added with ethanol until the solution reached 5 ml. The mixture then vortexed and allowed for about 30 minutes in a room temperature. The absorbance then measured at 518 nm. Radical scavenging activity was determined with the formula :

$$\% \text{ free radical scavenging activity} = \frac{\text{blank absorbance} - \text{sample absorbance}}{\text{blank absorbance}} \times 100\%$$

Total Microbial Counts

Total microbial counts was detected using Plate Count Agar (PCA) method [19, 20,21] with modification. From each treatment, 1 gram sample was taken. Each sample then put into a test tube contained 9 ml distilled water then shaken until suspension was formed at the

bottom of the tube. One ml of the suspension was taken from the tube and inserted into another tube and diluted to 10^4 dilution. One ml of the solution from the second tube was pipetted and put into a petri dish aseptically. After that, Fifteen ml liquid agar that had been cooled to 50°C was poured into the dish, then shaken carefully in order that the microbial cells could spread evenly. If the agar become solid, the petri dish then incubated in inverted position for approximately 48 hours at a temperature of 37°C . The total microbes was counted using the formula :

$$\left(\frac{\log}{\text{CFU}}\right) = \text{Number of colonies} \times 1/\text{dilution}$$

Statistical Analysis

Values shown are means \pm standard deviation from duplicate samples.

RESULT AND DISCUSSION

Anthocyanins

It shown in table 1. and figure 1 that the highest total anthocyanin content of buni fruit juice was found in the fresh fruit juice whether with or without added fructose that was 302.03 - 496.26 mg/ 100 g fresh fruit. It averagely similar with the anthocyanins content of cultivated blueberry and wild blueberry which were 365 mg and 487 mg per 100 g respectively [23]. The concentration of anthocyanin in buni fruit juice consistently reduced during storage time. After seven days the anthocyanin concentrations decreased to 135.55-166.48 mg/ 100 g . Then further decreased to 94.61-136.46; 64.59-100.37; 59.44-81.57 mg/ 100 g after 2, 3, and 4 weeks storage respectively.

The reduction of anthocyanin concentrations during storage time are influence by many factors. It is known that anthocyanins are unstable compounds. Light, storage, oxygen, heat-humidity, enzymes, and the presence of ascorbic acid, sugar, sulfur dioxide, sulfite salts, metal ions and copigments influence the concentration of anthocyanin in food [24,25,26].

Changes of Anthocyanin Contents, Radical Scavenging Activities and Total Microbial Counts of Buni Fruit (Antidesmabunius) Juice during Different Storage Times

Table.1. Changes of anthocyanin content (mg/ 100 gr) of buni fruit (Antidesmabunius) during 4 weeks storage time

Fructose concentration (%)	Storage time (Week)				
	0	1	2	3	4
0	496.26±21.38	166.48±3.64	136.46±0.91	83.09±35.78	81.57±37.30
5	379.36±99.16	151.92±19.1	126.45±35.48	100.37±6.37	67.32±5.46
10	371.17±58.22	135.55±40.94	94.61±4.55	79.75±23.35	58.53±21.53
15	302.03±14.56	120.54±15.01	84.18±3.21	64.59±33.66	59.44±26.69

** mean value± sd (n = 2)




Figure 1. The decrease of anthocyanin content of buni fruit (Antidesmabunius) during 4 weeks storage time

10 Antioxidant capacity

Antioxidants are chemical compounds that can donate one or more electrons to free radicals, so the free radicals that can be neutralised. Antioxidant activity testing methods used in this study are DPPH (1,1-Diphenyl-2-picrylhydrazyl). DPPH method is a simple method to determine the antioxidant activity of a food using DPPH radical [18]. DPPH test is a colorimetric method that is effective and quick to estimate anti-radical activity. DPPH radical is an organic compound containing nitrogen unstable with strong absorbance at = 517 nm and dark purple [17].




Figure 2. Changes of DPPH radical scavenging activity of buni fruit juice during storage time

The value of antioxidant activity of buni fruit juice with different concentrations and different storage time can be seen in Figure 2. The graph shows that the highest antioxidant activity is in the fresh fruit juice that is 66-70%, while the lowest antioxidant activity is in fruit juice that stored for three weeks that is 2-12%. The antioxidant activity is likely to experience continued decline over time. The ability of the fruit juice to neutralize free radicals depends on the antioxidant concentrations in it. In buni fruit, even though many plant phenolic compounds were present, however, the prominent antioxidant found was anthocyanin. The radical scavenging activity of buni fruit juice reduced concurrently with the reduction of the anthocyanin concentrations. On the other hand, the concentration of added sugar did not influence the antioxidant capacity. As a part of phenolic compounds, anthocyanins has aromatic rings that can donate its electron to stabilize the free radicals in a solution. However, the contribution of other phenolic compounds in buni fruit juice in the antioxidant activities could not be overseen. The phenolics contents of

colorful fruits and vegetables are proven to be beneficial for health [24,27].

Total microbial counts

From figure 3, it can be seen that the effect of fructose concentrations and storage time significantly high to the total microbial counts. In fresh fruit juice, the total microbial counts is 0, therefore, the fresh fruit juice after processing without storage is still considered safe from microbial growth. However, the microbes had started to grow and continuously increased every week. After seven days storage, the total microbial counts was 1.7×10^4 CFU/ml. The density of microbial growth was the highest in the third week after storage: 2.4×10^5 CFU/ml. However, there was an inverted association between concentrations of added sugar with the growth of microbes. The less concentration of the sugars, the more likely the microbes to grow. There are two factors that affect microbial growth are intrinsic factors (pH, aw, nutrient content, biological structure and content of anti-microbial) and extrinsic factors (storage temperature, relative humidity, type and amount of gas in the environment).




Figure 3. Changes of total microbial counts during storage time

This is consistent with argument from Buckle, which stated that a sugar solution with a high concentration (TSS > 70 ° Brix) will be able to provide stability of microorganisms on a product [28]. Manufacture of a product with high sugar concentrations is one important food

preservation technique. This is because the decrease of water activity (aw) is associated with the increase in the concentration of sugar itself, so that the water becomes less available for the growth of microorganisms. However, one should consider other food preservation

methods according to the characteristics of the bioactive compounds in a product in order to preserve the maximal nutritional benefits of it.

CONCLUSION

Buni fruit (*Antidesma bunius*) has a relatively high anthocyanin content and antioxidant capacity with averagely similar with other fruits with high anthocyanin content such as blueberry. However the concentration of anthocyanin and the fruit radical scavenging activity showed a marked reduction during storage time in temperature of 10°C. Therefore, further studies have to be done to find out the appropriate processes and storage methods in order to preserved the anthocyanin content and it antioxidant capacity.

ACKNOWLEDGEMENT

This research was funded by Directorate of Higher Education , The Ministry of Research and High Education Grant.

REFERENCES

1. Mendis S, Puska P & Norrving B (eds) (2011), Global atlas on cardiovascular disease prevention and control, World Health Organization, Geneve.
2. Andreyev A. Yu, Yu.E. Kushnareva, A.A. Starkov (2004) Mitochondrial metabolism of Reactive oxygen species, *Biochemistry (Moscow)* vol.70 no.2 , pp.200-2004.
3. De Pascual-Teresa S, Diego A. Moreno, Garcia-Viguerra C (2011) Flavanols and anthocyanins in cardiovascular health: a review of current evidence, *The international Journal of Molecular Science* 11: 1679-1703 ; doi: 10.3390/ijms 11041679.
4. Qin Xia, En, et all (2011) Biological activities of poyphenols from grapes, *The international Journal of Molecular Science* 11: 622- 646; doi: 10.3390/ijms 11020622.
5. Duthie, SJ, McE. Jenkinson, A, Crozier, A, Mullen, W, Pirie, L, Kyle, J, Yap, LS, Christen, P & Duthie, GG, 2006, 'The effects of cranberry juice consumption on antioxidant status and biomarkers relating to heart disease and cancer in healthy human volunteers', *European Journal of Nutrition*, Vol.45, pp. 113–122.
6. Spormann, TM, Albert, FW, Rath, T, Dietrich, H, Will, F, Stockis, J, Eisenbrand, G & Janzowski, C (2008) 'Anthocyanin/Polyphenolic-Rich Fruit Juice Reduces Oxidative Cell Damage in an Intervention Study with Patients on Hemodialysis', *Cancer Epidemiology, Biomarkers & Prevention*, Vol. 17, pp. 3372-3380.
7. Erlund, I, Koli, R, Alfthan, G, Marniemi, J, Puukka, P, Mustonen, P, Mattila, P, & Jula, A, 2008, 'Favorable effects of berry consumption on platelet function, blood pressure, and HDL cholesterol', *American Journal of Clinical Nutrition*, Vol.87, pp. 323–31.
8. Wang, LS and Stoner GD (2008). Anthocyanins and their role in cancer prevention. *Cancer Lett.* 269(2): 281-290. Doi:10.1016/j.canlet.2008.05.020.
9. Qin, Y, Xia, M, Ma, J, Hao, Y, Liu, J, Mou, HY, Cao, L & Ling, WH (2009). Anthocyanin supplementation improves serum LDL- and HDL-cholesterol concentrations associated with the inhibition of cholesteryl ester transfer protein in dyslipidemic subjects', *American Journal of Clinical Nutrition*, Vol.90, pp. 485–92.
10. Riso, P, Klimis-Zacas, Del Bo', C, Martini, G, Campolo, J, Vendrame, S, Møller, P, Loft, S, De Maria, R & Porrini, M (2013) 'Effect of a wild blueberry (*Vaccinium angustifolium*) drink intervention on markers of oxidative stress, inflammation and endothelial function in humans with cardiovascular risk factors', *European Journal of Nutrition*, vol., pp.949–961.
11. Orwa C, AMutua, Kindt R, Jamnadass R, S Anthony (2009) *Agroforestry Database: a tree reference and selection guide version* (<http://www.worldagroforestry.org/sites/treedatabases.asp>).
12. Dutkhu L Samappito S (2008) *An analysis on flavonoids contents in Mao Luang fruits fifteen cultivars grown in Northeast*

- Thailand.Pakistan Journal of Biological Sciences 11 (7): 996-1002.
13. Samppitto S and Bukhtup L (2008) An analysis on organic acids contents in ripe fruits of fifteen Mao Luang(*Antidesmabunius*) cultivars, harvested from dipterocarp forest of Phupan valley in Northeast Thailand.Pakistan Journal of Biological Sciences 11 (7): 974-981.
 14. Wrolstad RE &Giusti MM. (2001) Characterization and measurement of anthocyanins by UV-visible spectroscopy. In: Food Analytical Chemistry (Wrolstad, R. E. ed.). New York:John Wiley & Sons, Inc. Unit F1.2.
 15. Prior RL, Wu S, Schaich K. (2005) Standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements. J. Agric. Food Chem 53: 4290-4302.
 16. Lee J, Rennaker C, Wrolstad RE. (2008) Correlation of two anthocyanin quantification methods: HPLC and spectrophotometric methods. Food Chemistry 110: 782-786.
 17. Kedare SG & Singh RP (2011). Genesis and development of DPPH method of antioxidant assay.J Food Sci Technol: 48(4):412-422 .DOI 10.1007/s13197-011-0251-1.
 18. Brand-Williams W, Cuvelier ME, Berset C (1995). Use of a Free Radical Method to Evaluate Antioxidant Activity. Lebensm.-Wiss. u.-Technol 28: 25-30.
 19. Buchbinder L, Baris Y, Alff E, Reynolds E, Dillon E, Essin V, Pincus L, Strauss A (1951) Studies To Formulate New Media for the Standard Plate Count of Dairy Products. Public Health Reports 66 No. 11 : 327-340.
 20. American Public Health Association (APHA) (1978). Standard Methods for the Examination of Dairy Products, 14th ed., APHA Inc., Washington, D.C.
 21. Sigma Aldrich .Plate Count Agar #70152. Product Information..Available from:<http://www.sigmaldrich.com/content/dam/sigma-aldrich/docs/Sigma-Aldrich/Datasheet/1/70152dat.pdf>.
 22. Akubor PI, Adejo EE (2000).Physicochemical, microbiological and sensory changes in stored plantain chips. Plant Foods for Human Nutrition 55: 139-146.
 23. Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, Prior RL (2006) Concentrations of anthocyanins of common foods in the United States and estimation of normal consumption. J. Agric. Food Chem 54: 4069-4075.
 24. De Pascual-Teresa S, Sanchez-Ballesta MT, Garcia-Viquera (2013) Anthocyanin. In. Ramawat KG &Merillo JM (eds) Natural Products. Springer-Verlag; Berlin Heidelberg.
 25. Jackman RL, Yada RY, Tung MA, Speers RA (1987) Anthocyanin as food colorants-a review. J Food Biochem 11: 201.
 26. Francis FJ (1989) Food colorants-Anthocyanin. Crit Rev Food SciNutr 28:273.
 27. Basu A, Rhone M, Lyons T (2010). 'Berries: emerging impact on cardiovascular health',Nutr Rev, Vol.68(3): 168-177.
 28. Buckle KA, et al (1978). Food Science (A course manual in food science). Australian Vice-Counsellors' Committee : Canberra.

Changes of Anthocyanin Contents, Radical Scavenging Activities and Total Microbial Counts of Buni Fruit (Antidesmabunius) Juice during Different Storage Times

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	jfoodpharmsci.com Internet Source	4%
2	Submitted to Universitas Hasanuddin Student Paper	1%
3	psasir.upm.edu.my Internet Source	1%
4	www.ifrj.upm.edu.my Internet Source	1%
5	www.lib.ncsu.edu Internet Source	1%
6	"Abstracts of the Asian Congress of Nutrition 2019", Annals of Nutrition and Metabolism, 2019 Publication	1%
7	eprints.arums.ac.ir Internet Source	1%
8	www.mdpi.com Internet Source	

1 %

9

link.springer.com

Internet Source

1 %

10

www.stikesayani.ac.id

Internet Source

1 %

11

Jin Dai. "Characterization of Blackberry Extract and Its Antiproliferative and Anti-Inflammatory Properties", *Journal of Medicinal Food*, 06/2007

Publication

1 %

12

G. Gradinaru, C.G. Biliaderis, S. Kallithraka, P. Kefalas, C. Garcia-Viguera. "Thermal stability of Hibiscus sabdariffa L. anthocyanins in solution and in solid state: effects of copigmentation and glass transition", *Food Chemistry*, 2003

Publication

<1 %

13

Henning, Susanne M., Yanjun Zhang, Victoria G. Rontoyanni et al. "Variability in the Antioxidant Activity of Dietary Supplements from Pomegranate, Milk Thistle, Green Tea, Grape Seed, Goji, and Acai: Effects of in Vitro Digestion", *Journal of Agricultural and Food Chemistry*

Publication

<1 %

- | | | |
|----|---|------|
| 14 | Kyu-Ho Han, Mitsuo Sekikawa, Ken-ichiro Shimada, Makoto Hashimoto et al.
"Anthocyanin-rich purple potato flake extract has antioxidant capacity and improves antioxidant potential in rats", British Journal of Nutrition, 2007
Publication | <1 % |
| 15 | arxiv.org
Internet Source | <1 % |
| 16 | dea.lib.unideb.hu
Internet Source | <1 % |
| 17 | hdl.handle.net
Internet Source | <1 % |
| 18 | www.coursehero.com
Internet Source | <1 % |
| 19 | "Natural Products", Springer Nature, 2013
Publication | <1 % |
| 20 | Claudia Drossard, Ute Alexy, Katja Bolzenius, Clemens Kunz, Mathilde Kersting.
"Anthocyanins in the diet of infants and toddlers: intake, sources and trends", European Journal of Nutrition, 2011
Publication | <1 % |
| 21 | Palayyan Saraswathy Benherlal. "Chemical composition and in vitro antioxidant studies | <1 % |

onSyzygium cumini fruit", Journal of the
Science of Food and Agriculture, 11/2007

Publication

22

S Marchellia, A Sutiono, J Lheman, Yanti, RR
Tjandrawinata. "Cosmeceutical potency of
functional ripe buni cider", IOP Conference
Series: Earth and Environmental Science, 2020

Publication

23

kb.psu.ac.th

Internet Source

24

www.scribd.com

Internet Source

25

Isti Ilmiati Fujiati, Harun Alrasyid Damanik,
Adang Bachtiar, Andi Armyn Nurdin, Paul
Ward. "Development and validation of
prediabetes risk score for predicting
prediabetes among Indonesian adults in
primary care: Cross-sectional diagnostic
study", Interventional Medicine and Applied
Science, 2017

Publication

26

S. Samappito, L. Butkhup. "An Analysis on
Flavonoids, Phenolics and Organic Acids
Contents in Brewed Red Wines of Both Non-
Skin Contact and Skin Contact Fermentation
Techniques of Mao Luang Ripe Fruits
(Antidesma bunius) Harvested From Phupan
Valley in Northeast Thailand", Pakistan Journal

<1 %

<1 %

<1 %

<1 %

<1 %

of Biological Sciences, 2008

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On