

Journal of Global Pharma Technology The Role of Lipopolysaccharide and Tight Junction Protein-1 (Zo- 1) Levels and Symbiosis of Black Rice Extract in Obese Patients

by Agussalim Buchari

Submission date: 16-Apr-2021 04:20AM (UTC+0700)

Submission ID: 1560333717

File name: Levels_and_Symbiosis_of_Black_Rice_Extract_in_Obese_Patients.pdf (345.18K)

Word count: 5042

Character count: 28498

30 See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/345386982>

Journal of Global Pharma Technology The Role of Lipopolysaccharide and Tight Junction Protein-1 (ZO-1) Levels and Symbiosis of Black Rice Extract in Obese Patients

Article in Journal of Global Pharma Technology - November 2020

CITATIONS
0


READS
21

5 authors, including:

 **Agussalim Bukhari**
Universitas Hasanuddin
39 PUBLICATIONS 920 CITATIONS
[SEE PROFILE](#)

 **Nurpudji A Taslim**
Universitas Hasanuddin
20 PUBLICATIONS 10 CITATIONS
[SEE PROFILE](#)

 **Aminuddin Aminuddin**
Faculty of Medicine University of Hasanuddin
13 PUBLICATIONS 295 CITATIONS
[SEE PROFILE](#)

 **Hasta Handayani Idrus**
Universitas Muslim Indonesia
34 PUBLICATIONS 17 CITATIONS
[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

 The relationship of mRNA IL23 expression to degree of hirschsprung associated enterocolitis [View project](#)

 Obesity [view project](#) 15

The Role of Lipopolysaccharide and Tight Junction Protein-1 (ZO-1) Levels and Symbiosis of Black Rice Extract in Obese Patients

Armanto Makmun¹, Agussalim Bukhari², Nurpudji A. Taslim², Aminuddin², Hasta Handayani Idrus^{1*}

¹Faculty of Medicine, Indonesian Muslim University, Makassar, Indonesia.

²Department of Nutrition, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.

*Corresponding Author: Hasta Handayani Idrus

Abstract

Excessive fat accumulation, which is commonly known as obesity, causes health problems for the community, such as metabolic disorders. Recent studies have shown that the gut microflora is involved in the occurrence of metabolic disorders in obesity. The gut micro biota can regulate body weight by influencing metabolic processes, regulating the expression of host genes, increasing the ability to extract and store energy. Intestinal micro biota imbalance and increased plasma lipopolysaccharide (LPS) contribute to inflammation in obesity. The latest safe approach in cases of obesity is to use active ingredients from local foods. In South Sulawesi, Indonesia and also in other Asian countries found black rice as the local product that is very good for health because the high level of bioactive compounds that are important for health such as anthocyanin, proteins, amino acids, and phenolic acids. The ingredients in black rice play a role in modulating the composition of the intestinal micro biota as indicated by the increasing composition of bifid bacterium and Lactobacillus. The increase in the composition of Bifid bacterium causes improvement in the expression of tight junction proteins in the intestinal epithelium, thereby contributing to prevent the translocation of LPS into the circulation. Apart from its role in modulating the composition of bacteria, it can also function to influence intracellular signaling.

Keywords: Obesity, Intestinal micro biota, LPS, tight junction, Black rice.

Introduction

The increase the prevalence of obesity that has occurred in the last few decades is related to lifestyle changes, namely the lack of physical activity and an increase in excessive consumption of foods containing high fat. Obesity is an abnormal or excessive fat accumulation that poses a risk to health. Obesity is considered to be a public health challenge whose prevalence continues to increase and underlies various metabolic and cardiovascular diseases [1]. The prevalence of obesity has spread out in most western countries over the past 30 years [2].

World Health Organization (WHO) data in 2016, more than 1.9 billion adults aged 18 years up are overweight; around 39% and 13% are from the world's population. The prevalence of obesity worldwide nearly triple

between 1975 and 2016 [3]. Obesity causes public health problems such as several chronic disorders, including type-2 diabetes, cardiovascular diseases (CVD), hypertension, musculoskeletal disorders (especially osteoarthritis), poor mental health, cancer, and causes negative effects on quality of life, work productivity, and health care costs [4].

New research shows that the gut micro biota is involved in obesity and metabolic disorders, with the decrease type of gut micro biota, the Bactericides / Firmicute ratio [5, 6]. The gut micro biota can regulate body weight by affecting the metabolic, neuroendocrine and immune functions of the host. The gut micro biota, as a whole, provides additional metabolic functions and regulates host gene expression, increases the ability to extract

and store energy while on a diet and contributes to weight gain. Imbalances in the gut micro biota and increase plasma lipopolysaccharides in obesity may also act as inflammatory factors associated with the development of atherosclerosis, insulin resistance and weight gain [7].

Given the impact of obesity due to excessive consumption of high calorie foods, various approaches to treating obesity have been widely researched so that being overweight obese can increase the risk of death. Aerobic training is the optimal mode of exercise for reducing fat mass. Resistance training is required to increase lean muscle mass in middle-aged and overweight / obese individuals. In addition, there is also cognitive behavioral therapy that directly deals with behaviors that require change in order to lose weight and maintain weight loss.

Pharmacological therapy can help patients to improve their risk of obesity-related diseases. The Food and Drug Administration (FDA) has approved certain drugs that can be used in the treatment of obesity for lifestyle changes. The drugs are Orlistat, Phentermine / Topiramate, Lorcaserin, Naltrexone / Bupropion, and Liraglutide. It should be noted that certain weight loss drugs have previously been withdrawn due to safety concerns and the drug effects of the various pharmacological agents used for the treatment of obesity [8-10].

The current approach that safe in treating the inflammatory process in obesity involves the use of active ingredients from local foods. Morais et al. finds that supplementing with anthocyanin-rich jussara powder can normalize weight gain, serum cholesterol, triglycerides, and restore gut bacteria Bifidobacterium and Lactobacillus in mice fed a trans fatty acid diet [11, 12].

Black rice (*Oryza sativa* L.) is a local food originating from South Sulawesi, Indonesia, and is also commonly found in Asia and exported to other countries. Consumption of black rice is in great demand because of the high level of bioactive compounds that are important for health [13].

The main bioactive compounds found in black rice include gallic, protocatechuic, hydroxybenzoic, vanilic acid, cyanidin 3-O-glucoside, peonidin-3-O-glucoside,

proanthocyanidin (C3G), flavanols, catechins, epicatechins, carotenoids, and γ -oryzanol [2]. Black rice, an anthocyanin-rich pigment rice, is widely consumed worldwide and is considered the most nutritious of all rice genotypes [13]. Many studies have shown that diets with black rice anthocyanin extract

weaken the development of obesity, hypercholesterolemia, liver steatosis, and atherosclerosis [14, 15]. Many studies have shown that these compounds can improve lipid profiles, anti-inflammatory, anti-cancer and reduce oxidative stress, can also treat heart disease, and prevent diabetes [16, 17]. Therefore, several studies have been interested in black rice in people with obesity and metabolic disorders.

Bioactive of Black Rice

Black rice is one type of rice plant with a distinctive color, black, more than 60% of the world's black rice is cultivated in Asian countries such as Indonesia, China, Thailand [18]. Black rice is characterized by its high anthocyanin content. The main anthocyanin is found in black rice include cyanidin 3-O-glucoside (6.3 mg / g) and peonidin-3-O-glucoside (3.6 mg / g).

Anthocyanidine, the basic structure of the anthocyanine, consists of an aromatic ring that is bound to a heterocyclic ring containing oxygen. This oxygen forms bonds with carbon bonds in the third aromatic ring. The structural characteristics of anthocyanins make it highly reactive with reactive oxygen species (ROS). The concentration of flavones and flavonols are also found higher in black rice compared to other rice variants.

Another study reported that the mean anthocyanin content is 35 times higher in black rice compared to other rice. In addition, the content of bioactive compounds, protein and essential amino acids that are higher than white rice. Amino acids play a major role in individual dietary factors to modify the risk of developing diabetes conditions. In addition, the anthocyanin content in black rice is reported to inhibit the levels of α -glucosidase and α -amylase enzymes, which are the main enzymes that are closely related to postprandial glucose levels.

Therefore black rice can be used in diabetes management. Anthocyanin from natural substances can also reach the colon and perform as candidates for nutritional therapy in gastrointestinal tract disorders. Anthocyanin in the intestine can be distorted as phenolic acids by ring cutting, dehydroxylation, and methylation reactions by the intestinal micro biota.

These phenolic compounds are proven to increase the growth of beneficial bacteria and inhibit the growth of bacteria that are bad for the body. Cyanidine-3-O-glucoside and β-glycoside provide energy for bacteria and increase prebiotic activity [20, 21]. Black rice is also rich in phenolic acids which are generally categorized into free phenolic acids and bound phenolic acids.

Free phenolic acids can be in the form of cinnamic, protocatechuic and gallic acids, while bound phenolic acids can be in the form of ferulic, coumaric and caffeic acids. Among free phenolic acids, protocatechuic acid is the most dominant (81-90% of the total free phenolic acid), while ferulic acid is the type of bound phenolic acid that is mostly found (about 60%) in black rice. Consumption of white rice has been associated with an increased risk of developing type 2 diabetes mellitus and an increased risk of impaired glucose homeostasis. However [22, 23].

Black Rice-Gut Micro-Biota

Black rice has high anthocyanin content. Anthocyanin is considered as substances with good health effects, such as antioxidants, anti-inflammatory, anti-cancer, anti-proliferative, and as substances that have the potential to affect the growth of susceptible microorganisms. Several in vitro studies revealed that the number of potentially beneficial bacteria (Bifidobacteria and Lactobacilli) increased after ingestion of purple sweet potato anthocyanin and grape seed extracts [18].

The gastrointestinal tract contains at least 10¹⁴ bacteria, with the highest density in the large intestine. The composition of the gastrointestinal tract micro biota is influenced by many factors including genetics, gender, age, height, weight, dietary factors, immune system, stool consistency, sleep, medical history, ethnogeographic and socioeconomic conditions, sanitary conditions, smoking habits, and use of antibiotics [24].

The intestinal microbiota functions as an anti-infection, immunomodulation and in the metabolic process. One of the main metabolic activities of the gastrointestinal tract microbiota is the production of non-gas short chain fatty acids (SCFA) [25]. These SCFAs can interact with the host via several pathways, thereby affecting important processes such as inflammation [26].

Like polyphenols, anthocyanin is difficult to absorb in the small intestine and can be converted into phenolic acid by the intestinal microbiota through edge ring, dehydroxylation and methylation reactions. The metabolism produced from polyphenols have proven that polyphenols can selectively motivate the growth of beneficial bacteria and inhibit the proliferation of harmful bacteria [26, 27]. Obesity is associated with phylum and specific changes in the gut microbiota, and reduce intestinal bacteria [28, 29].

The content in black rice is known to play a role in modulating the composition of the intestinal microbiota, indicated by increasing the composition of Bifidobacterium and Lactobacillus [18]. The increase in the composition of Bifidobacterium causes an improvement in the expression of tight junction proteins in the intestinal epithelium, thereby contributing to preventing translocation of lipopolysaccharides into the circulation.

Apart from playing a role in modulating the composition of bacteria, it can also work to influence intracellular signaling, works to inhibit the expression of MyD88 and NF-κB proteins, so that it can decrease the expression of proinflammatory cytokines induced by the bound between LPS and the complexes TLR4, CD14 and LBP [30, 31]. Anthocyanin content also specifically increases the micro biota population of Bifidobacterium spp. and Lactobacillus-Enterococcus spp [33].

Research conducted by Flores et al. (2015) shown that the 20 mg / L anthocyanin level is not significant enough to cause changes in the microbiota profile. Proliferative and inhibitory effects on the microbiota population can be observed in the administration of anthocyanins with higher levels, namely 200 mg / L. Another possible mechanism underlying the role of anthocyanins in the microbiota profile is that

anthocyanins can reduce oxygen tension in the intestinal lumen, thereby increasing the proliferation of oxygen-sensitive bacterial populations [34].

Black Rice-Lipopolysaccharide (LPS)

Obesity is caused by a high-fat diet and metabolic disorders characterized by inflammatory status associated with changes in the composition of the gut micro biota and increase plasma lipopolysaccharide (LPS) levels [68]. Anthocyanin content in LPS that cause changes in the immune system. In the research of Pornngarm et al. (2016) who test the anti-inflammatory effect of high polar fraction of black rice whole grain extracts (BR-WG-P) show that BR-WG-P significantly inhibit LPS-induced proinflammatory mediators, including nitric oxide (NO) production and iNOS expression, and COX-2.

In addition, the secretion of proinflammatory cytokines including TNF-α and IL-6 was also significantly inhibit [24]. BR-WG and anthocyanins inhibit translocation of NF-κB and AP-1 into the nucleus. BR-WG-P also decrease phosphorylation of ERK, p38 and JNK depending on the dose used [24].

These results suggest that BR-WG-P may suppress LPS-induced inflammation through inhibition of the MAPK signaling pathway leading to decrease translocation of NF-κB and AP-1 [36]. In the research of Sung-Won Min et al (2010), black rice suppresses the production of pro-inflammatory cytokines, TNF-α and IL-1β, and inflammatory mediators, NO and prostaglandin E2 (PGE2), as well as the expression of the nitric oxide synthase (iNOS) and cyclooxygenase genes. -2 (COX-2) in cell RAW 264.7 [37].

Black Rice-Tight Junction

Gut-derived bacterial LPS plays an important role in the intestinal, systemic inflammatory response and is involved in causing necrotizing enterocolitis and

inflammatory bowel disease. Interference with tight junctions in the intestine is an important factor in intestinal inflammation. Physiologically, LPS causes an increase in

the permeability of the intestinal tight junction through the TLR-4 dependent process [38]. LPS induces an increase in tight junction permeability which is mediated by an increase in enterocyte membrane expression TLR-4 and TLR-4 dependent on an increase in membrane colocalization of membrane associated protein CD14 [39]. Obesity is a condition associated with increased intestinal permeability and impaired intestinal protective function due to changes in the structure of the tight junction protein observed in an obesity experimental animal model [40].

In addition, there is increased intestinal permeability, higher plasma endotoxin levels and increased proinflammatory cytokines. A high-fat diet causes suppression of occludin, claudin-1, claudin-3, and JAM-1 levels, accompanied by an increase in plasma TNF-α levels in the intestinal mucosa [41]. Research by Lei Zhao et al. (2018) mice were given 100 mg / kg of black rice via oral gavage and given 2% DSS in drinking water for five days induce colitis in the group of mice given black rice showed a decrease in histological score, which indicates that there are less mucosal injury and edema compared to DSS treatment.

This study shows that anthocyanins can protect tight junctions by modulating the ratio of TJ-positive, negative protein and confirm the protective effect of anthocyanins from black rice in overcoming colonic inflammation. In addition, this study show a decrease in the expression levels of IL-6, IL-1β, TNF-α and myeloperoxidase (MPO) which linearly related to neutrophil infiltration. The anthocyanin content of purple tuberous root shows down-regulated TNF-α, IFN-γ, [40, 41].

Table 1: Black rice studies

Reference	Method	Result	Conclusion
Sung-Won Min et al. 2010(37)	Cyanidin-3-O-β-D-glycoside ((C3G) and cyanidin were isolated from Black Rice as previously reported by Han et al. (44). In addition, protocatechuic acid, RPMI 1640, penicillin-streptomycin, LPS purified from Escherichia coli O111: B4, and a radio-immunoprecipitation assay	The anti-inflammatory effect of C3G the main constituent of black rice, and its metabolites, cyaniding and protocatechuic acid (PA), were assessed in lipopolysaccharide (LPS) - RAW 264.7 cells induced and carrageenan-induced inflammation in air sacs in BALB / c mice. BR, C3G and their metabolites suppress the production of pro inflammatory cytokines, TNF-α and IL-	Based on these findings, when black rice is consumed orally, its main constituent, C3G, can be metabolized to cyaniding and / or PA, which expresses a potent anti-inflammatory effect by regulating the activation of NF-κB and MAPK.

	<p>65 PA) lysis buffer. Antibodies to COX-2, iNOS, TNF-α, IL-1β, β-actinwer. Antibodies for IκB-α, p-IκB-α, p65, p-p65, p38, p-p38, JNK, p-JNK, ERK, and p-ERK. T58 Cytokine ELISA Kit was obtained from R&D Systems. Carrageenan (Type IV). Bio-Rad Protein Assay Kit.</p>	<p>18, and inflammatory mediators, NO and prostaglandin E2 (PGE2), as well as gene expression of nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) in RAW 264.7 cells. This agent also 27 bits phosphorylation of IκB-α, nuclear translocation of NF-κB, and activation of mitogen-activated protein kinases. In addition, this agent sign27ntly inhibited the leucocyte count and levels of TNF-α, 1312, and protein in air-bag exudates in carrageenan-treated mice, as well as COX-2 expression and NF-κB activation. Among the test agents,</p>	
<p>Hao Wang et al. 2020(5)</p>	<p>BRAE, fecal, and urine anthocyanin7were analyzed by HPLC using column C18 (4.6 mm \times 250 mm, 5 μm23HI-MADZU, Japan), with A) water / formic acid (99: 1, v / v) and B) 100% acetonitrile as mobile phase monitored at 520 nm in the presence of a standard. Pathogen-free male 57BL6 / J mice (6 weeks old). C57BL / 6J mice were grouped into the normal chow diet (NCD) group, the high fat and cholesterol (HCD) diet group, and three treatment groups that were given HCD with various doses of black rice anthocyanin extract (BRAE) for 12 weeks. 39</p>	<p>BRAE reduces weight gain, serum triglycerides (TG), total cholesterol (TC), 5els of non-high-density lipoprotein (non-HDL-C) cholesterol, and increased excretion of fecal sterols and short-chain fatty acids. (SCFAs) concentrations in HCD-induced hypercholesterolemic rats. In addition, BRAE decrease the TC content of the liver through fundamental regulation of the body's energy balance gene, adenosine 5'-monophosphate activates protein kinase (AMPK). 14nwhile, BRAE increases the expression of genes involved in cholesterol uptake and excretion, and maintains CYP7A1, mRNA exp5sion of the 5/8 ATP-binding ATP, and the relative abundance of gut micro biota. Other than that 35</p>	<p>BRAE supplements can be a useful treatment option for preventing HCD-induced hypocholesterolemia and metabolic syndrome. 5</p>
<p>Lei Zhao et al. 2009(42)</p>	<p>Black rice (<i>Oryza sativa</i> L.) was purchased from a local market in Beijing, China. It was grounded to be fine powder by crushing and passes through a 2 sieve 60-mesh sieve. The black rice powder was t73 sealed in PET / Al / PE bags an55ored at 4 $^{\circ}$ C before use. The DSS (36-50 kDa, reagent class) was purchased from M12 Biomedicals. RA (\geq98%), cyanidin-3-O-β-glucoside (\geq98%), peonidin-3-O-β-glucoside (\geq98%), chlorogenic acid (\geq98%) 56 ferulic acid (\geq98 %) 8 purchased from Chengdu Must Bio-Technology Co., Ltd. . HPLC grade acetonitrile was purchased from Mallinckrodt Baker. The real time quantitative PCR primer comes from Beijing Genomics.</p>	<p>The administration of black rice anthocyanin-rich extract (BRAE) and rosmarinic acid (RA), alo11 and in combination, significantly reduce the disease activity index (DAI) and colonic histological score in colitis-induced DSS mice. In addition, administering BRAE and RA, apart and in 71bination, not only reduces levels of myeloperoxidase (MPO) and nitric oxide (NO), but 28 inhibits the expression of pro-inflammatory mediators including interleukin (IL) -6, IL-1β, tumor necrosis factor (TNF) -α, induced nitric oxide synthase (iNOS) and cyclooxygenase (COX) -2. Our results show th50BRAE decreases histologic score and TNF-α 2 mRNA expression in a dose-dependent manner, while BRAE + R74ttenuated histological score and IL-6 mRNA expression are dose-dependent 3 24. However,</p>	<p>The B7E and RA diets, apart and in combination, relieve the symptoms and inflammation of colitis due to DSS in mice, and ca7provide a promising dietary approach for the management of inflammatory bowel disease.</p>
<p>Yongsheng Zhu et al. (2018)(18)</p>	<p>Anthocyanins from black rice were obtained by membrane filtration and column chromatographic separation. 17 anthocyanin monomers in black rice extract were identified by HPLC-MS / MS, and the main anthocyanin monomer (cyanidin-3-glucoside, C3G) was purified by preparative HPLC (Pre-HPLC). The proliferative effect of anthocyanins on Bifidobacteria and Lactobacillus was investigated by determining the pH of the media, bacterial population and metabolic products. After anaerobic incubation at 37C for 48 hours,</p>	<p>Anthocyanins and anthocyanin monomers from black rice have prebiotic activity and are metabolized into several small molecules by Bifid bacteria and Lactobacillus.</p>	<p>Various phenolic acids are obtained after the metabolism of C3G bacteria. The modulating effect of C3G on the gut micro biota through in vivo fermentation provides strong evidence of the health benefits of the anthocyanin diet.</p>

	<p>not only was the pH of the 137 a containing C3G lower than black rice anthocyanin extract (BRAE), but the number of Bifid bacteria and Lactobacillus also increased significantly. Furthermore, hydroxyphenylpropionic acid, hydroxyphenylacetic acid, and hydroxybenzoate and other metabolites were detected by GC-MS in vitro.</p>		
--	--	--	--

Conclusion

Obesity affects almost all physiological functions of the body and it is significant on health problem. This increases the risk of several diseases such as metabolic disorders, diabetes mellitus, cardiovascular disease, cancer, a range of musculoskeletal disorders and poor mental health. Intestinal micro biota imbalance and increased plasma LPS

contribute to inflammation in obesity. Black rice has a high content of bioactive compounds such as anthocyanin, proteins, amino acids, and phenolic acids. The ingredients of black rice have role in modulating the composition of the intestinal micro biota, inhibiting LPS-induced pro-inflammatory mediators, can protect the tight junction so that it can protect inflammation in the colon.

References

- Liu X, Cervantes C, Liu F (2017) Common and distinct regulation of human and mouse brown and beige adipose tissues: a promising therapeutic target for obesity. *Protein Cell.*, 8(6):446-54.
- Zhang Y, Xie C, Wang H, Foss RM, Clare M, George EV, et al (2016) Irisin exerts dual effects on browning and adipogenesis of human white adipocytes. *Am J. Physiol - Endocrinol Metab.*, 311(2):E530-41.
- Organization WH (2018) Obesity and Overweight.
- Chooi YC, Ding C, Magkos F (2019) The epidemiology of obesity. *Metabolism*, 92: 6-10.
- Wang H, Liu D, Ji Y, Liu Y, Xu L, Guo Y (2020) Dietary Supplementation of Black Rice Anthocyanin Extract Regulates Cholesterol Metabolism and Improves Gut Microbiota Dysbiosis in C57BL / 6J Mice Fed a High-Fat and Cholesterol Diet. *Food Funct.*, 64(8):1-13.
- Zhang M, Yao J, Deng J, Duan Z (2019) Function saponin on intestinal health in antibiotic-treated. 4124-33.
- Sun L, Ma L, Ma Y, Zhang F, Zhao C, Nie Y (2018) Insights into the role of gut microbiota in obesity: pathogenesis, mechanisms, and therapeutic perspectives. *Protein Cell*, 9(5):397-403.
- Turner M, Jannah N, Kahan S, Gallagher C, Dietz W (2018) Current Knowledge of Obesity Treatment Guidelines by Health Care Professionals. *Obesity*, 26(4):665-71.
- Haslam DW, James WPT (2005) Obesity. *Lancet*, 366(9492):1197-209.
- Organization WH (2018) Obesity and overweight. In Switzerland: World Health Organization.
- Almeida C, Missae L, Moura R De, Vera V, Rosso D, Oller C, et al (2015) Polyphenols-rich fruit in maternal diet modulates inflammatory markers and the gut microbiota and improves colonic expression of ZO-1 in offspring. *Food Res Int.*, 77(2):186-93.
- Morais CA, Oyama LM, Oliveira JL De, Garcia MC, Rosso VV De, Sousa L, et al (2014) Jussara (*Euterpe edulis* Mart.) Supplementation during Pregnancy and Lactation Modulates the Gene and Protein Expression of Inflammation Biomarkers Induced by trans -Fatty Acids in the Colon of Offspring. *Mediat Inflamm.*, 2014(2014):987927.

- 13 Zhang MW, Zhang RF, Zhang FX, Liu RH (2010) Phenolic Profiles and Antioxidant Activity of Black Rice Bran of Different Commercially Available Varieties. *J. Agric. Food Chem.*, 58(13):7580-7.
- 14 Cremonini E, Daveri E, Mastaloudis A, Adamo AM, Mills D, Kalanetra K, et al (2019) Redox Biology Anthocyanins protect the gastrointestinal tract from high fat diet-induced alterations in redox signaling, barrier integrity and dysbiosis. *Redox Biol.*, 26(June):101-269.
- 15 Jang H, Park M, Kim H, Lee Y, Hwang K, Park J, et al (2012) Black rice (*Oryza sativa* L.) extract attenuates hepatic steatosis in C57BL/6 J mice fed a high-fat diet via fatty acid oxidation. *Nutr Metab.*, 9(27):1-11.
- 16 Chen PN, Kuo WH, Chiang CL, Chiou HL, Hsieh YS, Chu SC (2006) Black rice anthocyanins inhibit cancer cells invasion via repressions of MMPs and u-PA expression. *Chem Biol Interact.*, 163(3):218-29.
- 17 Prasad BJ, Sharavanan PS, Sivaraj R (2019) Health benefits of black rice -A review. *Grain Oil Sci. Technol.*, 2(4):109-13.
- 18 Zhu Y, Sun H, He S, Lou Q, Yu M, Tang M, et al (2018) Metabolism and prebiotics activity of anthocyanins from black rice (*Oryza sativa* L.) in vitro. *PLoS One*, 13: 29
- 19 Javadzadeh Y, Hamedeyaz S (2014) Floating Drug Delivery Systems for Eradication of *Helicobacter pylori* in Treatment of Peptic Ulcer Disease. *Trends Helicobacter pylori Infect.*, 304-7.
- 20 Kang MY, Kim JH, Rico CW, Nam SH (2011) A comparative study on the physicochemical characteristics of black rice varieties. *Int. J. Food Prop.*, 14(6):1241-54.
- 21 Mirmiran P, Bahadoran Z, Esfandyari S, Azizi F (2017) Dietary protein and amino acid profiles in relation to risk of dysglycemia: Findings from a prospective population-based study. *Nutrients*, 9(9):1-9.
- 22 Sumczynski D, Kotásková E, Družbiková H, Mlček J (2016) Determination of contents and antioxidant activity of free and bound phenolics compounds and in vitro digestibility of commercial black and red rice (*Oryza sativa* L.) varieties. *Food Chem.*, 211: 339-46.
- 23 Ito VC, Lacerda LG (2019) Black rice (*Oryza sativa* L.): A review of its historical aspects, chemical composition, nutritional and functional properties, and applications and processing technologies. *Food Chem.*, 301(April):125-304.
- 24 Idrus HH, Hatta M, Kasim VN, Achmad AF, Mangarengi Y (2020) Molecular Impact on High Motility Group Box-1 (HMGB-1) in Pamps and Damp. *Indian J. Public Health.*, 11(1):1-8.
- 25 Boerner BP, Sarvetnick NE (2011) Type 1 diabetes: Role of intestinal microbiome in humans and mice. *Ann N Y Acad. Sci.*, 1243(1):103-18.
- 26 Wang LL, Guo HH, Huang S, Feng CL, Han YX, Jiang JD (2017) Comprehensive evaluation of SCFA production in the intestinal bacteria regulated by berberine using gas-chromatography combined with polymerase chain reaction. *J. Chromatogr. B Anal. Technol. Biomed Life Sci.*, 1057(May):70-80.
- 27 Cueva C, Moreno-Arribas MV, Martín-Álvarez PJ, Bills G, Vicente MF, Basilio A, et al (2010) Antimicrobial activity of phenolic acids against commensal, probiotic and pathogenic bacteria. *Res Microbiol.*, 161(5):372-82.
- 28 Lee HC, Jenner AM, Low CS, Lee YK (2006) Effect of tea phenolics and their aromatic fecal bacterial metabolites on intestinal microbiota. *Res Microbiol.*, 157(9):876-84.
- 29 Waldram A, Holmes E, Wang Y, Rantalainen M, Wilson ID, Tuohy KM, et al (2009) Top-down systems biology modeling of host metabotype-microbiome associations in obese rodents. *J. Proteome Res.*, 8(5):2361-75.
- 30 Turnbaugh PJ, Hamady M, Yatsunenkov T, Cantarel BL, Ley RE, Sogin ML, et al (2009) A core gut microbiome between lean and obesity twins. *Nature*, 457(7228):480-4.
- 31 Lu YC, Yeh WC, Ohashi PS (2008) LPS/TLR4 signal transduction pathway. *Cytokine*, 42(2):145-51.
- 32 Villena J, Kitazawa H (2013) Modulation of intestinal TLR4-inflammatory signaling pathways by probiotic

- microorganisms: Lessons learned from *Lactobacillus jensenii* TL2937. *Front Immunol.*, 4(DEC):1-13.
- 33 Saez-Lara MJ, Gomez-Llorente C, Plaza-Diaz J, Gil A (2015) The role of probiotic lactic acid bacteria and bifidobacteria in the prevention and treatment of inflammatory bowel disease and other related diseases: A systematic review of randomized human clinical trials. *Biomed Res Int.*, 2015.
- 34 Flores G, Luisa M, Costabile A, Klee A, Bigetti K, Gibson GR (2015) In vitro fermentation of anthocyanins encapsulated with cyclodextrins: Release, metabolism and influence on gut microbiota growth. *J. Funct. Foods*, 16(2015):50-7.
- 35 Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, et al (2007) Metabolic Endotoxemia Initiates Obesity and Insulin Resistance. *Diabetes*, 56(July):1761-72.
- 36 Limtrakul P, Yodkeeree S, Pitchakarn P, Punfa W (2015) Suppression of Inflammatory Responses by Black Rice Extract in RAW 264.7 Macrophage Cells via Downregulation of NF- κ B and AP-1 Signaling Pathways, 16:4277-83.
- 37 Min SW, Ryu SN, Kim DH (2010) Anti-inflammatory effects of black rice, cyanidin-3-O- β -D-glucoside, and its metabolites, cyanidin and protocatechuic acid. *Int Immunopharmacol.*, 10(8):959-66.
- 38 Guo S, Nighot M, Al-sadi R, Alhmod T, Nighot P, Affairs V (2016) Lipopolysaccharide regulation of intestinal tight junction permeability is mediated by TLR-4 signal transduction pathway activation of FAK and MyD88. *J. Immunol.*, 195(10):4999-5010.
- 39 Guo S, Al-sadi R, Said HM, Ma TY (2013) Lipopolysaccharide Causes an Increase in Intestinal Tight Junction Permeability in Vitro and in Vivo by Inducing Enterocyte Membrane Expression and Localization of. *Am J. Pathol.*, 182(2):375-87.
- 40 Idrus HH, Mangarengi Y, Mustajar NS (2018) Test of Polymerase Chain Reaction (PCR) Detection and The Specificity in Gen Hd *Salmonella typhi* in RS. Ibnu Sina. In: The 8th Annual Basic Science International Conference "Convergence of Basic Sciences, Toward the World's Sustainability Challenges." 1.
- 41 Ma TY, Boivin MA, Ye D, Pedram A, Said HM, Thomas Y, et al (2005) Mechanism of TNF-intestinal epithelial tight junction barrier: role of myosin light-chain kinase protein expression, 0001: 422-30.
- 42 Zhao L, Zhang Y, Liu G, Hao S, Wang C, Wang Y (2018) Black rice anthocyanin-rich extract and rosmarinic acid, alone and in combination, protect against DSS-induced colitis in mice. *Food Funct.*, 9(5):2796-808.
- 43 Chen T, Hu S, Zhang H, Guan Q (2017) Anti-inflammatory effects of *Dioscorea alata* L. anthocyanins in a TNBS-induced colitis model. *Food Funct.*, 8(2):659-69.
- 44 Han SJ, Ryu S, Trinh HT, Joh EH, Jang SY, Han MJ, et al (2009) Metabolism of cyanidin-3-o- β -d-glucoside isolated from black colored rice and its antiscratching behavioral effect in mice. *J. Food Sci.*, 74: 8.

Journal of Global Pharma Technology The Role of Lipopolysaccharide and Tight Junction Protein-1 (Zo- 1) Levels and Symbiosis of Black Rice Extract in Obese Patients

ORIGINALITY REPORT

42%
SIMILARITY INDEX

30%
INTERNET SOURCES

36%
PUBLICATIONS

12%
STUDENT PAPERS

PRIMARY SOURCES

1 Yolanda Sanz, Arlette Santacruz, Paola Gauffin. "Gut microbiota in obesity and metabolic disorders", Proceedings of the Nutrition Society, 2010 **2%**
Publication

2 res.mdpi.com **2%**
Internet Source

3 library.cmu.ac.th **2%**
Internet Source

4 www.intechopen.com **2%**
Internet Source

5 Xu Si, Jinlong Tian, Chi Shu, Yuehua Wang, Ersheng Gong, Ye Zhang, Weijia Zhang, Huijun Cui, Bin Li. "Serum Ceramide Reduction by Blueberry Anthocyanin-Rich Extract Alleviates Insulin Resistance in Hyperlipidemia Mice", Journal of Agricultural and Food Chemistry, 2020 **2%**
Publication

6	"Plant Biotechnology: Recent Advancements and Developments", Springer Science and Business Media LLC, 2017 Publication	2%
7	www.tandfonline.com Internet Source	1%
8	repository.ubaya.ac.id Internet Source	1%
9	coek.info Internet Source	1%
10	Filomena Nazzaro, Florinda Fratianni, Vincenzo De Feo, Alberto Battistelli, Adriano Gomes Da Cruz, Raffaele Coppola. "Polyphenols, the new frontiers of prebiotics", Elsevier BV, 2020 Publication	1%
11	www.science.gov Internet Source	1%
12	Black Rice, 2016. Publication	1%
13	Iramaia Angélica Neri-Numa, Cinthia Baú Betim Cazarin, Ana Lúcia Tasca Gois Ruiz, Bruno Nicolau Paulino et al. "Targeting flavonoids on modulation of metabolic syndrome", Journal of Functional Foods, 2020 Publication	1%

- | | | |
|----|--|-----|
| 14 | www.hindawi.com
Internet Source | 1 % |
| 15 | Submitted to Universiti Teknologi MARA
Student Paper | 1 % |
| 16 | Submitted to Universitas Airlangga
Student Paper | 1 % |
| 17 | Tingting Qiu, Yong Sun, Xiaoya Wang, Liufeng Zheng, Hua Zhang, Li Jiang, Xuemei Zhu, Hua Xiong. "Drum drying-and extrusion-black rice anthocyanins exert anti-inflammatory effects via suppression of the NF- κ B /MAPKs signaling pathways in LPS-induced RAW 264.7 cells", Food Bioscience, 2021
Publication | 1 % |
| 18 | www.frontiersin.org
Internet Source | 1 % |
| 19 | www.scribd.com
Internet Source | 1 % |
| 20 | Vivian Cristina Ito, Luiz Gustavo Lacerda. "Black rice (<i>Oryza sativa</i> L.): A review of its historical aspects, chemical composition, nutritional and functional properties, and applications and processing technologies", Food Chemistry, 2019
Publication | 1 % |
| 21 | www.cambridge.org | |

Internet Source

1 %

22

Submitted to Western Governors University

Student Paper

1 %

23

Eun-Ha Joh, Jin-Ju Jeong, Dong-Hyun Kim.
"Kalopanaxsaponin B inhibits LPS-induced
inflammation by inhibiting IRAK1 Kinase",
Cellular Immunology, 2012

Publication

1 %

24

Submitted to University of Queensland

Student Paper

1 %

25

wellnesslabinfo.blogspot.com

Internet Source

1 %

26

www.ncbi.nlm.nih.gov

Internet Source

1 %

27

onlinelibrary.wiley.com

Internet Source

1 %

28

www.degruyter.com

Internet Source

1 %

29

Submitted to Queen Mary and Westfield
College

Student Paper

1 %

30

Submitted to Pennsylvania State System of
Higher Education

Student Paper

<1 %

31 Antonio Speciale, Antonella Saija, Romina Bashllari, Maria Sofia Molonia et al. "Anthocyanins As Modulators of Cell Redox-Dependent Pathways in Non-Communicable Diseases", Current Medicinal Chemistry, 2020
Publication <1 %

32 Eun-Ha Joh. "Lancemaside a inhibits lipopolysaccharide-induced inflammation by targeting LPS/TLR4 complex", Journal of Cellular Biochemistry, 2010
Publication <1 %

33 apnews.com
Internet Source <1 %

34 krishikosh.egranth.ac.in
Internet Source <1 %

35 www.semanticscholar.org
Internet Source <1 %

36 E. O. Igwe, K. E. Charlton, Y. C. Probst, K. Kent, M. E. Netzel. "A systematic literature review of the effect of anthocyanins on gut microbiota populations", Journal of Human Nutrition and Dietetics, 2018
Publication <1 %

37 Submitted to University of Bedfordshire
Student Paper <1 %

38

Balasubramaniam Jaya Prasad, Pazhaniyandi Subramania Sharavanan, Rengaraj Sivaraj. "Health benefits of black rice – A review", Grain & Oil Science and Technology, 2019

Publication

<1 %

39

Jared Gabriel Lopez DELA ROSA, Paul Mark Baco MEDINA. "Philippine rice wine (Tapuy) made from Ballatinao black rice and traditional starter culture (Bubod) showed high alcohol content, total phenolic content, and antioxidant activity", Food Science and Technology, 2021

Publication

<1 %

40

Mendy Hatibie Oley, Maximillian Christian Oley, Lily Lucia Loho, Dicky Panduwinata, Deanette Michelle R. Aling, Muhammad Faruk. "Trichilemmal carcinoma in an unexposed area of the nose: A case report", International Journal of Surgery Case Reports, 2021

Publication

<1 %

41

Shiyu Li, Binning Wu, Wenyi Fu, Lavanya Reddivari. "The Anti-inflammatory Effects of Dietary Anthocyanins against Ulcerative Colitis", International Journal of Molecular Sciences, 2019

Publication

<1 %

42

Shuang Xie, Shiyi Zhao, Lan Jiang, Linhao Lu, Qian Yang, Qinghua Yu. " Stimulates Intestinal

<1 %

Epithelial Proliferation and Induces
Differentiation into Goblet Cells in Young
Chickens ", Journal of Agricultural and Food
Chemistry, 2019

Publication

43

s3.amazonaws.com

Internet Source

<1 %

44

koreascience.or.kr

Internet Source

<1 %

45

www.biomedcentral.com

Internet Source

<1 %

46

repository.umy.ac.id

Internet Source

<1 %

47

www.biorxiv.org

Internet Source

<1 %

48

Fengfeng Wu, Na Yang, Alhassane Touré,
Zhengyu Jin, Xueming Xu. "Germinated Brown
Rice and Its Role in Human Health", Critical
Reviews in Food Science and Nutrition, 2013

Publication

<1 %

49

Hui Zhang, Guoyin Kai, Yongjun Xia,
Guangqiang Wang, Lianzhong Ai. "Antioxidant
and in vitro digestion property of black rice
(Oryza sativa L.): a comparison study between
whole grain and rice bran", International
Journal of Food Engineering, 2020

Publication

<1 %

- 50 Jie Ren, Sung H. Chung. "Anti-inflammatory Effect of α -Linolenic Acid and Its Mode of Action through the Inhibition of Nitric Oxide Production and Inducible Nitric Oxide Synthase Gene Expression via NF- κ B and Mitogen-Activated Protein Kinase Pathways", *Journal of Agricultural and Food Chemistry*, 2007
Publication <1 %
-
- 51 link.springer.com
Internet Source <1 %
-
- 52 scielo.conicyt.cl
Internet Source <1 %
-
- 53 www.sochob.cl
Internet Source <1 %
-
- 54 Aécio L. de S. Dias, Barbara Pachikian, Yvan Larondelle, Joëlle Quetin-Leclercq. "Recent advances on bioactivities of black rice", *Current Opinion in Clinical Nutrition & Metabolic Care*, 2017
Publication <1 %
-
- 55 Beita Zhao, Bing Xia, Xiaohan Li, Li Zhang, Xiaoning Liu, Renjie Shi, Rongwei Kou, Zhigang Liu, Xuebo Liu. "Sesamol Supplementation Attenuates DSS-Induced Colitis via Mediating Gut Barrier Integrity, Inflammatory Responses, and Reshaping Gut Microbiome", <1 %

56

Hongying Qin, Guangming Zhou, Guilong Peng, Junping Li, Junhua Chen. "Application of Ionic Liquid-Based Ultrasound-Assisted Extraction of Five Phenolic Compounds from Fig (*Ficus carica* L.) for HPLC-UV", Food Analytical Methods, 2014

Publication

<1 %

57

Jiří Pospíšil, Daniela Konrádová, Miroslav Strnad. "Chapter 3 Antileishmanial Activity of Lignans, Neolignans, and Other Plant Phenols", Springer Science and Business Media LLC, 2021

Publication

<1 %

58

Sung-Won Min, Young-Jun Park, Dong-Hyun Kim. "Kakkalide and Its Metabolite Irisolidone Ameliorate Carrageenan-Induced Inflammation in Mice by Inhibiting NF-κB Pathway", Inflammation, 2010

Publication

<1 %

59

Y Hamanaka. "Expression of human beta-defensin 2 (hBD-2) in *Helicobacter pylori* induced gastritis: antibacterial effect of hBD-2 against *Helicobacter pylori*", Gut, 2001

Publication

<1 %

60

bmcchem.biomedcentral.com

Internet Source

<1 %

61

bpspubs.pericles-prod.literatumonline.com

Internet Source

<1 %

62

go.gale.com

Internet Source

<1 %

63

worldwidescience.org

Internet Source

<1 %

64

"Obesity and Diabetes", Springer Science and Business Media LLC, 2020

Publication

<1 %

65

Lee, In-Ah, Young-Jun Park, Eun-Ha Joh, and Dong-Hyun Kim. "Soyasaponin Ab Ameliorates Colitis by Inhibiting the Binding of Lipopolysaccharide (LPS) to Toll-like Receptor (TLR)4 on Macrophages", Journal of Agricultural and Food Chemistry, 2011.

Publication

<1 %

66

Yange Wei, Haixia Yang, Chenhui Zhu, Jianjun Deng, Daidi Fan. "Hypoglycemic Effect of Ginsenoside Rg5 Mediated Partly by Modulating Gut Microbiota Dysbiosis in Diabetic db/db Mice", Journal of Agricultural and Food Chemistry, 2020

Publication

<1 %

67

www.nursingce.com

Internet Source

<1 %

68

"Biomarkers of Human Aging", Springer Science and Business Media LLC, 2019

Publication

<1 %

69

"Natural Products for Cancer Chemoprevention", Springer Science and Business Media LLC, 2020

Publication

<1 %

70

Dietary Components and Immune Function, 2010.

Publication

<1 %

71

K.V. Surangi Dharmawansa, David W. Hoskin, H. P. Vasantha Rupasinghe.

"Chemopreventive Effect of Dietary Anthocyanins against Gastrointestinal Cancers: A Review of Recent Advances and Perspectives", International Journal of Molecular Sciences, 2020

Publication

<1 %

72

Veronica Lazar, Lia-Mara Ditu, Gratiela G. Pircalabioru, Ariana Picu, Laura Petcu, Natalia Cucu, Mariana Carmen Chifiriuc. "Gut Microbiota, Host Organism, and Diet Triologue in Diabetes and Obesity", Frontiers in Nutrition, 2019

Publication

<1 %

73

Yawadio, R.. "Identification of phenolic compounds isolated from pigmented rices

<1 %

and their aldose reductase inhibitory activities", Food Chemistry, 2007

Publication

74

You-Chang Oh, Won-Kyung Cho, Yun Hee Jeong, Ga Young Im, Min Cheol Yang, Youn-Hwan Hwang, Jin Yeul Ma. "Anti-Inflammatory Effect of Citrus Unshiu Peel in LPS-Stimulated RAW 264.7 Macrophage Cells", The American Journal of Chinese Medicine, 2012

Publication

<1 %

75

journals.usamvcluj.ro

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On