

Minocycline improves recognition memory and attenuates microglial activation in Gunn rat: A possible hyperbilirubinemia-induced animal model of schizophrenia

by Erlyn Limoa

FILE	YCLINE_IMPROVES_RECOGNITION_MEMORY_AND_ATTENUATES_MICROGLIAL.PDF (1.32M)		
TIME SUBMITTED	06-JAN-2020 09:23AM (UTC+0700)	WORD COUNT	7067
SUBMISSION ID	1239424360	CHARACTER COUNT	38327



9
Minocycline improves recognition memory and attenuates microglial
activation in Gunn rat: A possible hyperbilirubinemia-induced animal
model of schizophrenia



44
Kristian Liary^{a,d}, Tsuyoshi Miyaoka^{a,*}, Toshiko Tsumori^{b,c}, Motohide Furuya^a, Sadayuki Hashioka^a,
Rei Wake^a, Keiko Tsuchie^a, Michiyo Fukushima^a, Erlyn Limoa^{a,d}, Andi Jayalangka Tanra^d, Jun Horiguchi^a

43
^a Department of Psychiatry, Shimane University Faculty of Medicine, 89-1 Enya-cho, Izumo 693-8501, Japan

^b Department of Anatomical Neuroscience, Shimane University Faculty of Medicine, 89-1 Enya-cho, Izumo 693-8501, Japan

^c Department of Nursing, Faculty of Health and Welfare, Prefectural University of Hiroshima, 1-1 38-en-cho, Mihara 723-0053, Japan

^d Department of Psychiatry, Hasanudin University Faculty of Medicine, Jl. Perintis Kemerdekaan Km. 10, Makassar 90245, South Sulawesi, Indonesia

ARTICLE INFO

Article history:

Received 11 September 2013

Received in revised form 17 December 2013

Accepted 18 December 2013

Available online 2 January 2014

Keywords:

Activated microglia

Behavioral test

Gunn rat

Minocycline

Schizophrenia

Unconjugated bilirubin

ABSTRACT

81
Background: Accumulating evidence indicates that neuroinflammation plays a significant role in the pathophysiology of schizophrenia. We previously reported evidence of schizophreni-a-like behaviors and microglial activation in Gunn rats. We concluded that the Gunn rat, which exhibits a high concentration of unconjugated bilirubin, may be useful as an animal model of schizophrenia. On the other hand, there have been numerous reports that minocycline is effective in treating schizophrenia.

Methods: In the present study, we investigated the effects of minocycline on performance of behavioral tests (prepulse inhibition (PPI) and novel object recognition test (NORT)) after animals received either 40 mg/kg/d of minocycline or vehicle by intraperitoneal (i.p.) injection for 14 consecutive days. Furthermore, we examined the effects of minocycline on microglial activation in the hippocampal dentate gyrus of Gunn rats and Wistar rats. **Results:** We found that administration of minocycline for 14 days significantly increased the exploratory preference in retention sessions and tended to improve the PPI deficits in Gunn rats. Immunohistochemistry analysis revealed that microglial cells in the minocycline-treated Gunn rat group showed less expression of CD11b compared to vehicle-treated Gunn and Wistar groups.

Conclusion: Our findings suggest that minocycline improves recognition memory and attenuates microglial activation in the hippocampal dentate gyrus of Gunn rats. Therefore, minocycline may be a potential therapeutic drug for schizophrenia.

© 2013 Published by Elsevier Inc.

1. Introduction

37
Schizophrenia is one of the most mysterious and costliest mental disorders in terms of human suffering and societal expenditures (van Os and Kapur, 2009). Despite extensive research and remarkable

Abbreviations: UCB, unconjugated bilirubin; PPI, prepulse inhibition; NORT, novel object recognition test; i.p., intraperitoneal; Iba1, ionized calcium binding adaptor molecule 1; Iba2, integrin alpha M; CD11b, cluster of differentiation molecule 11b; DG, dentate gyrus; SGZ, subgranular zone; GL, granular layer; ML, molecular layer; PB, phosphate buffer; PBS, phosphate buffer saline; ABC, avidin–biotin peroxidase complex; ANOVA, analysis of variance.

* Corresponding author at: Department of Psychiatry, Shimane University Faculty of Medicine, 89-1 Enya-cho, Izumo City 693-8501, Shimane, Japan. Tel.: +81 853 20 2262; fax: +81 853 20 2260.

E-mail addresses: krizbox@hotmail.com (K. Liary), miyami@med.shimane-u.ac.jp (T. Miyaoka), t-tsumori@pu-hiroshima.ac.jp (T. Tsumori), pancho@med.shimane-u.ac.jp (M. Furuya), hashioka@med.shimane-u.ac.jp (S. Hashioka), reiwake@med.shimane-u.ac.jp (R. Wake), keits@med.shimane-u.ac.jp (K. Tsuchie), mtaki@med.shimane-u.ac.jp (M. Fukushima), erlynliem@yahoo.com (E. Limoa), ajtanra@yahoo.com (A.J. Tanra), jhoriguchi@med.shimane-u.ac.jp (J. Horiguchi).

0278-5846/\$ – see front matter © 2013 Published by Elsevier Inc.
<http://dx.doi.org/10.1016/j.pnpbp.2013.12.017>

30
advances in the neurobiological, neurochemical and genetic aspects of this disabling mental illness, the underlying etiological processes remain a challenge for clinicians and basic researchers alike (Meyer-Lindenberg et al., 2013). Moreover, this imperfect understanding of the etiology has been associated with poor treatment outcomes for many individuals (Dean et al., 2012). Therefore, a novel insight into the underlying mechanism of this illness is an urge to allow new treatment targets to be explored.

23
An accumulating body of evidence points to the significant role of neuroinflammation and the immune system in the pathophysiology of schizophrenia (Takahashi and Sakurai, 2013). There are also numerous reports that support the hypothesis that inflammation is a risk factor for developing schizophrenia later in life (Kneeland and Fatemi, 2013; Miller et al., 1988; Miller et al., 2013). Moreover, evidence from genomic (Ripke et al., 2011; Schwab et al., 2002), blood (Erbagci et al., 2001; Miller et al., 2011), postmortem (Fisman, 1975; Radewicz et al., 2000; Steiner et al., 2008), and in vivo imaging (Doorduyn et al., 2009; van Berckel et al., 2008) studies are leading toward a greater consensus that immune activation is involved in the pathophysiology of

schizophrenia. Interestingly, some anti-psychotics may have anti-inflammatory effects (Bian et al., 2008; Kato et al., 2007) and anti-inflammatory agents may have anti-psychotic effects (Laan et al., 2010; Muller and Schwarz, 2008).

Minocycline is a second generation of the antibiotic, tetracycline, and is one of the most promising neuroprotective and anti-inflammatory agents currently in clinical and experimental trials (Miyaoaka, 2012). Our previous clinical studies showed the potential of minocycline as an adjunctive treatment in schizophrenia patients (Miyaoaka et al., 2007, 2008). Similar results were also found in various animal models of psychosis or schizophrenia. Minocycline was reported to attenuate behavioral impairment as well as neurotoxicity after administration of methamphetamine (Mizoguchi et al., 2008; Zhang et al., 2006a), methylenedioxymethamphetamine (MDMA) (Zhang et al., 2006b), N-methyl-D-aspartate (NMDA) receptor antagonist dizocilpine (Levkovitz et al., 2007; Zhang et al., 2007), and viral mimetic polyriboinosinic-polyribocytidilic acid (Poly(I:C)) (Juckel et al., 2011).

Our previous studies generated Gunn rat (Gunn, 1944) as a possible animal model of schizophrenia. First, we reported behavioral abnormalities, deficits in prepulse inhibition (PPI), and neuropathological changes in Gunn rats that are similar to the characteristics of schizophrenia (Hayashida et al., 2009). Then, we demonstrated evidence of activated microglial cells in the hippocampal dentate gyrus (DG) of Gunn rats (Liury et al., 2012). Furthermore, anti-psychotic medications affected Gunn rat behaviors in a way similar to their effect on schizophrenic behaviors (Tsuchie et al., 2013). We proposed that hyperbilirubinemia (a high level of unconjugated bilirubin, UCB) leads to chronic neuroinflammation and has a pathogenic effect on the development of the brain and concluded that the Gunn rat may be used as an animal model of schizophrenia.

In the present study, we sought to investigate the effects of minocycline on Gunn rat, a possible hyperbilirubinemia-induced animal model of schizophrenia. We hypothesized that minocycline may improve behavioral disturbances and attenuate microglial activation in Gunn rats. First, we performed behavior tests (prepulse inhibition (PPI) and novel object recognition test (NORT)). Then, using immunohistochemistry analysis, we examined the microglial activation in hippocampal DG of Gunn rats.

2. Methods

2.1. Animals

The animals were male homozygous (j/j) Gunn rats (6 weeks old, 160–200 g body weight) and male Wistar rats (6 weeks old, 200–240 g body weight) at the beginning of the experiment (Japan SLC, Inc., Shizuoka, Japan). Animals were housed under standard conditions with a room temperature of $23 \pm 2^\circ\text{C}$, humidity $55 \pm 5\%$, and 12 h light/12 h dark cycle (light phase 7:00 to 19:00) with free access to food and water. All procedures were performed with the approval of the Shimane University Animal Ethics Committee, under the guidelines of the National Health and Medical Research Council of Japan.

2.2. Drug administration

Minocycline hydrochloride (Sigma-Aldrich, St. Louis, Mo., USA) was freshly prepared as a stock solution of 5 mg/ml (10 mM) in filtered PBS and stored frozen at -80°C every 7 days. We adjusted the acidic pH (pH 4.0) of minocycline hydrochloride to neutrality by adding sodium hydroxide as described by (Ferretti et al., 2012). As the vehicle, we used the filtered PBS, pH 7.4.

Animals were divided into four groups: Wistar + vehicle (WV) group, Wistar + minocycline (WM) group, Gunn + vehicle (GV) group, and Gunn + minocycline (GM) group. Each animal received either 40 mg/kg/d of minocycline or vehicle by intraperitoneal (i.p.)

injection for 14 consecutive days. The dose (40 mg/kg) of minocycline was selected based on the fact that this dose was effective in improving the methamphetamine-induced impairment of recognition memory (Mizoguchi et al., 2008) as well as prepulse inhibition deficits after the administration of the NMDA receptor antagonist dizocilpine (Zhang et al., 2007). To avoid stress or pain due to the intraperitoneal injection, animals were anesthetized by putting them into a halothane inhalation chamber for less than 1 min before the daily injection.

2.3. Prepulse inhibition (PPI) of startle response

A startle response system was used (SR-LAB, San Diego Instrument) as reported previously (Hayashida et al., 2009). Briefly, each rat was placed in a Plexiglass cylinder where it was exposed to white background noise at 65 dB for a 5 min acclimatization period. This was followed by four types of trials: (1) pulse alone, consisting of a 20 ms burst of white noise at 120 dB; (2) a 20 ms burst of white noise at 24 dB followed by a 20 ms white noise at 120 dB (70 PP + P); (3) a 20 ms burst of white noise at 80 dB followed by a 20 ms white noise at 120 dB (80 PP + P); and (4) background noise only (no stimulus). Our pre-experiment (unpublished data) found that Gunn rat showed a sensitive response and behavior, therefore the use of three or four different prepulse intensities affected the behavior test and biased the result. In consequence, we decided to use just 2 prepulse intensities (70 dB and 80 dB). The interval between prepulse and pulse was set at 100 ms. Trials were given in a pseudo-random order with variable intervals (20 s–60 s, average 40 s) between each trial. Startle response was measured in sessions consisting of 50 trials. The percentage of PPI (%PPI) was defined as the magnitude of inhibition due to the startle amplitude that was induced by the prepulse. $\%PPI = (1 - (\text{startle magnitude after prepulse-pulse pair} / \text{startle magnitude after pulse-only})) \times 100$.

2.4. Novel object recognition test (NORT)

The NORT procedure consisted of three sessions: habituation, training, and retention. The experimental apparatus consisted of a Plexiglass open field box (42 × 42 × 42 cm) with a dust-covered floor located in a sound attenuated room. Each animal was individually habituated to the box, with 15 min of exploration in the absence of objects for two consecutive days (habituation session, days 1–2). During the training session (day 3), two identical objects were symmetrically fixed to the floor of the box, 5 cm from the walls, and each animal was allowed to explore freely for 2 × 5 min. An animal was considered to be exploring the object when its head was facing the object (a distance between the head and object of approximately 1 cm or less) or was touching or sniffing the object. In the retention session (day 4), one of the familiar objects used during the training session had been replaced with a novel object. The novel object was different in shape and color, but similar in size to the familiar object. The animal was then allowed to explore freely for 5 min. All sessions were videotaped for later examination. The exploratory preference index in the retention session, a ratio of the amount of time spent exploring the novel object over the total time exploring both objects, was used to measure cognitive function. In the training session, the exploratory preference index was calculated as the ratio of time spent exploring the object that was replaced by the novel object in the retention session over the total exploration time.

2.5. Immunohistochemistry for light and confocal laser scanning microscopy

Soon after finishing the behavior test, animals underwent deep intraperitoneal anesthesia with sodium pentobarbital (80 mg/kg body weight) and were perfused transcardially with 500 ml of physiological

saline, followed by 10 ml of 4% paraformaldehyde in 0.1 M phosphate buffer (PB; pH 7.4). The brain was quickly removed, postfixed in a solution of 4% paraformaldehyde at room temperature for 4 h, and then immersed overnight in a cold solution of 20% sucrose. Later, the brains were cut at 50 μ m thickness in the frontal plane using a freezing microtome.

The immunohistochemistry procedure was carried out as described previously (Liairy et al., 2012). After incubation in 1% H₂O₂ solution, floating sections were preincubated with 1.5% normal goat serum and 0.2% Triton-X-100 in 0.1 M PB (pH 7.4) for 1 h rotating at room temperature (RT), and then incubated overnight with the primary antibody, mouse anti-CD11b (1:500, AbSerotec, Oxford, UK). Subsequently, the sections were incubated for 1 h in biotinylated anti-mouse IgG (1:200, standard ABC Kit, Vector Labs, Burlingame, CA, USA), then incubated for 1 h in PBS containing avidin–biotin peroxidase complex (ABC), and then the sections were developed by incubating in PBS containing 10 mg diaminobenzidine (DAB) and 5 μ l of 30% hydrogen peroxide for 10 min.

For double immunofluorescent labeling, sections were incubated in the primary antibody, rabbit anti-ionized calcium binding adaptor molecule 1 (Iba1, 1:4000, Wako Ltd., Osaka, Japan), followed by Cy3-conjugated anti-rabbit IgG (1:1000, Amersham Bioscience Ltd., Piscataway, NJ, USA). For the secondary antibody, we used mouse anti-CD11b (1:500), followed by Alexa488-conjugated anti-mouse IgG (1:1000, Invitrogen, Carlsbad, CA, USA).

2.6. Measuring CD11b-labeled microglial cells immunoreactive area

The immunoreactive area was measured using a computer-assisted image analysis program (ImageTool V. 3.0, Department of Dental Diagnostic Science, University of Texas Health Science Center, San Antonio, TX, USA). Images were captured from four areas within the hippocampal DG: hilus, subgranular zone (SGZ), granular layer (GL), and molecular layer (ML). Sections containing CD11b-labeled cells were examined under a light microscope (Nikon, ECLIPSE E800). Images (N = 80 per region, per time point) were randomly captured from the same region of interest from interaural 5.88 mm, bregma 3.12 mm, to interaural 4.20 mm, bregma –4.80 mm, based on the rat brain atlas (Paxinos and Watson, 2007). The images were analyzed using the software that automates converting all immunolabeled elements that fall within a threshold range into pure black pixels and the rest of the image into pure white pixels. The software then quantifies the total numbers and percentages of black and white pixels for statistical analysis of the data.

2.7. Statistical analysis

The data are presented as the mean \pm standard error of the mean (SEM). The PPI data were analyzed by three-way analysis of variance (ANOVA) using R version 3.0.2 (The R Foundation for Statistical Computing, Vienna, Austria) which analyzed the interaction between three factors (strain \times treatment \times prepulse intensity). When appropriate, group means at individual levels were compared by post-hoc Bonferroni analysis to determine differences among groups. The NORT and immunohistochemistry data were analyzed by two-way ANOVA (strain \times treatment) and followed by the post hoc Bonferroni test. This analysis was performed with SPSS software (Dr. SPSS Inc for Windows V. 11.0, SPSS Inc., Chicago, IL, USA). Significance for the result was set at $p < 0.05$.

3. Results

3.1. Effects of minocycline on PPI deficits

The PPI test was conducted with two different prepulse intensities, 70 dB and 80 dB. The three-way ANOVA revealed that there was no significant three-way interaction between strain, treatment

and prepulse intensity ($F_{(1,48)} = 0.025, p > 0.05$). As for two-way interaction, no interaction between strain and prepulse intensity was significant ($F_{(1,48)} = 4.792, p < 0.05$). We found significant main effects for all factors included. The %PPI results were significantly affected by strain, treatment and prepulse intensity. %PPI was significantly lower for Gunn rats than Wistar rats ($F_{(1,48)} = 34.433, p < 0.001$). Treatment with minocycline produced significantly higher %PPI result than treatment with saline ($F_{(1,48)} = 4.413, p < 0.05$). As shown in Fig. 1, post hoc Bonferroni test revealed that the GV group had a significant lower %PPI compared to the WV group in both 70 dB ($F_{(1,24)} = 20.886, p < 0.001$) and 80 dB ($F_{(1,24)} = 9.040, p = 0.006$) tests. Treatment with minocycline for 14 days significantly increased the %PPI in the GM group compared to the GV group at 80 dB ($F_{(1,24)} = 3.60, p = 0.048$) but produced no significant difference at 70 dB ($F_{(1,24)} = 3.545, p = 0.072$). Meanwhile, no significant difference was found in WM compared to WV groups both at 70 dB ($F_{(1,24)} = 0.006, p = 0.941$) and at 80 dB ($F_{(1,24)} = 0.044, p = 0.835$).

3.2. Effects of minocycline on recognition memory

As shown in Fig. 2, in the training session, two-way ANOVA revealed that there were no significant interaction between strain \times treatment in both total time ($F_{(3,38)} = 2.272, p = 0.913$) and in exploratory preference ($F_{(3,38)} = 2.945, p = 0.412$). In the retention session, two-way ANOVA revealed significant differences among the groups studied in total time ($F_{(3,38)} = 0.257, p = 0.005$) and in exploratory preference ($F_{(3,38)} = 0.224, p = 0.004$). Post hoc analysis revealed that GV group showed significant reduction in total time ($F_{(1,38)} = 4.701, p = 0.036$) and exploratory preference ($F_{(1,38)} = 10.975, p = 0.002$) compared to the WV group. Treatment with minocycline in the GM group significantly improved the exploratory preference ($F_{(1,38)} = 8.126, p = 0.007$) without affecting the total exploration time ($F_{(1,38)} = 0.121, p = 0.730$) compared to the GV group. Meanwhile, treatment with minocycline showed no significant difference in the GM group compared to the WV group on total exploration time and exploratory preference in both the training or retention sessions.

3.3. Effects of minocycline on microglial activation

Next, we examined the effects of minocycline treatment on microglial activation in the hippocampal DG. CD11b expression in the hippocampal DG of the GV group was consistently high. In many cases, the CD11b immunoreactivity was detected not only in the thick processes, but also strongly expressed within the cell bodies of the Iba1-labeled microglial cells (Fig. 3-1b, -1c). Statistical analysis by two-way ANOVA revealed a significant interaction between strain \times treatment in hilus ($F_{(3,20)} = 1.977, p = 0.006$), SGZ ($F_{(3,20)} = 1.902, p = 0.003$), granular layer ($F_{(3,20)} = 7.135, p = 0.045$), but not in molecular layer ($F_{(3,20)} = 6.986, p = 0.644$). Post hoc analysis indicated a significant increase in percentage of CD11b immunoreactivity of the GV group in the hilus ($F_{(1,20)} = 43.506, p < 0.001$), SGZ ($F_{(1,20)} = 57.156, p < 0.001$), granular layer ($F_{(1,20)} = 18.619, p < 0.001$), and molecular layer ($F_{(1,20)} = 11.323, p = 0.003$) compared to the WV group (Fig. 4a–d).

The consecutive administration of minocycline (40 mg/kg) for 14 days decreased the CD11b expression in the DG of the GM group. In this case, most CD11b immunoreactivity was expressed only on the thin ramified processes with a slight expression within the cell bodies of the Iba1-labeled microglial cells (Fig. 3-2b, 3-2c). A post hoc Bonferroni test revealed that minocycline significantly decreased the percentage of CD11b immunoreactivity in the hilus ($F_{(1,20)} = 23.510, p < 0.001$), SGZ ($F_{(1,20)} = 18.473, p < 0.001$), and granular layer ($F_{(1,20)} = 8.907, p = 0.007$) compared to the GV group (Fig. 4a–c). No significant difference was found between WV and WM groups.




Fig. 1. Effects of minocycline on PPI test at 70 dB and 80 dB. Each value is the mean \pm S.E.M. ($n = 7$ per group). * $p < 0.01$, ** $p < 0.001$ as compared to WV control group; # $p < 0.05$ as compared to GV group.




Fig. 2. Effects of minocycline on novel object recognition test (NORT). A. Total exploration time; B. exploratory preference. Each value is the mean \pm S.E.M. ($n = 9$ each in Wistar rat group; $n = 12$ each in Gunn rat group). * $p < 0.05$, ** $p < 0.005$ as compared to WV control group; # $p < 0.01$ as compared to GV group.

4. Discussion

The major findings of the present study are that minocycline significantly improved the impairment of recognition memory in Gunn rats, and that minocycline significantly attenuated the activation of microglial cells in the hippocampal DG. To our knowledge, this is the first report demonstrating the effects of minocycline on behavioral changes and on microglial activation in the Gunn rat, a possible hyperbilirubinemia-induced animal model of schizophrenia.

Our results showed that Gunn rats had lower %PPI compared to Wistar control rats, indicating that the Gunn rat has impaired sensorimotor gating. Moreover, Gunn rats also showed a reduction of exploratory preference after long-term retention interval (24 h) during NORT, indicating that the Gunn rat has impaired recognition memory. Our findings are in agreement with other animal models of schizophrenia on the association with PPI deficits (Dieckmann et al., 2007; Geyer et al., 1990) and NORT impairment (Kamei et al., 2006; Mizoguchi et al., 2008). Taking these factors together, we herein propose the Gunn rat as a potential animal model in relation to the cognitive deficits in schizophrenia.

To detect and measure microglial activation in the hippocampal DG, we used a marker of integrin α M (ITGAM) cluster of differentiation molecule 11b (CD11b), which is expressed on all types of microglia, and expression of which is significantly increased on activated microglial cells (He et al., 1997). Furthermore, one of the earliest changes in microglia cells after facial nerve transection was the up-regulation of CD11b (Graeber et al., 1988). Therefore, CD11b expression appears to be a sensitive marker of microglial activation. The hippocampal region was our main interest due to its potential role in learning and memory with the dentate gyrus (DG) as the primary gateway for inputs into the hippocampus with a continuously developing structure with birth of new neurons – ‘neurogenesis’ – throughout mammalian life (Gage, 2002).




Fig. 3. CD11b expression in ionized calcium binding adaptor molecule 1 (Iba1)-labeled microglial cells. A confocal Z stack image depicting Iba1-labeled (red), CD11b expression (green), and merged (yellow) cells of vehicle-treated Gunn rat (GV) group (1a–1c) compared to those of minocycline-treated Gunn rat (GM) group (2b,c). Microglial cells in the GM (2b,c) group expressed low levels of CD11b immunoreactivity compared to the GV group (1b,c). Scale bars: 200 μm (1,2), 20 μm (1a–1c, 2a–2c).

Moreover, hippocampus is also one of the brain regions whose function is altered in schizophrenia (Tamminga et al., 2010).

Consistent with our previous report, we found a significant high level of CD11b immunoreactivity in all areas of the hippocampal DG, including the hilus, SGZ, granular layer, and molecular layer in Gunn rats. The high level of CD11b immunoreactivity was accompanied by altered morphology, suggesting that microglial cells in Gunn rats were in an activated state. Microglial cell is well-known as a key player in the function of the cerebral innate immune system to pathological changes. Threats

to CNS homeostasis can trigger a rapid transformation from a normally “ramified” state into alerted and “activated” states. Microglial cells primarily serve in tissue defense and protection when participating in the mechanisms of innate and adaptive immunity. Conversely, excessive acute or chronic microglial activation can provoke severe neuronal and glial damage by carrying or fuelling destructive cascades [for review, see (Kettenmann et al., 2011; van Rossum and Hanisch, 2004)]. Microglial cells may also contribute to synaptic modulation, learning, and memory processes. Disturbance of the functions or alterations in




Fig. 4. Mean percentage of black pixels indicating CD11b immunoreactivity in the hippocampal dentate gyrus (DG). Each value is the mean ± S.E.M. (n = 6 per group). *p < 0.005, **p < 0.001 as compared to WV control group; #p < 0.01, ##p < 0.001 as compared to GV group.

the number and/or morphology of microglial cells in brain regions involved in cognitive and emotional aspects of behavior may represent an etiological factor in the onset or progression of memory impairment and neuropsychiatric disorders (Blank and Prinz, 2013). Accordingly, we assumed that the chronic microglial activation in adult Gunn rat is potentially more maladaptive than neuroprotective (Liaury et al., 2012), and based on our present results, we suggest that the microglial activation in the hippocampal DG in Gunn rat may be involved in promoting cognitive impairment, as was shown in the behavior test.

Consecutive administration of minocycline for 14 days significantly improved the impaired recognition memory in Gunn rats, which was indicated by the increase of exploratory preference in the retention session during SORT. Moreover, minocycline significantly attenuated the activation of microglial cells in the hippocampal DG of Gunn rats, which was indicated by the decreased immunoreactivity of CD11b marker. According to previous studies showing about the neuroprotective effects of minocycline (Miyaka, 2012; Yrjanheikki et al., 1998), we assume that this phenomenon is part of the mechanism underlying minocycline's anti-inflammatory properties through inhibition of microglial activation. Similar results were reported that minocycline significantly improved the recognition memory of mice treated with methamphetamine (Mizoguchi et al., 2008) and phencyclidine (Fujita et al., 2008). Minocycline was also reported to significantly attenuate the microglial activation in mouse brains induced by methamphetamine and 3,4-methylenedioxymethamphetamine (Zhang et al., 2006a, 2006b). Taking these facts and our results into consideration, we speculate that the inhibition of microglial activation by minocycline in the hippocampal DG may, in part, be implicated in the mechanism of minocycline action to improve the recognition memory in Gunn rats. Although we demonstrated that minocycline attenuated microglial activation, we did not evaluate its effects on inflammatory markers, such as pro-inflammatory cytokines and chemokines. Therefore, it may be of interest to further study and establish the anti-inflammatory effects of minocycline in Gunn rats. On the other hand, the PPI deficit in Gunn rats was not fully rescued by minocycline administration. While the results at the 80 dB showed a significant improvement in the PPI deficits, later we only found a tendency toward improvement at the 70 dB. We speculate that this may be due to the small number of animals used for the PPI test ($n = 7$, per group). Other possibility is that it may be due to the incomplete inhibition of microglial activation, since minocycline did not reduce the immunoreactivity of CD11b in the molecular layer of the hippocampal DG in Gunn rats. Another study also reported the partial improvement of minocycline on olfactory bulbectomized (OBX) rats, a model of cognitive and behavioral impairments arising from neurodegenerative processes (Borre et al., 2012). Minocycline normalized OBX-induced hyperactivity in the open field, protected against hippocampal dependent spatial memory deficit, but failed to prevent fear memory loss. They suggested that minocycline may have ameliorated the OBX-induced cognitive deficits in a region specific manner and that treatment with minocycline may be effective in early phase of a neurodegenerative disease.

However, the exact cellular and molecular mechanisms underlying this complex UCB–neuron–glia interaction in Gunn rats still remain elusive. We presume that another mechanism may also be involved, which is mediated through the glutamatergic system in the brain. Exposure of glia cells to UCB increases the extracellular concentration of glutamate by decreased uptake (Silva et al., 1999) and/or enhanced secretion (Fernandes et al., 2004), engendering overstimulation of glutamate and N-methyl-D-aspartate (NMDA) receptor (Ostrow et al., 2004). Moreover, high levels of bilirubin can lower the brain's threshold and enhance its vulnerability to NMDA-triggered excitotoxic brain injury (McDonald et al., 1998). Release of pro-inflammatory cytokines may affect gliogenesis and neurogenesis, and may lead to deficits in learning and memory. Dysregulation of glutamate metabolism and over-expression of tumor necrosis factor- α (TNF- α) and interleukin (IL)-1 β are consistent with schizophrenia neuropathology. Increased

proinflammatory status of the brain also interacts with glutamatergic and dopaminergic neurotransmission, which can induce or aggravate positive, negative, and cognitive symptoms of schizophrenia (Muller and Schwarz, 2006). Interestingly, some studies have shown that minocycline has properties that affect glutamatergic pathways. Administration of minocycline to neurons in-vitro and to mice in-vivo increased GluR1 phosphorylation and membrane insertion (Imbesi et al., 2008). GluR1 receptors, which appear to be critical for cognitive processes that are impaired in schizophrenia, may be crucially involved in the pathobiology of schizophrenia (Wiedholz et al., 2008). Moreover, minocycline attenuated behavioral changes after administration of a non-competitive N-methyl-D-aspartate (NMDA) receptor antagonist (Levkovitz et al., 2007; Zhang et al., 2007).

Inhibition of chronic neuroinflammation, particularly of microglial activation, has been suggested to be a practical strategy in the treatment of major psychiatry diseases, including schizophrenia (Dean et al., 2012). Two studies reported the effectiveness of minocycline adjunctive therapy in early schizophrenia for negative symptoms (Chaudhry et al., 2012; Levkovitz et al., 2010). In addition, Levkovitz et al. also reported the improvement on cognitive function. Another report showed two cases with persistent schizophrenia symptoms despite long term clozapine treatment that were treated successfully with adjunct minocycline (Liaury et al., 2011). Moreover, long term treatment with minocycline is generally safe and well tolerated in humans (Bonelli et al., 2003; Stone et al., 2003). Taken together, it might be worthwhile to further explore the effects of minocycline in schizophrenia.

5. Conclusion

The present study indicates that minocycline improves recognition memory and attenuates the activation of microglial cells in the hippocampal DG of Gunn rat, a possible hyperbilirubine induced animal model of schizophrenia. Therefore, minocycline may be a potential therapeutic drug for schizophrenia. Our results may also provide crucial information to elucidate the etiology of schizophrenia and support the possibility of using the Gunn rat as an animal model of schizophrenia.

Acknowledgments

This work was supported in part by a Grant-in-Aid for Scientific Research (C) (No. 13201342) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and a Health and Labor Sciences Research Grant for Clinical Research (No. 04T-580) from the Ministry of Health, Labor, and Welfare of Japan. Part of KL's work was also supported by a grant from the Hashiya Scholarship Foundation. All authors have read and approved the final manuscript. All authors declare no competing interest. We thank Dr. Hideaki Yasuda (Department of Psychiatry, Shimane University Faculty of Medicine) and Akira Yasuda (Department of Medical Biostatistic, Shimane University Faculty of Medicine) for their valuable suggestions on statistical analysis. We are particularly grateful to Katherine Ono for editorial assistance.

References

- Bian Q, Kato T, Monji A, Hashioka S, Mizoguchi Y, Horikawa H, et al. The effect of atypical antipsychotics, perospirone, ziprasidone and quetiapine on microglial activation induced by interferon-gamma. *Prog Neuropsychopharmacol Biol Psychiatry* 2008;32(1):42–8.
- Blank T, Prinz M. Microglia as modulators of cognition and neuropsychiatric disorders. *Glia* 2013;65(1):62–70.
- Bonelli RM, Heuberger C, Reisecker F. Minocycline for Huntington's disease: an open label study. *Neurology* 2003;60(5):883–4.
- Borre Y, Sir V, de Kivit S, Westphal KG, Olivier B, Oosting RS. Minocycline restores spatial but not fear memory in olfactory bulbectomized rats. *Eur J Pharmacol* 2012;697(1–3):59–64.
- Chaudhry IB, Hallak J, Husain N, Minhas F, Stirling J, Richardson P, et al. Minocycline benefits negative symptoms in early schizophrenia: a randomised double-blind placebo-controlled clinical trial in patients on standard treatment. *J Psychopharmacol* 2012;26(9):1185–93.

- Dean OM, Data-Franco J, Giorlando F, Berk M. Minocycline: therapeutic potential in psychiatry. *CNS Drugs* 2012;26(5):391–401.
- Diekmann M, Freudenberg F, Klein S, Koch M, Schwabe K. Disturbed social behavior and motivation in rats selectively bred for deficient sensorimotor gating. *Schizophr Res* 2007;97(1–3):250–3.
- Doorduyn J, de Vries EF, Willemsen AT, de Groot JC, Dierckx RA, Klein HC. Neuroinflammation in schizophrenia-related psychosis: a PET study. *J Nucl Med* 2009;50(11):1801–7.
- Erbagci AB, Herken H, Koyluoglu O, Yilmaz N, Tarakcioglu M. Serum IL-1beta, sIL-2R, IL-6, IL-8 and TNF-alpha in schizophrenic patients, relation with symptomatology and responsiveness to risperidone treatment. *Mediators Inflamm* 2001;10(3):109–15.
- Fernandes A, Silva RF, Fakao AS, Brito MA, Brites D. Cytokine production, glutamate release and cell death in rat cultured astrocytes treated with unconjugated bilirubin and IPS. *J Neuroimmunol* 2004;153(1–2):64–75.
- Ferretti MT, Allard S, Partridge V, Ducatenzeiler A, Cuello AC. Minocycline corrects early, pre-plaque neuroinflammation and inhibits BACE-1 in a transgenic model of Alzheimer's disease-like amyloid pathology. *J Neuroinflammation* 2012;9(62):1742–2094.
- Fisman M. The brain stem in psychosis. *Br J Psychiatry* 1975;126:414–22.
- Fujita Y, Ishima T, Kunitachi S, Hagiwara H, Zhang L, Iyo M, et al. Phencyclidine-induced cognitive deficits in mice are improved by subsequent subchronic administration of the antibiotic drug minocycline. *Prog Neuropsychopharmacol Biol Psychiatry* 2008;32(2):336–9.
- Gage FH. Neurogenesis in the adult brain. *J Neurosci* 2002;22(3):612–3. [2002 Feb 1].
- Geyer MA, Swerdlow NR, Mansbach RS, Braff DL. Startle response models of sensorimotor gating and habituation deficits in schizophrenia. *Brain Res Bull* 1990;25(3):485–98.
- Graeber MB, Streit WJ, Kreutzberg GW. Axotomy of the rat facial nerve leads to increased CR3 complement receptor expression by activated microglial cells. *J Neurosci Res* 1988;21(1):18–24.
- Gunn CK. Hereditary acholuric jaundice in the rat. *Can Med Assoc J* 1944;50(3):230–7.
- Hayashida M, Miyaoka T, Tsuchie K, Yasuda H, Wake R, Nishida A, et al. Hyperbilirubinemia-related behavioral and neuropathological changes in rats: a possible schizophrenia animal model. *Prog Neuropsychopharmacol Biol Psychiatry* 2009;33(4):581–8.
- He BP, Tay SS, Leong SK. Microglia responses in the CNS following sciatic nerve transection in C57BL/6J and BALB/c mice. *Exp Neurol* 1997;146(2):587–95.
- Imbesi M, Uz T, Manev R, Sharma RP, Manev H. Minocycline increases phosphorylation and membrane insertion of neuronal GluR1 receptors. *Neurosci Lett* 2008;447(2–3):134–7.
- Juckel G, Manitz MP, Brune M, Friebe A, Heneka MT, Wolf RJ. Microglial activation in a neuroinflammatory animal model of schizophrenia – a pilot study. *Schizophr Res* 2011;131(1–3):96–100.
- Kamei H, Nagai T, Nakano H, Togan Y, Takayangi M, Takahashi K, et al. Repeated methamphetamine treatment impairs recognition memory through a failure of novelty induced ERK1/2 activation in the prefrontal cortex of mice. *Biol Psychiatry* 2006;59(1):75–84.
- Kato T, Monji A, Hashioka S, Kanba S. Risperidone significantly inhibits interferon-gamma-induced microglial activation in vitro. *Schizophr Res* 2007;92(1–3):108–15.
- Kelly DL, Vyas G, Richardson CM, Koola M, McMahon RP, Buchanan RW, et al. Adjunct minocycline to clozapine treated patients with persistent schizophrenia symptoms. *Schizophr Res* 2011;133(1–3):257–8. <http://dx.doi.org/10.1016/j.schres.2011.08.005>. [2011 Dec, Epub 2011 Aug 26].
- Kettenmann H, Hanisch UK, Noda M, Verkhratsky A. Physiology of microglia. *Physiol Rev* 2011;91(2):461–553.
- Kneeland RE, Fatemi SH. Viral infection, inflammation and schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2013;42:35–48.
- Laan W, Grobbee DE, Selten JP, Heijnen CJ, Kahn RS, Burger H. Adjuvant aspirin therapy reduces symptoms of schizophrenia spectrum disorders: results from a randomized, double-blind, placebo-controlled trial. *J Clin Psychiatry* 2010;71(5):520–7.
- Levkovitz Y, Levi U, Braw Y, Cohen H. Minocycline, a second-generation tetracycline, as a neuroprotective agent in an animal model of schizophrenia. *Brain Res* 2007;18:154–62.
- Levkovitz Y, Mendlovich S, Riwkes S, Braw Y, Levkovitch-Verbin H, Gal G, et al. A double-blind, randomized study of minocycline for the treatment of negative and cognitive symptoms in early-phase schizophrenia. *J Clin Psychiatry* 2010;71(2):138–49.
- Liury K, Miyaoka T, Tsumori T, Furuya M, Wake R, Ieda M, et al. Morphological features of microglial cells in the hippocampal dentate gyrus of Gunn rat: a possible schizophrenia animal model. *J Neuroinflammation* 2012;9(56):1742–2094.
- McDonald JW, Shapiro SM, Silverstein FS, Johnston MV. Role of glutamate receptor-mediated excitotoxicity in bilirubin-induced brain injury in the Gunn rat model. *Exp Neurol* 1998;150(1):21–9.
- Mednick SA, Machon RA, Huttunen MO, Bonett D. Adult schizophrenia following prenatal exposure to an influenza epidemic. *Arch Gen Psychiatry* 1988;45(2):189–92.
- Meyer U. Developmental neuroinflammation and schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2013;42:20–34.
- Miller BJ, Buckley P, Seabolt W, Mellor A, Kirkpatrick B. Meta-analysis of cytokine alterations in schizophrenia: clinical status and antipsychotic effects. *Biol Psychiatry* 2011;70(7):663–71.
- Miller BJ, Culeppner N, Rapaport MH, Buckley P. Prenatal inflammation and neurodevelopment in schizophrenia: a review of human studies. *Prog Neuropsychopharmacol Biol Psychiatry* 2013;42:92–100.
- Miyaoka T. Minocycline for schizophrenia: a critical review. *Open J Psychiatry* 2012;2:399–406.
- Miyaoka T, Yasukawa R, Yasuda H, Hayashida M, Inagaki T, Horiguchi J. Possible antipsychotic effects of minocycline in patients with schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2007;31(11):304–7.
- Miyaoka T, Yasukawa R, Yasuda H, Hayashida M, Inagaki T, Horiguchi J. Minocycline as adjunctive therapy for schizophrenia: an open-label study. *Clin Neuropharmacol* 2008;31(5):287–92.
- Mizoguchi H, Takuma K, Fukakusa A, Ito Y, Nakatani A, Ibi D, et al. Improvement by minocycline of methamphetamine-induced impairment of recognition memory in mice. *Psychopharmacology (Berl)* 2008;196(2):233–41.
- Muller N, Schwarz M. Schizophrenia as an inflammation-mediated dysbalance of glutamatergic neurotransmission. *Neurotox Res* 2006;10(2):131–48.
- Muller N, Schwarz MJ. COX-2 inhibition in schizophrenia and major depression. *Curr Pharm Des* 2008;14(14):1452–65.
- Ostrow JD, Pascolo L, Brites D, Tiribelli C. Molecular basis of bilirubin-induced neurotoxicity. *Trends Mol Med* 2004;10(2):65–70.
- Paxinos G, Watson C. The rat brain in stereotaxic coordinates. London: Elsevier Inc.; 2007 [Vol 6th Edition].
- Radevich K, Garey LJ, Gentleman SM, Reynolds R. Increase in HLA-DR immunoreactive microglia in frontal and temporal cortex of chronic schizophrenics. *J Neuropathol Exp Neurol* 2000;59(2):137–50.
- Ripke S, Sanders AR, Kendler KS, Levinson DF, Sklar P, Holmans PA, et al. Genome-wide association study identifies five new schizophrenia loci. *Nat Genet* 2011;43(10):969–76.
- Schwab SG, Hallmayer J, Freimann J, Lerer B, Albus M, Bormann-Hassenbach M, et al. Investigation of linkage and association/linkage disequilibrium of HLA A-, DQA1-, DQB1-, and DRB1- alleles in 69 sib-pair- and 89 trio-families with schizophrenia. *Am J Med Genet* 2002;114(3):315–20.
- Silva R, Mata LR, Gulbenkian S, Brito MA, Tinelli C, Brites D. Inhibition of glutamate uptake by unconjugated bilirubin in cultured cortical rat astrocytes: role of concentration and pH. *Biochem Biophys Res Commun* 1999;265(1):67–72.
- Steiner J, Bielau H, Brisch R, Danos P, Ullrich O, Mawrin C, et al. Immunological aspects in the neurobiology of suicide: elevated microglial density in schizophrenia and depression is associated with suicide. *J Psychiatr Res* 2008;42(2):151–7.
- Stone M, Fortin PR, Pacheco-Tena C, Inman RD. Should tetracycline treatment be used more extensively for rheumatoid arthritis? Meta-analysis demonstrates clinical benefit with reduction in disease activity. *J Rheumatol* 2003;30(10):2112–22.
- Takahashi N, Sakurai T. Roles of glial cells in schizophrenia: possible targets for therapeutic approaches. *Neurobiol Dis* 2013;53:49–60.
- Tamminga CA, Stan AD, Wagner AD. The hippocampal formation in schizophrenia. *Am J Psychiatry* 2010;167(10):1178–93.
- Tsuchie K, Miyaoka T, Furuya M, Liury K, Ieda M, Wake R, et al. The effects of antipsychotics on behavioral abnormalities of the Gunn rat (unconjugated hyperbilirubinemia rat), a rat model of schizophrenia. *Asian Journal of Psychiatry* 2013;6(2):119–23.
- van Berckel BN, Bossong MG, Boellaard R, Kloet R, Schuitmaker A, Caspers E, et al. Microglia activation in recent-onset schizophrenia: a quantitative [¹¹C]PK11195 positron emission tomography study. *Biol Psychiatry* 2008;64(9):820–2.
- van Os J, Kapur S. Schizophrenia. *Lancet* 2009;374(9690):635–45.
- van Rossum D, Hanisch UK. Microglia. *Metab Brain Dis* 2004;19(3–4):393–411.
- Wiedholz LM, Owens WA, Horton RE, Feyder M, Karlsson RM, Hefner K, et al. Mice lacking the AMPA GluR1 receptor exhibit striatal hyperdopaminergia and 'schizophrenia-related' behaviors. *Mol Psychiatry* 2008;13(6):631–40.
- Yrjanheikki J, Keinanen R, Pellikka M, Hokfelt T, Koistinaho J. Tetracyclines inhibit microglial activation and are neuroprotective in global brain ischemia. *Proc Natl Acad Sci U S A* 1998;95(26):15769–74.
- Zhang L, Kitaichi K, Fujimoto Y, Nakayama H, Shimizu E, Iyo M, et al. Protective effects of minocycline on behavioral changes and neurotoxicity in mice after administration of methamphetamine. *Prog Neuropsychopharmacol Biol Psychiatry* 2006a;30(8):1381–93.
- Zhang L, Shirayama Y, Shimizu E, Iyo M, Hashimoto K. Protective effects of minocycline on 3,4-methylenedioxymethamphetamine-induced neurotoxicity in serotonergic and dopaminergic neurons of mouse brain. *Eur J Pharmacol* 2006b;544(1–3):1–9.
- Zhang L, Shirayama Y, Iyo M, Hashimoto K. Minocycline attenuates hyperlocomotion and prepulse inhibition deficits in mice after administration of the NMDA receptor antagonist dizocilpine. *Neuropsychopharmacology* 2007;32(9):2004–10.

Minocycline improves recognition memory and attenuates microglial activation in Gunn rat: A possible hyperbilirubinemia-induced animal model of schizophrenia

ORIGINALITY REPORT

%**24**

SIMILARITY INDEX

%**3**

INTERNET SOURCES

%**22**

PUBLICATIONS

%**28**

STUDENT PAPERS

PRIMARY SOURCES

1

link.springer.com

Internet Source

%**3**

2

Jian-Dong Long, Yao Liu, Dong-Liang Jiao, Yu-Jun Wang, Gui-Ying Zan, Yun-Yue Ju, Min Zhao, Jing-Gen Liu. "The neuroprotective effect of memantine on methamphetamine-induced cognitive deficits", Behavioural Brain Research, 2017

Publication

%**2**

3

Zhang, L.. "Protective effects of minocycline on behavioral changes and neurotoxicity in mice after administration of methamphetamine", Progress in Neuropsychopharmacology & Biological Psychiatry, 20061230

Publication

%**1**

4

Daisuke Ibi, Taku Nagai, Hiroyuki Koike, Yuko Kitahara et al. "Combined effect of neonatal immune activation and mutant DISC1 on phenotypic changes in adulthood", Behavioural

%**1**

Brain Research, 2010

Publication

-
- | | | |
|---|---|-----|
| 5 | www.ncbi.nlm.nih.gov
Internet Source | % 1 |
|---|---|-----|
-
- | | | |
|---|---|-----|
| 6 | Zhu, Furong, Yingjun Zheng, Yong Liu, Xianghui Zhang, and Jingping Zhao. "Minocycline alleviates behavioral deficits and inhibits microglial activation in the offspring of pregnant mice after administration of polyriboinosinic–polyribocytidilic acid", <i>Psychiatry Research</i> , 2014.
Publication | % 1 |
|---|---|-----|
-
- | | | |
|---|---|-----|
| 7 | Yuliya Borre, Volkan Sir, Sander de Kivit, Koen G. Westphal, Berend Olivier, Ronald S. Oosting. "Minocycline restores spatial but not fear memory in olfactory bulbectomized rats", <i>European Journal of Pharmacology</i> , 2012
Publication | % 1 |
|---|---|-----|
-
- | | | |
|---|---|-----|
| 8 | www.tandfonline.com
Internet Source | % 1 |
|---|---|-----|
-
- | | | |
|---|---|-----|
| 9 | www.dovepress.com
Internet Source | % 1 |
|---|---|-----|
-
- | | | |
|----|--|-----|
| 10 | Submitted to West Virginia University
Student Paper | % 1 |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 11 | hdl.handle.net
Internet Source | % 1 |
|----|---|-----|
-

12

Radmila Manev, Hari Manev. "Minocycline, schizophrenia and GluR1 glutamate receptors", *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 2009

Publication

% 1

13

Submitted to CSU Northridge

Student Paper

% 1

14

www.jneuroinflammation.com

Internet Source

% 1

15

Hiroyuki Mizoguchi, Kazuhiro Takuma, Ayumi Fukakusa, Yukio Ito, Akiko Nakatani, Daisuke Ibi, Hyoung-Chun Kim, Kiyofumi Yamada.

"Improvement by minocycline of methamphetamine-induced impairment of recognition memory in mice", *Psychopharmacology*, 2007

Publication

% 1

16

D Brites, A Fernandes, A S Falcão, A C Gordo, R F M Silva, M A Brito. "Biological risks for neurological abnormalities associated with hyperbilirubinemia", *Journal of Perinatology*, 2009

Publication

% 1

17

onlinelibrary.wiley.com

Internet Source

% 1

18

Submitted to American Public University System

19

bmcneurosci.biomedcentral.com

Internet Source

% 1

20

Blank, Thomas, and Marco Prinz. "Microglia as modulators of cognition and neuropsychiatric disorders", *Glia*, 2013.

Publication

% 1

21

Maria Teresa Ferretti, Simon Allard, Vanessa Partridge, Adriana Ducatenzeiler, A Claudio Cuello. "Minocycline corrects early, pre-plaque neuroinflammation and inhibits BACE-1 in a transgenic model of Alzheimer's disease-like amyloid pathology", *Journal of Neuroinflammation*, 2012

Publication

% 1

22

journals.sagepub.com

Internet Source

% 1

23

www.frontiersin.org

Internet Source

% 1

24

Kazuhiro Ishii, Taku Nagai, Yuki Hirota, Mariko Noda, Toshitaka Nabeshima, Kiyofumi Yamada, Ken-ichiro Kubo, Kazunori Nakajima. "Reelin has a preventive effect on phencyclidine-induced cognitive and sensory-motor gating deficits", *Neuroscience Research*, 2015

% 1

25

"Poster presentations", Schizophrenia
Research, 200601

Publication

% 1

26

repositorio.unifesp.br

Internet Source

% 1

27

academic.oup.com

Internet Source

% 1

28

www.nature.com

Internet Source

% 1

29

journals.plos.org

Internet Source

% 1

30

www.ijcem.com

Internet Source

% 1

31

Yuji Uchio, Naotaka Shu, Mitsuo Ochi.
"ADVERSE EFFECTS OF SALINE
NEUROLYSIS ON A CHRONICALLY
COMPRESSED RAT SCIATIC NERVE",
Scandinavian Journal of Plastic and
Reconstructive Surgery and Hand Surgery,
8/23/2000

Publication

<% 1

32

Manoj Kumar Mishra. "Minocycline
neuroprotects, reduces microglial activation,
inhibits caspase 3 induction, and viral replication

<% 1

following Japanese encephalitis", Journal of
Neurochemistry, 2/18/2008

Publication

33

www.wfsbp.org

Internet Source

<% 1

34

jpet.aspetjournals.org

Internet Source

<% 1

35

Submitted to Trinity College Dublin

Student Paper

<% 1

36

opac.ll.chiba-u.jp

Internet Source

<% 1

37

ddd.uab.cat

Internet Source

<% 1

38

repository.unhas.ac.id

Internet Source

<% 1

39

A.S. Falcão, C. Bellarosa, A. Fernandes, M.A. Brito, R.F.M. Silva, C. Tiribelli, D. Brites. "Role of multidrug resistance-associated protein 1 expression in the in vitro susceptibility of rat nerve cell to unconjugated bilirubin", Neuroscience, 2007

Publication

<% 1

40

Mutiara, S.. "The involvement of phosphatidylinositol 3-kinase in gonadotropin-releasing hormone-induced gonadotropin @a-

<% 1

and FSH@b-subunit genes expression in clonal gonadotroph L@bT2 cells", Molecular and Cellular Endocrinology, 20080213

Publication

41

estudogeral.sib.uc.pt

Internet Source

<% 1

42

digitalcommons.wustl.edu

Internet Source

<% 1

43

Wake, R.. "Characteristic brain hypoperfusion by 99mTc-ECD single photon emission computed tomography (SPECT) in patients with the first-episode schizophrenia", European Psychiatry, 201010

Publication

<% 1

44

M. B. Graeber, W. J. Streit, G. W. Kreutzberg. "Axotomy of the rat facial nerve leads to increased CR3 complement receptor expression by activated microglial cells", Journal of Neuroscience Research, 1988

Publication

<% 1

45

qsciences.com

Internet Source

<% 1

46

Submitted to University of Pittsburgh

Student Paper

<% 1

47

Suk-Chan Hahm, Young Wook Yoon, Junesun Kim. "High-Frequency Transcutaneous Electrical

<% 1

Nerve Stimulation Alleviates Spasticity After Spinal Contusion by Inhibiting Activated Microglia in Rats", Neurorehabilitation and Neural Repair, 2014

Publication

48

Keisuke Shima, Koji Shimatani, Yuichi Kurita, Toshio Tsuji. "Novel method for mitigation of body sway and preliminary results for tandem standing", MHS2013, 2013

Publication

<% 1

49

"Abstracts for the 13th International Congress on Schizophrenia Research (ICOSR)", Schizophrenia Bulletin, 03/01/2011

Publication

<% 1

50

Yulug, B., and W.-R. Schabitz. "Towards A Multimodal Treatment of Depression: A Minireview on the Potential Role of Antibiotics", Mini Reviews in Medicinal Chemistry, 2011.

Publication

<% 1

51

Takahashi, Nagahide, and Takeshi Sakurai. "Roles of glial cells in schizophrenia: Possible targets for therapeutic approaches", Neurobiology of Disease, 2013.

Publication

<% 1

52

Sung-Sen Yang, Chuen-Lin Huang, Huei-E Chen, Che-Se Tung, Han-Peng Shih, Yia-Ping Liu. "Effects of SPAK knockout on sensorimotor

<% 1

gating, novelty exploration, and brain area-dependent expressions of NKCC1 and KCC2 in a mouse model of schizophrenia", Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2015

Publication

53

McDonald, J.W.. "Role of Glutamate Receptor-Mediated Excitotoxicity in Bilirubin-Induced Brain Injury in the Gunn Rat Model", Experimental Neurology, 199803

Publication

<% 1

54

T. Breidert. "Protective action of the peroxisome proliferator-activated receptor-gamma agonist pioglitazone in a mouse model of Parkinson's disease", Journal of Neurochemistry, 8/2002

Publication

<% 1

55

mediterranean.seagrassonline.org

Internet Source

<% 1

56

Submitted to University of Bath

Student Paper

<% 1

57

Xiaohong Chen, Xiaomeng Ma, Ying Jiang, Rongbiao Pi, Yingying Liu, Lili Ma. "The prospects of minocycline in multiple sclerosis", Journal of Neuroimmunology, 2011

Publication

<% 1

58

L.G. Staples, I.S. McGregor, R. Apfelbach, G.E. Hunt. "Cat odor, but not trimethylthiazoline (fox

<% 1

odor), activates accessory olfactory and defense-related brain regions in rats", *Neuroscience*, 2008

Publication

59

d-nb.info

Internet Source

<% 1

60

Kenji Hashimoto. "Microglial activation in schizophrenia and minocycline treatment", *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 2008

Publication

<% 1

61

Yi He, Stanley Appel, Weidong Le. "Minocycline inhibits microglial activation and protects nigral cells after 6-hydroxydopamine injection into mouse striatum", *Brain Research*, 2001

Publication

<% 1

62

bio.jsnu.edu.cn

Internet Source

<% 1

63

Kelly, Deanna L., Kelli M. Sullivan, Joseph P. McEvoy, Robert P. McMahon, Heidi J. Wehring, James M. Gold, Fang Liu, Dale Warfel, Gopal Vyas, Charles M. Richardson, Bernard A. Fischer, William R. Keller, Maju Mathew Koola, Stephanie M. Feldman, Jessica C. Russ, Richard SE Keefe, Jennifer Osing, Leeka Hubzin, Sharon August, Trina M. Walker, and Robert W. Buchanan. "Adjunctive Minocycline in

<% 1

Clozapine-Treated Schizophrenia Patients With Persistent Symptoms :", Journal of Clinical Psychopharmacology, 2015.

Publication

64

Arango, Celso, George Garibaldi, and Stephen R. Marder. "Pharmacological approaches to treating negative symptoms: A review of clinical trials", Schizophrenia Research, 2013.

Publication

<% 1

65

escholarship.org

Internet Source

<% 1

66

S.-I. Yeo, J.-E. Kim, H.J. Ryu, C.H. Seo, B.C. Lee, I.-G. Choi, D.-S. Kim, T.-C. Kang. "The roles of fractalkine/CX3CR1 system in neuronal death following pilocarpine-induced status epilepticus", Journal of Neuroimmunology, 2011

Publication

<% 1

67

Volker Arolt, Oliver Ambrée. "The Question of Pro-Inflammatory Immune Activity in Schizophrenia and the Potential Importance of Anti-Inflammatory Drugs", S. Karger AG, 2013

Publication

<% 1

68

Yamamoto, Yasushi, Masahisa Onoguchi, Kazunori Kawakami, Masuo Haramoto, Rei Wake, Jun Horiguchi, and Hajime Kitagaki. "Evaluation of the difference-correction effect of the gamma camera systems used by easy Z-

<% 1

score Imaging System (eZIS) analysis", *Annals of Nuclear Medicine*, 2014.

Publication

69

Kelly, D.L.. "Adjunct minocycline to clozapine treated patients with persistent schizophrenia symptoms", *Schizophrenia Research*, 201112

Publication

<% 1

70

Yuliya E. Borre, Theodora Panagaki, Pim J. Koelink, Mary. E. Morgan et al. "Neuroprotective and cognitive enhancing effects of a multi-targeted food intervention in an animal model of neurodegeneration and depression", *Neuropharmacology*, 2014

Publication

<% 1

71

www.hopkinsmedicine.org

Internet Source

<% 1

72

apsjournals.apsnet.org

Internet Source

<% 1

73

David A Taylor. "Cannabidiol Reverses MK-801-Induced Disruption of Prepulse Inhibition in Mice", *Neuropsychopharmacology*, 04/2006

Publication

<% 1

74

www.karger.com

Internet Source

<% 1

75

Sestito, R. S., L. B. Trindade, R. G. de Souza, L. N. Kerbauy, M. M. Iyomasa, and M. L. Rosa.

<% 1