

# Variations in seasonal solar insolation are associated with a history of suicide attempts in bipolar I disorder

*by* Andi Jayalangkara

---

**Submission date:** 13-Mar-2022 08:50PM (UTC+0700)

**Submission ID:** 1783110636

**File name:** ted\_with\_a\_history\_of\_suicide\_attempts\_in\_bipolar\_I\_disorder.pdf (823.53K)

**Word count:** 11375

**Character count:** 62796

## RESEARCH

## Open Access

41

## Variations in seasonal solar insolation are associated with a history of suicide attempts in bipolar I disorder



Michael Bauer<sup>1\*</sup>, Tasha Glenn<sup>2</sup>, Eric D. Achtyes<sup>3</sup>, Martin Alda<sup>4</sup>, Esen Agaoglu<sup>5</sup>, Kürşat Altınbaş<sup>6</sup>, Ole A. Andreassen<sup>7</sup>, Elias Angelopoulos<sup>8</sup>, Raffaella Ardu<sup>9</sup>, Edgar Arrua Vares<sup>1</sup>, Memduha Aydin<sup>10</sup>, Yavuz Ayhan<sup>5</sup>, Christopher Baethge<sup>11</sup>, Rita Bauer<sup>1</sup>, Bernhard T. Baune<sup>12,13,14</sup>, Ceylan Balaban<sup>15</sup>, Claudia Becerra-Palars<sup>16</sup>, Aniruddh P. Behere<sup>17</sup>, Prakash B. Behere<sup>18</sup>, Habte Belete<sup>19</sup>, Tilahun Belete<sup>19</sup>, Gabriel Okawa Belizario<sup>20</sup>, Frank Bellivier<sup>21</sup>, Robert H. Belmaker<sup>22</sup>, Francesco Benedetti<sup>23,24</sup>, Michael Berk<sup>25,26</sup>, Yuly Bersudsky<sup>27</sup>, Şule Bicakci<sup>5,28</sup>, Harriet Birabwa-Oketcho<sup>29</sup>, Thomas D. Bjella<sup>7</sup>, Conan Brady<sup>30</sup>, Jorge Cabrera<sup>31</sup>, Marco Cappucciati<sup>32</sup>, Angela Marianne Paredes Castro<sup>25</sup>, Wei-Ling Chen<sup>33</sup>, Eric Y. Wo Cheung<sup>34</sup>, Silvia Chiesa<sup>32</sup>, Marie Crowe<sup>35</sup>, Alessandro Cuomo<sup>36</sup>, Sara Dallaspezia<sup>24</sup>, Maria Del Zompo<sup>9</sup>, Pratikumar Desai<sup>37</sup>, Seetal Dodd<sup>25,38</sup>, Markus Donix<sup>1</sup>, Bruno Etain<sup>21</sup>, Andrea Fagiolini<sup>36</sup>, Frederike T. Fellendorf<sup>39</sup>, Ewa Ferencztajn-Rochowiak<sup>40</sup>, Jess G. Fiedorowicz<sup>41</sup>, Kostas N. Fountoulakis<sup>42</sup>, Mark A. Frye<sup>43</sup>, Pierre A. Geoffroy<sup>44,45,46</sup>, Ana Gonzalez-Pinto<sup>47</sup>, John F. Gottlieb<sup>48</sup>, Paul Grof<sup>49</sup>, Bartholomeus C. M. Haarman<sup>50</sup>, Hirohiko Harima<sup>51</sup>, Mathias Hasse-Sousa<sup>52</sup>, Chantal Henry<sup>53,54</sup>, Lone Höfdding<sup>55</sup>, Josselin Houenou<sup>56,57</sup>, Massimiliano Imbesi<sup>32</sup>, Erkki T. Isometsä<sup>58,59</sup>, Maja Ivkovic<sup>60</sup>, Sven Janno<sup>61</sup>, Simon Johnsen<sup>62</sup>, Flávio Kapczinski<sup>52</sup>, Gregory N. Karakatsoulis<sup>42</sup>, Mathias Kardell<sup>63</sup>, Lars Vedel Kessing<sup>64</sup>, Seong Jae Kim<sup>65</sup>, Barbara König<sup>66</sup>, Timur L. Kot<sup>67</sup>, Michael Koval<sup>68</sup>, Mauricio Kunz<sup>52</sup>, Beny Lafer<sup>20</sup>, Mikael Landén<sup>63,69</sup>, Erik R. Larsen<sup>70</sup>, Melanie Lenger<sup>39</sup>, Ute Lewitzka<sup>1</sup>, Rasmus W. Licht<sup>71,72</sup>, Carlos Lopez-Jaramillo<sup>73</sup>, Alan MacKenzie<sup>74</sup>, Helle Østergaard Madsen<sup>75</sup>, Simone Alberte Kongstad A. Madsen<sup>62</sup>, Jayant Mahadevan<sup>76</sup>, Augustine Mahardika<sup>77</sup>, Mirko Manchia<sup>78,79,80</sup>, Wendy Marsh<sup>81</sup>, Monica Martinez-Cengotitabengoa<sup>82</sup>, Klaus Martiny<sup>75</sup>, Yuki Mashima<sup>83</sup>, Declan M. McLoughlin<sup>84</sup>, Ybe Meesters<sup>50</sup>, Ingrid Melle<sup>7</sup>, Fátima Meza-Urzuá<sup>16</sup>, k Yee Ming<sup>85</sup>, Scott Monteith<sup>86</sup>, Muthukumar Moorthy<sup>76</sup>, Gunnar Morken<sup>87,88</sup>, Enrica Mosca<sup>9</sup>, Anton A. Mozzhegorov<sup>89</sup>, Rodrigo Munoz<sup>90</sup>, Starlin V. Mythri<sup>91</sup>, Fethi Nacef<sup>92</sup>, Ravi K. Nadella<sup>76</sup>, Takako Nakanotani<sup>93</sup>, René Ernst Nielsen<sup>71,72</sup>, Claire O'Donovan<sup>4</sup>, Adel Omrani<sup>94</sup>, Yamima Osher<sup>27</sup>, Uta Ouali<sup>92</sup>, Maja Pantovic-Stefanovic<sup>60</sup>, Pornjira Pariwatcharakul<sup>95</sup>, Joanne Petite<sup>4</sup>, Andrea Pfennig<sup>1</sup>, Yolanda Pica Ruiz<sup>96</sup>, Maximilian Pilhatsch<sup>1,97</sup>, Marco Pinna<sup>79,98</sup>, Maurizio Pompili<sup>99</sup>, Richard Porter<sup>35</sup>, Danilo Quiroz<sup>100</sup>, Francisco Diego Rabelo-da-Ponte<sup>52</sup>, Raj Ramesar<sup>101</sup>, Natalie Rasgon<sup>102</sup>, Woraphat Ratta-apha<sup>95</sup>, Michaela Ratzenhofer<sup>39</sup>, Maria Redahan<sup>30</sup>, M. S. Reddy<sup>91</sup>, Andreas Reif<sup>15</sup>, Eva Z. Reininghaus<sup>39</sup>, Jenny Gringer Richards<sup>103</sup>, Philipp Ritter<sup>1</sup>, Janusz K. Rybakowski<sup>40</sup>, Leela Sathyaputri<sup>103</sup>, Ângela M. Scippa<sup>104</sup>, Christian Simhandl<sup>105</sup>, Emanuel Severus<sup>1</sup>, Daniel Smith<sup>106</sup>, José Smith<sup>107</sup>, Paul W. Stackhouse Jr.<sup>108</sup>

5

\*Correspondence: Michael.Bauer@uniklinikum-dresden.de

<sup>1</sup>Department of Psychiatry and Psychotherapy, Faculty of Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Dan J. Stein<sup>109</sup>, Kellen Stilwell<sup>37</sup>, Sergio Strejilevich<sup>107</sup>, Kuan-Pin Su<sup>110,111</sup>, Mythily Subramaniam<sup>112</sup>, Ahmad Hatim Sulaiman<sup>113</sup>, Kirsi Suominen<sup>114</sup>, Andi J. Tanra<sup>115</sup>, Yoshitaka Tatebayashi<sup>93</sup>, Wen Lin Teh<sup>112</sup>, Leonardo Tondo<sup>116,117</sup>, Carla Torrent<sup>118</sup>, Daniel Tuinstra<sup>37</sup>, Takahito Uchida<sup>83</sup>, Arne E. Vaaler<sup>87,88</sup>, Julia Veeh<sup>15</sup>, Eduard Vieta<sup>118</sup>, Biju Viswanath<sup>76</sup>, Maria Yoldi-Negrete<sup>119</sup>, Oguz Kaan Yalcinkaya<sup>5</sup>, Allan H. Young<sup>120</sup>, Yosra Zgueb<sup>92</sup> and Peter C. Whybrow<sup>121</sup>

## Abstract

**Background:** Bipolar disorder is associated with circadian disruption and a high risk of suicidal behavior. In a previous exploratory study of patients with bipolar I disorder, we found that a history of suicide attempts was associated with differences between winter and summer levels of solar insolation. The purpose of this study was to confirm this finding using international data from 42% more collection sites and 25% more countries.

**Methods:** Data analyzed were from 71 prior and new collection sites in 40 countries at a wide range of latitudes. The analysis included 4876 patients with bipolar I disorder, 45% more data than previously analyzed. Of the patients, 1496 (30.7%) had a history of suicide attempt. Solar insolation data, the amount of the sun's electromagnetic energy striking the surface of the earth, was obtained for each onset location (479 locations in 64 countries).

**Results:** This analysis confirmed results of the exploratory study with the same best model and slightly better statistical significance. There was a significant inverse association between a history of suicide attempts and the ratio of mean winter insolation to mean summer insolation (mean winter insolation/mean summer insolation). This ratio is largest near the equator which has little change in solar insolation over the year, and smallest near the poles where the winter insolation is very small compared to the summer insolation. Other variables in the model associated with an increased risk of suicide attempts were a history of alcohol or substance abuse, female gender, and younger birth cohort. The winter/summer insolation ratio was also replaced with the ratio of minimum mean monthly insolation to the maximum mean monthly insolation to accommodate insolation patterns in the tropics, and nearly identical results were found. All estimated coefficients were significant at  $p < 0.01$ .

**Conclusion:** A large change in solar insolation, both between winter and summer and between the minimum and maximum monthly insolation, may increase the risk of suicide attempts in bipolar I disorder. With frequent circadian rhythm dysfunction and suicidal behavior in bipolar disorder, greater understanding of the optimal roles of daylight and electric lighting in circadian entrainment is needed.

**Keywords:** Bipolar disorder, Suicide, Sunlight, Solar insolation, Psychiatry, Circadian, Seasonal variation

## Introduction

The risk of suicidal behavior for those with bipolar disorder is estimated to be 20–30 times higher than for the general population (Pompili et al. 2013; Shaffer 2015; Dong et al. 2019; Plans et al. 2019). Risk factors for suicidal behavior in bipolar disorder include depression, agitation, impulsivity, comorbid alcohol or substance abuse, prior suicidal acts, recent discharge from a psychiatric hospital, along with genetic, demographic, sociodemographic and cultural factors, and stressful life events (Pompili et al. 2013; Shaffer 2015; Tondo et al. 2021; Bachmann 2018; Plans et al. 2019; Tidemalm et al. 2014). Additionally, international epidemiology studies of the general population spanning several decades report seasonality in suicide attempts and deaths with a peak in spring or summer (Galvão et al. 2018; Woo et al. 2012; Su et al. 2020; Postolache et al. 2010; Oladunjoye et al. 2020;

Christodoulou et al. 2012; Coimbra et al. 2016; Petridou et al. 2002).

There is increasing recognition of the profound and diverse impacts of daylight on human physiology and behavior, and the complexity of the mechanisms underlying the human response to light (Münch et al. 2017; Aranda and Schmidt 2021; Foster 2020). In addition to vision, daylight modulates circadian timing, the sleep-wake cycle, daily neuroendocrine functions, alertness, performance, mood and thermoregulation (Fitz-Jeffries et al. 2020; Paul and Brown 2019; Prayag et al. 2019; Cajochen 2007; Fisk et al. 2018; LeGates et al. 2014). Alterations in circadian rhythm are a major component of mood disorders (Logan and McClung 2019; Jones and Benca 2015; McClung 2013; Ketchesin et al. 2020), with disruptions in sleep, hormonal secretion, mood

regulation and social rhythms occurring frequently in bipolar disorder (Melo et al. 2017; Takaesu 2018; McCarthy 2019; Gonzalez 2014). The effects of circadian disruptions in bipolar disorder are interrelated and can both trigger and exacerbate symptoms (Harvey 2008; Walker et al. 2020; Geoffroy 2018). About 25% of patients exhibit a seasonal pattern in the course of bipolar disorder (Geoffroy et al. 2014; Mani et al. 2018).

In a prior exploratory study of patients with bipolar I disorder, we found that a history of suicide attempts was associated with living in locations with a large change in solar insolation between winter and summer (Bauer et al. 2019). Solar insolation (incoming solar radiation) is defined as the amount of electromagnetic energy from the sun striking a surface area on earth (Stackhouse et al. 2018). The aim of the current study was to investigate whether a repeat analysis with more data would confirm or contradict the results of the exploratory study. In addition, an analysis using the ratio of the minimum mean monthly insolation to the maximum mean monthly insolation was added to accommodate the insolation patterns in the tropics. The prior analysis included data from 50 collection sites in 32 countries. This analysis used 45% more data both from new and prior collection sites, including data from 71 collection sites in 40 countries with diverse cultures, healthcare systems, and climates.

## Methods

### Data collection

Data were collected by direct questioning, reviewing records, or both. All patients had a diagnosis of bipolar disorder from a psychiatrist according to DSM-IV or DSM-5 criteria. Study approval was obtained from local institutional review boards, following local requirements. This analysis includes the data used in the exploratory study that were collected between 2010 and 2017 and additional data collected between 2019 and 2020. Details about the project methodology were published previously (Bauer et al. 2012, 2014, 2017).

### Data collection sites

Researchers from 71 collection sites in 40 countries provided the data, including those at university medical centers, specialty clinics and individual practitioners. Collection sites located in the northern hemisphere were: Aalborg, Denmark; Aarhus, Denmark; Ankara, Turkey; Athens, Greece; Bangkok, Thailand; Barcelona, Spain; Barhir Dar, Ethiopia; Beer Sheva, Israel; Belgrade, Serbia; Bengaluru, India; Cagliari, Sardinia, Italy (2 sites); Calgary, Canada; Dresden, Germany; Dublin, Ireland; Frankfurt, Germany; Halifax, Canada; Helsinki, Finland; Glasgow, UK; Gothenburg, Sweden; Grand Rapids, MI, USA; Hong Kong, China; Hyderabad, India;

Iowa City, Iowa, USA; Jincheon, South Korea; Kampala, Uganda; Kansas City, KS, USA; Khanti-Mansiysk, Russia; Konya, Turkey; Kuala Lumpur, Malaysia; Los Angeles, CA, USA; Medellín, Colombia; Mexico City, Mexico; Milan, Italy; Oslo, Norway; Ottawa, Canada; Piacenza, Italy; Palo Alto, CA, USA; Paris, France (2 sites); Poznan, Poland; Rochester, MN, USA; Rome, Italy; San Diego, CA, USA; Siena, Italy; Singapore; Stockholm, Sweden; Tartu, Estonia; Thessaloniki, Greece (2 sites); Tokyo, Japan (3 sites); Taichung, Taiwan; Trondheim, Norway; Tunis, Tunisia; Vitoria, Spain; Wardha, India; Wiener Neustadt, Austria; Worcester, MA, USA, and Würzburg, Germany. Collection sites located in the southern hemisphere were: Adelaide, Australia; Melbourne/Geelong, Australia; Buenos Aires, Argentina; Cape Town, South Africa; Christchurch, New Zealand; Mataram, Indonesia; Porto Alegre, Brazil; Salvador, Brazil; Santiago, Chile (2 sites); and São Paulo, Brazil.

### Patient data collected

To facilitate international participation, minimal clinical data were collected for each patient. The patient data collected included gender, age of onset, polarity of first episode, family history of mood disorders, history of psychosis, episode course, history of alcohol and substance abuse, and history of suicide attempts. Three locations were also collected for each patient: birth location, onset location and current location. The same birth cohort groups were used as in the exploratory analysis, and in prior research (Bauer et al. 2014, 2015, 2017; Chengappa et al. 2003).

### Country specific data

Country specific socioeconomic data were obtained for all onset locations, including physician density per 1000 population, country median age, unemployment rate, poverty rate, gross domestic product (GDP) per capita (CIA World Factbook 2020), psychiatrists per 100,000 (WHO 2019a), Gini index of income inequality, percent Internet users (World Bank 2020a, b), gender inequality index (UN 2020), and if the country has a state-sponsored or officially favored religion (Pew Research 2017).

### Solar insolation

The NASA POWER database provides average monthly solar insolation expressed in kilowatt hours/square meter/day (kWh/m<sup>2</sup>/day) based on satellite observations collected between 1983 and the present (Stackhouse et al. 2018; NASA 2020). As in the exploratory study, a 22-year climatology of insolation spanning Jan 1984–December 2013 at spatial resolution of 1° × 1° latitude/longitude was used in this analysis. The actual onset locations were grouped into reference onset locations representing all

onset locations within a  $1^\circ \times 1^\circ$  grid of latitude and longitude. For example, Dresden, Germany at latitude of  $51.1^\circ$  north and  $13.8^\circ$  east is the reference onset location for all locations between  $51^\circ$  and  $52^\circ$  north, and  $13^\circ$  and  $14^\circ$  east. The latitude and longitude of the reference onset location were used to identify solar insolation values for each patient.

During a year, the pattern of mean monthly solar insolation varies by latitude, with little change near the equator and large changes near the north and south poles. Solar insolation values for locations at the same latitude may vary significantly due to local conditions including cloud cover, aerosols (including dust and pollution), water vapor amounts, and altitude. Locations in the tropics (less than  $23.5^\circ$  north or south of the equator), may have a wet season where clouds reduce solar insolation and a dry season with clear skies rather than a winter/summer insolation pattern. To summarize the changes in solar insolation throughout the year at each reference onset location, two variables were created: (1) the ratio of the mean northern hemisphere winter (December, January, February) to the mean summer (June, July, August) insolation, and (2) the ratio of the minimum mean monthly insolation to the maximum mean monthly insolation. The insolation data from the southern hemisphere were shifted by 6 months for comparison to data from the northern hemisphere to account for the seasonal cycle.

### Statistics

The same statistical approach was used as the exploratory study (Bauer et al. 2019), in which the generalized estimating equations (GEE) statistical technique was used to account for both the correlated data and unbalanced number of patients at reference onset locations. The GEE technique estimates the dependent variable as a function of the entire population, producing a population averaged or marginal estimates of model coefficient (Zeger and Liang 1986). All GEE models in this study were estimated using a binomial distribution, an exchangeable working correlation matrix and a logit link function where a patient history of suicide attempts was the dependent binary variable. The selection process of the best model of a history of suicide attempts first identified individual independent variables with an estimated coefficient significant at the 0.05 level in a univariate GEE model. Significant independent variables from univariate models and variables found in prior suicide research were then combined into multivariate GEE models of a history of suicide attempts. To identify the best model, the multivariate model estimates were compared using the corrected quasi-likelihood independence model criterion (Pan 2001) and confidence intervals at the 0.01 significance level to reduce the chance of type 1 errors. Based

on the logit link function, the exponential coefficient can be interpreted as the effect size (Li et al. 2019). Demographic variables were reported using descriptive statistics. SPSS version 26.0 was used for all analyses.

## Results

### Available data

Data for 10,771 patients were available from the 71 collection sites, including 3379 new patients, 46% more patients than in the exploratory analysis. Of these, 7844 patients had a diagnosis of bipolar I disorder. Of the 7844 patients with bipolar I disorder, a history of suicide attempts was available for 6064 patients. Of the 6064 patients with data on a history of suicide attempts, all 5 variables in the best model were only available for 4876 patients, with 81% of the excluded patients missing data for a history of alcohol or substance abuse. Although 19.6% of the patients with data on a history of suicide attempts were excluded, the other demographics were similar to those included in the best model. The demographics of the 4876 patients included in the best model are shown in Table 1. Of the 4876 patients, 2760 patients (56.6%) were female, and 1496 patients (30.7%) had a history of suicide attempts.

### Onset locations

For the 4876 patients analyzed in the best model, there were 479 reference onset locations in 64 countries. The onset location was in the northern hemisphere for 4176 patients (85.6%), and in the southern hemisphere for 700 patients (14.4%), similar to estimates that about 12.5% of the world population lives in the southern hemisphere (Kummu and Varis 2011). Of the 4876 patients, 912 (18.7%) had an onset location in the tropics. For the 4876 patients, 97.6% of the onset locations were in the same country as the current country, and 83.3% of the onset cities were the same as the current city. The average number of patients in each onset location was 10.2, with 256 (5.3%) of the 4876 patients in an onset location with a single patient. As with the exploratory study, the much larger number of onset locations than collection sites reflects worldwide urbanization (WHO 2019b). Figure 1 provides a comparison of the range of latitudes for the onset locations between this analysis and the exploratory study.


### Ratio of mean winter solar insolation to mean summer solar insolation

At locations near the equator, there is little change in solar insolation between winter and summer and the ratio of mean winter solar insolation to mean summer solar insolation is large (near 1). At locations near the poles, solar insolation is very small in winter when compared to

**Table 1** Demographics of the patients with bipolar I disorder (N = 4876)

Parameter	Value	N	%
Gender	Female	2760	56.6
	Male	2116	43.4
Polarity of first episode <sup>a</sup>	Manic/hypomanic	2302	48.8
	Depressed	2419	51.2
Family history of mood disorder <sup>a</sup>	No	2026	45.3
	Yes	2448	54.7
Alcohol or substance abuse	No	3369	69.1
	Yes	1507	30.9
State sponsored religion in country of onset	No	2662	54.6
	Yes	2214	45.4
History of suicide attempt	No	3380	69.3
	Yes	1496	30.7
Cohort group	DOB < 1940	179	3.7
	DOB ≥ 1940 and DOB < 1960	1241	25.5
	DOB ≥ 1960	3456	70.9
<b>Parameter</b>		<b>Mean</b>	<b>SD</b>
Age at time of data collection		47.8	14.4
Age of onset		25.7	10.6

<sup>a</sup> Missing values excluded



**Fig. 1** Comparison of range of onset location latitudes for current and exploratory analyses for patients with bipolar I disorder (N = 4876)

the summer, and the ratio is small (near 0). The ratio of mean winter solar insolation to mean summer solar insolation by latitude groups is shown in Table 2. The ratio of mean winter solar insolation to mean summer solar insolation for example onset locations in the latitude groups is shown in Table 3.

#### Best model results

The best fitting model for a history of suicide attempts uses the ratio of mean winter solar insolation to the mean summer solar insolation and is shown in Table 4. This is the same model that was selected in the exploratory study as the best model, and the estimated coefficients are similar in value to those in the exploratory analysis. The inclusion of 4876 patients in the best model was a 45% increase over the 3365 patients included in the exploratory analysis.

The estimated coefficients for the model suggest that the odds of a suicide attempt will decrease by 4.8% for every 0.1 increase in the ratio of mean winter to summer insolation. Alternatively stated, comparing a ratio of 1 (near the equator) to a ratio of 0 (near a pole), there was a 48% difference in the odds of a suicide attempt with the lowest odds at the equator. The model estimates that being male will decrease the odds of a suicide attempt by 54%, and living in a country with a state sponsored or favored religion will decrease the odds by 65%. The model also estimates that having a history of alcohol or substance abuse will increase the odds of a suicide attempt by 58%, and being in the youngest cohort will increase the odds of a suicide attempt by 127%.

#### Ratio of minimum mean monthly insolation to the maximum mean monthly insolation

The ratio of minimum mean monthly insolation to the maximum mean monthly insolation by latitude groups is shown in Table 2. The ratio of minimum mean

monthly insolation to the maximum mean monthly insolation for selected onset locations in the latitude groups is shown in Table 3. A second model that substituted the ratio of minimum mean monthly insolation to the maximum mean monthly insolation for the ratio of mean winter solar insolation to mean summer solar insolation was estimated using the same data as with the best model.

The estimated coefficients for the model using the minimum mean monthly insolation to the maximum mean monthly insolation are shown in Table 5, and are very similar to those in the best model. The estimated coefficients for the monthly model suggest that the odds of a suicide attempt will decrease by 4.4% for every 0.1 increase in the ratio of mean winter to summer insolation. Alternatively stated, comparing a ratio of 1 (near the equator) to a ratio of 0 (near a pole), there was a 44% difference in the odds of a suicide attempt with the lowest odds at the equator. The model estimates that being male will decrease the odds of a suicide attempt by 54%, and living in a country with a state sponsored or favored religion will decrease the odds by 69%. The model also estimates that having a history of alcohol or substance will increase the odds of a suicide attempt by 59%, and being in the youngest cohort will increase the odds of a suicide attempt by 124%.

The collection site was thought to be an adequate proxy for the onset location for some or all patients from Barcelona, Cape Town, Christchurch, Frankfurt, Helsinki, Melbourne/Geelong, Porto Alegre, São Paulo, Salvador, Vitoria, and Würzburg, where the patient onset location was not provided. To test the effect of using the current location as a proxy for the onset location, the best model and the minimum mean monthly insolation to the maximum mean monthly insolation model were also estimated excluding these patients. The magnitude of the estimated coefficients did not change substantially and

**Table 2** Ratio of mean winter solar insolation/mean summer solar insolation and ratio of minimum mean monthly insolation/maximum mean monthly insolation by latitude for patient onset locations (N = 4876)

Degrees latitude north + south	N	%	Ratio mean winter insolation/mean summer insolation	Ratio minimum mean monthly insolation/maximum mean monthly insolation
0–9	268	5.5	1.0313	0.8076
10–19	420	8.6	1.1074	0.6744
20–29	254	5.2	0.7772	0.6093
30–39	1333	27.3	0.4075	0.3165
40–49	1921	39.4	0.3023	0.2119
50–59	444	9.1	0.1662	0.0903
60+	236	4.8	0.0857	0.0220
Total	4876	100.0	0.4423	0.3135

**Table 3** Ratio of mean winter solar insolation/mean summer solar insolation and ratio of minimum mean monthly insolation/maximum mean monthly insolation: example onset locations by latitude group (N = 4876)

Degrees latitude north + south	Onset location	Ratio mean winter insolation/mean summer insolation	Ratio minimum mean monthly insolation/maximum mean monthly insolation
0–9	Kampala, Uganda	1.1400	0.8197
	Kuala Lumpur, Malaysia	0.9702	0.7694
	Mataram, Indonesia	1.0125	0.7831
	Medellin, Columbia	0.9065	0.8370
	Singapore	1.0560	0.7797
10–19	Bahir Dar, Ethiopia	1.1639	0.7713
	Bangkok, Thailand	1.0680	0.7207
	Bengaluru, India	1.1702	0.6814
	Hyderabad, India	1.1762	0.6421
	Mexico City, Mexico	0.9074	0.6855
	Salvador, Brazil	0.6844	0.6246
20–29	Hong Kong, China	0.6603	0.6016
	São Paulo, Brazil	0.7419	0.6050
	Taichung, Taiwan	0.4492	0.3931
	Wardha, India	1.1545	0.5750
30–39	Ankara, Turkey	0.3266	0.2374
	Athens, Greece	0.3148	0.2319
	Beer Sheva, Israel	0.4246	0.3556
	Buenos Aires, Argentina	0.3978	0.3149
	58 Jari, Italy	0.3066	0.2328
	Cape Town, South Africa	0.3873	0.3227
	Los Angeles, CA, USA	0.4235	0.3503
	Melbourne, Australia	0.3628	0.2913
	San Francisco, CA, USA	0.4163	0.3137
	Santiago, Chile	0.3537	0.2879
	Seoul, South Korea	0.6406	0.4404
	Tokyo, Japan	0.7201	0.5574
Tunis, Tunisia	0.3695	0.2859	
40–49	Belgrade, Serbia	0.2832	0.1960
	Barcelona, Spain	0.3622	0.2603
	Boston, MA, USA	0.3626	0.2662
	Christchurch, New Zealand	0.3225	0.2461
	Grand Rapids, MI, USA	0.3281	0.2256
	Halifax, Canada	0.3300	0.2270
	Minneapolis, MN, USA	0.3339	0.2371
	Paris, France	0.2317	0.1540
	Rome, Italy	0.2993	0.2203
	Siena, Italy	0.2988	0.2077
	Vienna, Austria	0.2631	0.1667
	Würzburg, Germany	0.2381	0.1477
	50–59	Aarhus, Denmark	0.1432
Calgary, Canada		0.2269	0.1454
Dresden, Germany		0.2255	0.1379
Dublin, Ireland		0.1927	0.1149
Oslo, Norway		0.1126	0.0433
Poznan, Poland		0.2127	0.1290
Stockholm, Sweden		0.1087	0.0427
Tartu, Estonia		0.1353	0.0562
60+	Helsinki, Finland	0.1095	0.0359
	Khanti-Mansiysk, Russia	0.0951	0.0243
	Trondheim, Norway	0.0673	0.0116

remained significant at the 0.01 level. Estimated models including other patient, country and solar insolation variables were not as significant, or not as meaningful.

### Discussion

This analysis confirmed the results of the exploratory study after including 45% more international patient data. Living in locations with a large change in solar

**Table 4** Estimated parameters for best model explaining a history of suicide attempts for patients with bipolar I disorder (N = 4876)

Parameters	Coefficient estimate ( $\beta$ )	Standard error	Exp ( $\beta$ )	99% Confidence interval		Coefficient significance	
				Lower	Upper	Wald Chi-squared	P
Intercept	-0.935	0.2279	0.393	-1.522	-0.348	16.815	<0.001
Ratio mean winter insolation/mean summer insolation	-0.730	0.1752	0.482	-1.181	-0.279	17.357	<0.001
State sponsored religion in onset country	-0.438	0.1145	0.645	-0.733	-0.143	14.655	<0.001
Male	-0.609	0.0792	0.544	-0.813	-0.405	59.096	<0.001
History of alcohol or substance abuse	0.459	0.0726	1.582	0.272	0.646	39.978	<0.001
DOB $\geq$ 1960	0.822	0.2289	2.275	0.232	1.414	12.890	<0.001 <sup>a</sup>
DOB $\geq$ 1940 and DOB < 1960	0.681	0.2064	1.975	0.149	1.212	10.872	0.001 <sup>a</sup>

Dependent variable: history of suicide attempts (yes/no). Model: intercept, ratio of mean winter insolation/mean summer insolation at onset location, gender, state sponsored religion in onset country (yes/no), alcohol or substance abuse (yes/no) and birth cohort group (DOB < 1940, DOB  $\geq$  1940 and DOB < 1960, DOB  $\geq$  1960)

<sup>a</sup> Individual parameters Wald Chi-square statistics and significance. The model effects Wald Chi-square and significance for the cohort parameter was 12.904 and 0.002, respectively with 2 degrees of freedom

**Table 5** Estimated parameters for alternative model explaining a history of suicide attempts for patients with bipolar I disorder (N = 4876)

Parameters	Coefficient estimate ( $\beta$ )	Standard error	Exp ( $\beta$ )	99% Confidence interval		Coefficient significance	
				Lower	Upper	Wald Chi-squared	P
Intercept	-1.026	0.2302	0.358	-1.619	-0.434	19.885	<0.001
Ratio minimum mean monthly insolation/maximum mean monthly insolation	-0.813	0.2552	0.444	-1.470	-0.155	10.136	0.001
State sponsored religion in onset country	-0.378	0.1127	0.685	-0.668	-0.088	11.252	0.001
Male	-0.612	0.0794	0.542	-0.816	-0.407	59.438	<0.001
History of alcohol or substance abuse	0.466	0.0730	1.594	0.278	0.655	40.760	<0.001
DOB $\geq$ 1960	0.808	0.2312	2.244	0.213	1.404	12.224	<0.001 <sup>a</sup>
DOB $\geq$ 1940 and DOB < 1960	0.679	0.2085	1.972	0.142	1.216	10.612	0.001 <sup>a</sup>

Dependent variable: history of suicide attempts (yes/no). Model: intercept, ratio minimum mean monthly insolation/maximum mean monthly insolation at onset location, gender, state sponsored religion in onset country (yes/no), alcohol or substance abuse (yes/no) and birth cohort group (DOB < 1940, DOB  $\geq$  1940 and DOB < 1960, DOB  $\geq$  1960)

<sup>a</sup> Individual parameters Wald Chi-square statistics and significance. The model effects Wald chi-square and significance for the cohort parameter was 12.224 and 0.002, respectively with 2 degrees of freedom

insolation between winter and summer was associated with increased history of suicide attempts in patients with bipolar I disorder. The onset locations in this analysis were distributed across all latitudes in both hemispheres, and represent a wide range of solar insolation profiles and climatic conditions. The exploratory study results were confirmed in this study in two ways: by identifying the same GEE model as the best model, and by estimating a nearly identical relationship between solar insolation and a history of suicide attempts with slightly better statistical significance. In addition, the estimated coefficients for all other contributing variables in the model, history of alcohol or substance abuse, female gender, birth cohort and

state sponsored religion, were similar and slightly more significant. The finding of nearly identical results with an alternative measure of variation in solar insolation, which applies to all locations including the tropics, further confirms the association between a change in solar insolation and a history of suicide attempts.

The largest change in solar insolation between winter and summer occurs at locations near the poles. Suicide is a serious public health problem in the 8 countries with Arctic communities above 60°N (Pollock et al. 2020; Young et al. 2015). For example, in 2017 the suicide rate for the state of Alaska was nearly double the US national suicide rate, and nearly triple for Alaska native people

(AK-IBIS 2019). Additionally, seasonality in suicide is associated with latitude, with little monthly variation or seasonality in suicide rates near the equator, and spring and summer peaks in suicide rates with increasing latitudes north or south (Davis and Lowell 2002; Schwartz 2019).

There is related evidence involving patterns of solar radiation from studies within individual countries. In Finland, an increased suicide risk was associated with the cumulative low solar radiation over the long northern winter (Ruuhela et al. 2009). Several studies reported that an increasing risk of suicidal behavior was associated with increasing solar radiation. In South Korea, increased solar radiation in spring and summer was associated with an increased suicide rate (Jee et al. 2017). In Germany and Greece, increased solar insolation may precede suicidal acts (Müller et al. 2011; Papadopoulos et al. 2005). In Italy, higher solar radiation was associated with an increase in patients admitted to an emergency psychiatric unit with a primary diagnosis of bipolar disorder (Aguglia et al. 2019).

#### Consistency with prior research

The demographics of the patients are consistent with prior international studies of bipolar disorder, 35% having a history of suicide attempts (Tondo et al. 2016; Dong et al. 2019; Bobo et al. 2018), and 32% a history of alcohol or substance abuse (Toftdahl et al. 2016; Hunt et al. 2016; Grant et al. 2012; Nesvåg et al. 2015). Although we previously found a strong, inverse relation between the maximum monthly increase in solar insolation in springtime and the age of onset of bipolar I disorder (Bauer et al. 2017, 2014, 2012), the unadjusted mean age of onset of 21 is also similar to international studies (Baldessarini et al. 2012; Morselli et al. 2003; Kalman et al. 2019).

The other variables included in the best model also agree with prior suicide research in bipolar disorder and the general population. Alcohol and substance abuse (Schaffer et al. 2015; Carrà et al. 2014; Østergaard et al. 2017; Bobo 2018; Yuodelis-Flores and Ries 2015; Westström and Rossow 2016), and being female (Schaffer et al. 2015; Dong et al. 2019; Tondo et al. 2016; Bobo 2018) are associated with an increased risk of suicidal behavior. Increased suicide attempts or deaths are reported internationally in younger birth cohorts (Twenge et al. 2019; Odagi et al. 2011; Page et al. 2013; Yu and Chen 2019; Kwon et al. 2009; Gunnell et al. 2003; Phillips 2014; Chung et al. 2016). Studies involving all major world religions find that religion may be protective against suicidal behavior (Eskin et al. 2020; Wu et al. 2015; Van Weele et al. 2016; Stack and Kposowa 2011; Dervic et al. 2011; Caribe et al. 2015; Jacob et al. 2019).

#### Specific importance of daylight

The findings of this study highlight the importance of daylight to human wellbeing and behavior. In repeated surveys, people preferred daylight over electric lighting as the source of illumination, although the reasons for the strong daylight preference are not fully established (Knoop et al. 2020; Boyce et al. 2003; Haans 2014). Daylight differs from electric lighting in many fundamental properties, including the spectrum, intensity, temporal characteristics, flicker, and polarization, and the properties of daylight change throughout the day, month and year (Knoop et al. 2020; Aaarts et al. 2017). Many additional factors influence the physiological effects of light. These include individual characteristics such as age, lifestyle, health status, and genetics, environmental issues such as the season, climate, latitude and building design, and the duration of exposure and prior light exposure (Münch et al. 2017, 2020; Turner and Mainster 2008; Prayag et al. 2019).

Researchers emphasize the need to better understand how people respond to light and electric lighting in real-life settings (Knoop et al. 2020; Webler et al. 2019; Münch et al. 2020; Foster et al. 2020). Knowledge of non-image forming visual functions including circadian entrainment has grown rapidly. However, many findings are from small studies of healthy young adults exposed to electric lighting in controlled settings, or from animal studies. Even in controlled settings, considerable individual variability in sensitivity to light was detected (Phillips et al. 2019; McGlashan et al. 2018; Chellepa 2020). Understanding of how light intensity and duration of exposure interact for circadian entrainment is limited (Foster et al. 2020). Studies are needed that measure naturally occurring entrainment in large numbers of people of all ages and occupations, including mixed exposure to daylight and electric light in the day as well as electric lighting at night (Knoop et al. 2020; Webler et al. 2019; Münch et al. 2020; Foster et al. 2020). It is also not clear how applicable these findings are to patients with bipolar disorder. The optimal mix of daylight and electric lighting for circadian entrainment needs to be clarified to increase understanding of bipolar disorder and suicide risk, and improve the efficacy of chronic therapeutic treatments (Geoffroy and Palagini 2021; Gottlieb et al. 2019; Münch et al. 2020; Wang et al. 2020; Wirz-Justice and Benedetti 2020).

#### Limitations

Data in this project were collected as a convenience sample. The diagnosis was based on DSM-IV or DSM-5 criteria, but data collection methods and sources were not standardized, including the definition of suicide attempts. Although convenience samples can contain inadvertent

biases, this study repeated the results of the exploratory study using substantially more international patient data. This suggests either sample biases in the exploratory study were duplicated in the data collection for this study from 71 international collection sites or, more likely, the relationship found between solar insolation and a history of suicide attempts was confirmed.

Although a large percentage of patients had the same onset city and current city (83.3%), and the same onset and current country (97.6%), there was no confirmation that the suicide attempt occurred at the onset location. There was no data on individual risk factors for suicide attempts, the phase of bipolar disorder when the suicide attempt occurred, or treatments received for bipolar disorder, including those that may lower the risk of suicide such as lithium. There was no data on suicide deaths. The risks for attempted versus completed suicides could not be analyzed, although there are known distinctions (Hansson et al. 2018; Nock et al. 2008). There was no data on individuals who did not seek treatment. There was no individual data on sun exposure, sun-related activities, or lifestyle issues such as shift work. This analysis does not demonstrate causality or predict individual behavior. Characteristics of the forms of electric lighting were not discussed and other environmental variables were not included. Data from the southern hemisphere were shifted by 6 months, disregarding cultural dimensions of seasonality. Religious and cultural differences may influence data collection related to suicide, and to alcohol and drug abuse. The premature mortality from general medical illness (Roshanaei-Moghaddam and Katon 2009), completed suicides, treatment dropout rates, and the increased rate of diagnosis of bipolar disorder over time (Blader and Carlson 2007) may bias findings related to the birth cohort.

We previously noted two issues related to solar insolation that should be investigated in relation to suicide attempts: the potential impacts of continuous low solar insolation in areas near the poles with winters that last longer than 3 months, and of regional variance in insolation that has occurred over decadal timeframes (Wild 2012). However, we felt it was important to first confirm the results of the exploratory study.

## Conclusion

A history of suicide attempts in patients with bipolar I disorder was associated with living in locations with a large change in solar insolation, both between winter and summer and between the minimum and maximum monthly values. Given the frequent presence of circadian rhythm dysfunction and suicidal behavior in bipolar disorder, and the fundamental importance of daylight to human health, greater understanding of the optimal roles

of daylight and electric lighting in circadian entrainment in both the normal population and bipolar disorder is needed.

## Acknowledgements

None.

## Authors' contributions

MB and TG completed the initial draft, which was reviewed by all authors. All authors read and approved the final manuscript.

## Funding

Open Access funding enabled and organized by Projekt DEAL. Michael Berk is supported by a NHMRC Senior Principal Research Fellowship (1156072). Pierre A. Geoffroy, Chantal Henry and Josselin Houenou received grants from the French Agence Nationale pour la Recherche (ANR-11-IDEX-0004 Labex BioPsy "Olfaction and Bipolar Disorder" collaborative project, ANR-10-COHO-10-01 psyCOH and ANR-DFG ANR-14-CE35-0035 FUNDO). Mok Ying, Mythily Subramaniam, and Wen Lin Teh received funding from the National Medical Research Centre (NMRC) Centre Grant (Ref No: NMRC/CG/M002/2017\_IMH).

## Availability of data and materials

The data will not be shared or made publicly available.

## Declarations

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

The authors provide consent for publication.

### Competing interests

Rasmus W. Licht has received research Grants from Glaxo Smith Kline, honoraria for lecturing from Pfizer, Glaxo Smith Kline, Eli Lilly, Astra-Zeneca, Bristol-Myers Squibb, Janssen Cilag, Lundbeck, Otsuka, Servier and honoraria from advisory board activity from Glaxo Smith Kline, Eli Lilly, Astra-Zeneca, Bristol-Myers Squibb, Janssen Cilag, Sunovion and Sage. All other authors report no competing interests.

## Author details

<sup>1</sup>Department of Psychiatry and Psychotherapy, Faculty of Medicine, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany. <sup>2</sup>ChronoRecord Association, Fullerton, CA, USA. <sup>3</sup>Division of Psychiatry and Behavioral Medicine, Michigan State University College of Human Medicine, Grand Rapids, MI, USA. <sup>4</sup>Department of Psychiatry, Dalhousie University, Halifax, NS, Canada. <sup>5</sup>Department of Psychiatry, Hacettepe University Faculty of Medicine, Ankara, Turkey. <sup>6</sup>Department of Psychiatry, Selcuk University Faculty of Medicine, Mazhar Osman Mood Center, Konya, Turkey. <sup>7</sup>NORMENT Centre, Division of Mental Health and Addiction, Oslo University Hospital and Institute of Clinical Medicine, University of Oslo, Oslo, Norway. <sup>8</sup>Department of Psychiatry, National and Capodistrian University of Athens, Athens, Greece. <sup>9</sup>Department of Psychiatry, National and Capodistrian University of Athens, Athens, Greece. <sup>10</sup>Department of Biomedical Sciences, University of Cagliari, Sardinia, Italy. <sup>11</sup>Department of Psychiatry, Selcuk University Faculty of Medicine, Konya, Turkey. <sup>12</sup>Department of Psychiatry and Psychotherapy, University of Cologne Medical School, Cologne, Germany. <sup>13</sup>Department of Psychiatry, University of Münster, Münster, Germany. <sup>14</sup>Department of Psychiatry, Melbourne Medical School, The University of Melbourne, Melbourne, Australia. <sup>15</sup>The Florey Institute of Neuroscience and Mental Health, The University of Melbourne, Parkville, VIC, Australia. <sup>16</sup>Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt, Johann Wolfgang Goethe-Universität Frankfurt am Main, Frankfurt am Main, Germany. <sup>17</sup>National Institute of Psychiatry "Ramón de la Fuente Muñiz", Mexico City, Mexico. <sup>18</sup>Child and Adolescent Psychiatry, Helen DeVos Children's Hospital, Michigan State University-CHM, Grand Rapids, MI, USA. <sup>19</sup>Department of Psychiatry, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences (Deemed University), Wardha, India. <sup>20</sup>Department of Psychiatry, College of Medicine

- 80
- 17 Health Sciences, Bahir Dar University, Bahir Dar, Ethiopia. <sup>20</sup> Bipolar Disorder Research Program, Department of Psychology, University of São Paulo Medical School, São Paulo, Brazil. <sup>21</sup> Département de Psychiatrie et de Médecine Addictologique, Assistance Publique-Hôpitaux de Paris, INSERM UMR-S1144, Université Paris, FondaMental Foundation, Paris, France. <sup>22</sup> Professor Emeritus of Psychiatry, Ben Gurion University of the Negev, Beer Sheva, Israel. <sup>23</sup> University Vita-Salute San Raffaele, Milan, Italy. <sup>24</sup> Psychiatry and Clinical Psychobiology, Division of Neuroscience, San Raffaele Scientific Institute, Milan, Italy. <sup>25</sup> Deakin University, IMPACT-The Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Barwon Health, Geelong, Australia. <sup>26</sup> Orygen, The National Centre of Excellence in Youth Mental Health, Centre for Youth Mental Health, Florey Institute for Neuroscience and Mental Health, Department of Psychiatry, The University of Melbourne, Melbourne, Australia. <sup>27</sup> Department of Psychiatry, Faculty of Health Sciences, Beer Sheva Mental Health Center, Ben Gurion University of the Negev, Beer Sheva, Israel. <sup>28</sup> Department of Psychiatry, Baskent University Faculty of Medicine, Ankara, Turkey. <sup>29</sup> Butabika Hospital, Kampala, Uganda. <sup>30</sup> Department of Psychiatry, Trinity College Dublin, St Patrick's University Hospital, Dublin, Ireland. <sup>31</sup> Mood Disorders Clinic, Dr. Jose Horowitz Psychiatric Institute, Santiago de Chile, Chile. <sup>32</sup> Department of Mental Health and Substance Abuse, Piacenza, Italy. <sup>33</sup> Department of Psychiatry, Chiayi Branch, Taichung Veterans General Hospital, Chiayi, Taiwan. <sup>34</sup> Private Practice, Central, Hong Kong. <sup>35</sup> Department of Psychological Medicine, University of Otago, Christchurch, New Zealand. <sup>36</sup> Department of Molecular Medicine, University of Siena School of Medicine, Siena, Italy. <sup>37</sup> West Christian Mental Health Services, Grand Rapids, MI, USA. <sup>38</sup> Department of Psychiatry, University of Melbourne, Parkville, VIC, Australia. <sup>39</sup> Department of Psychiatry and Psychosomatic Medicine, Medical University Graz, Graz, Austria. <sup>40</sup> Department of Adult Psychiatry, Poznan University of Medical Sciences, Poznan, Poland. <sup>41</sup> Department of Psychiatry, School of Epidemiology and Public Health, University of Ottawa, Ottawa, ON, Canada. <sup>42</sup> Department of Psychiatry, School of Medicine, Faculty of Health Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece. <sup>43</sup> Department of Psychiatry and Psychology, Mayo Clinic Depression Center, Mayo Clinic, Rochester, MN, USA. <sup>44</sup> Département de Psychiatrie et d'addictologie, AP-HP, Hôpital Bichat-Crochin, Paris, France. <sup>45</sup> GHU Paris-Psychiatry and Neurosciences, 75014 Paris, France. <sup>46</sup> Université de Paris, NeuroDiderot, Inserm, Paris, France. <sup>47</sup> BIOARABA, Department of Psychiatry, University Hospital of Alava, University of the Basque Country, CIBERSAM, Vitoria, Spain. <sup>48</sup> Department of Psychiatry, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA. <sup>49</sup> Mood Disorders Center of Ottawa and the Department of Psychiatry, University of Toronto, Ottawa, Canada. <sup>50</sup> Department of Psychiatry, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands. <sup>51</sup> Department of Psychiatry, Tokyo Metropolitan Matsuzawa Hospital, Setagaya, Tokyo, Japan. <sup>52</sup> Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil. <sup>53</sup> Department of Psychiatry, GHU Paris Psychiatrie et Neurosciences, 75020 Paris, France. <sup>54</sup> Université de Paris, 75006 Paris, France. <sup>55</sup> Department of Clinical Research, University of Southern Denmark, Odense, Denmark. <sup>56</sup> Université Paris Est Créteil, INSERM, IMRB, Translational Neuropsychiatry, Fondation FondaMental, 94010 Créteil, France. <sup>57</sup> Université Paris Saclay, INSERM, Neurospin, 91191 Gif-sur-Yvette, France. <sup>58</sup> Department of Psychiatry, University of Helsinki and Helsinki University Hospital, Helsinki, Finland. <sup>59</sup> National Institute for Health and Welfare, Helsinki, Finland. <sup>60</sup> University Clinical Center of Serbia, Clinic for Psychiatry, Belgrade, Serbia. <sup>61</sup> Department of Psychiatry, University of Tartu, Tartu, Estonia. <sup>62</sup> Unit for Psychiatric Research, Aalborg University Hospital, Aalborg, Denmark. <sup>63</sup> Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. <sup>64</sup> Copenhagen Affective Disorder Research Center (CADIC), Psychiatric Center Copenhagen, Rigshospitalet, Copenhagen, Denmark. <sup>65</sup> Department of Psychiatry, Cheongju Hospital, Cheongju, South Korea. <sup>66</sup> BIPOLAR Zentrum Wiener Neustadt, Wiener Neustadt, Austria. <sup>67</sup> Khanty-Mansiysk Clinical Psychoneurological Hospital, Khanty-Mansiysk, Russia. <sup>68</sup> Department of Neuroscience, Michigan State University, East Lansing, MI, USA. <sup>69</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. <sup>70</sup> Mental Health Department Odense University Clinic and Department of Regional Health Research, University of Southern Denmark, Esbjerg, Denmark. <sup>71</sup> Psychiatry, Aalborg University Hospital, Aalborg, Denmark. <sup>72</sup> Department of Clinical Medicine, Aalborg University, Aalborg, Denmark. <sup>73</sup> Mood Disorders Program, Hospital Universitario San Vicente Fundación, Research Group in Psychiatry, Department of Psychiatry, Faculty of Medicine, Universidad Antioquia, Medellín, Colombia. <sup>74</sup> Forensic Psychiatry, University of Glasgow, NHS Greater Glasgow and Clyde, Glasgow, UK. <sup>75</sup> Copenhagen University Hospital, Psychiatric Centre Copenhagen, Copenhagen, Denmark. <sup>76</sup> Department of Psychiatry, National Institute of Mental Health and Neuro Sciences (NIMHANS), Bengaluru, India. <sup>77</sup> Department of Psychiatry, Faculty of Medicine, Mataram University, Mataram, Indonesia. <sup>78</sup> Department of Pharmacology, Dalhousie University, Halifax, NS, Canada. <sup>79</sup> Section of Psychiatry, Department of Medical Science and Public Health, University of Cagliari, Cagliari, Italy. <sup>80</sup> Unit of Clinical Psychiatry, University Hospital Agency of Cagliari, Cagliari, Italy. <sup>81</sup> Department of Psychiatry, University of Massachusetts Medical School, Worcester, MA, USA. <sup>82</sup> Osakidetza, Basque Health Service, BioAraba Research Institute, University of the Basque Country, Vitoria, Spain. <sup>83</sup> Department of Psychiatry, Keio University School of Medicine, Tokyo, Japan. <sup>84</sup> Department of Psychiatry, Trinity College Institute of Neuroscience, Trinity College Dublin, St Patrick's University Hospital, Dublin, Ireland. <sup>85</sup> Department of General Psychiatry, Mood Disorders Unit, Institute of Mental Health, Singapore City, Singapore. <sup>86</sup> Michigan State University College of Human Medicine, Traverse City Campus, Traverse City, MI, USA. <sup>87</sup> Department of Mental Health, Norwegian University of Science and Technology-NTNU, Trondheim, Norway. <sup>88</sup> Department of Psychiatry, St. Olavs' University Hospital, Trondheim, Norway. <sup>89</sup> Soviet Psychoneurological Hospital, Ural, Russia. <sup>90</sup> Department of Psychiatry, University of California San Diego, San Diego, CA, USA. <sup>91</sup> Asha Bipolar Clinic, Asha Hospital, Hyderabad, Telangana, India. <sup>92</sup> Razi Hospital, Faculty of Medicine, University of Tunis-El Manar, Tunis, Tunisia. <sup>93</sup> Affective Disorders Research Project, Tokyo Metropolitan Institute of Medical Science, Setagaya, Tokyo, Japan. <sup>94</sup> Tunisian Bipolar Clinic, Érable Médical Cabinet 324, Lac 2, Tunis, Tunisia. <sup>95</sup> Department of Psychiatry, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand. <sup>96</sup> Hospital 'Ángeles del Pedregal', Mexico City, Mexico. <sup>97</sup> Department of Psychiatry and Psychotherapy, Elblandklinikum Radebeul, Radebeul, Germany. <sup>98</sup> Lucio Bini Mood Disorder Center, Cagliari, Italy. <sup>99</sup> Department of Neurosciences, Mental Health and Sensory Organs, Sant'Andrea Hospital, Sapienza University of Rome, Rome, Italy. <sup>100</sup> Department of Psychiatry, Diego Portales University, Santiago de Chile, Chile. <sup>101</sup> SA MRC Genomic and Precision Medicine Research Unit, Division of Human Genetics, Department of Pathology, Institute of Infectious Diseases and Molecular Medicine, University of Cape Town, Cape Town, South Africa. <sup>102</sup> Department of Psychiatry and Behavioral Sciences, Stanford School of Medicine, Palo Alto, CA, USA. <sup>103</sup> Departments of Psychiatry, Epidemiology, and Internal Medicine, Iowa Neuroscience Institute, The University of Iowa, Iowa City, IA, USA. <sup>104</sup> Department of Neuroscience and Mental Health, Federal University of Bahia, Salvador, Brazil. <sup>105</sup> Bipolar Zentrum Wierzbustadt, Sigmund Freud Privat Universität, Vienna, Austria. <sup>106</sup> Centre for Clinical Neuroscience, University of Edinburgh, Edinburgh, Scotland, UK. <sup>107</sup> Bipolar Disorder Program, Neuroscience Institute, Favaloro University, Buenos Aires, Argentina. <sup>108</sup> Science Directorate/Climate Science Branch, NASA Langley Research Center, Hampton, VA, USA. <sup>109</sup> Department of Psychiatry, MRC Unit on Risk and Resilience in Mental Disorders, University of Cape Town, Cape Town, South Africa. <sup>110</sup> College of Medicine, China Medical University (CMU), Taichung, Taiwan. <sup>111</sup> An-Nan Hospital, China Medical University, Tainan, Taiwan. <sup>112</sup> Research Division, Institute of Mental Health, Singapore, Singapore. <sup>113</sup> Department of Psychological Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia. <sup>114</sup> Department of Social Services and Health Care, Psychiatry, City of Helsinki, Helsinki, Finland. <sup>115</sup> Department of Psychiatry, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia. <sup>116</sup> McLean Hospital/Harvard Medical School, Boston, MA, USA. <sup>117</sup> Mood Disorder Lucio Bini Centers, Cagliari e Roma, Italy. <sup>118</sup> Clinical Institute of Neuroscience, Hospital Clinic, University of Catalonia, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain. <sup>119</sup> Subdirección de Investigaciones Clínicas, Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, Mexico City, Mexico. <sup>120</sup> Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. <sup>121</sup> Department of Psychiatry and Biobehavioral Sciences, Semel Institute for Neuroscience and Human Behavior, University of California Los Angeles (UCLA), Los Angeles, CA, USA.

Received: 6 May 2021 Accepted: 13 July 2021

Published online: 01 September 2021

References

Aarts MP, Brown S, Bueno B, Gjedde A, Mersch D, Münch M, et al. Reinventing daylight. In: Changing perspectives on daylight: Science, technology and culture. Sponsored supplement to Science/AAS; Science/AAAS Custom Publishing Office: Washington, DC, USA, 2017. p. 33–37.

Aguglia A, Serafini G, Escelsior A, Canepa G, Amore M, Maina G. Maximum temperature and solar radiation as predictors of bipolar patient admission in an emergency psychiatric ward. *Int J Environ Res Public Health*. 2019;16:1140.

AK-IBIS (Alaska Department of Health and Social Services). Complete health indicator report of suicide mortality rate—all ages. 2019. [http://ibis.dhss.alaska.gov/indicator/complete\\_profile/SuicDth.html](http://ibis.dhss.alaska.gov/indicator/complete_profile/SuicDth.html). Accessed 3 Feb 2021.

Aranda ML, Schmidt TM. Diversity of intrinsically photosensitive retinal ganglion cells: circuits and functions. *Cell Mol Life Sci*. 2021;78889–907.

Bachmann S. Epidemiology of suicide and the psychiatric perspective. *Int J Environ Res Public Health*. 2018;15:1425.

Baldessarini RJ, Tondo L, Vazquez GH, Undurraga J, Bolzani L, Yildiz A, et al. Age at onset versus family history and clinical outcomes in 1,665 international bipolar-I disorder patients. *World Psychiatry*. 2012;11:40–6.

Bauer M, Glenn T, Alda M, Andreassen OA, Ardu R, Bellivier F, et al. Impact of sunlight on the age of onset of bipolar disorder. *Bipolar Disord*. 2012;14:654–63.

Bauer M, Glenn T, Alda M, Andreassen OA, Angelopoulos E, Ardu R, et al. Relationship between sunlight and the age of onset of bipolar disorder: an international multisite study. *J Affect Disord*. 2014;167:104–11.

Bauer M, Glenn T, Alda M, Andreassen OA, Angelopoulos E, Ardu R, et al. Influence of birth cohort on age of onset cluster analysis in bipolar I disorder. *Eur Psychiatry*. 2015;30:99–105.

Bauer M, Glenn T, Alda M, Aleksandrovich MA, Andreassen OA, Angelopoulos E, et al. Solar insolation in springtime influences age of onset of bipolar I disorder. *Acta Psychiatr Scand*. 2017;136:571–82.

Bauer M, Glenn T, Alda M, Andreassen OA, Angelopoulos E, Ardu R, et al. Association between solar insolation and a history of suicide attempts in bipolar I disorder. *J Psychiatr Res*. 2019;113:1–9.

Blader JC, Carlson GA. Increased rates of bipolar disorder diagnoses among US child, adolescent, and adult inpatients, 1996–2004. *Biol Psychiatry*. 2007;62:107–14.

Bobo WW, Na PJ, Geske JR, McElroy SL, Frye MA, Biernacka JM. The relative influence of individual risk factors for attempted suicide in patients with bipolar I versus bipolar II disorder. *J Affect Disord*. 2018;225:489–94.

Boyce P, Hunter C. Howlett O (2003) The benefits of daylight through windows. Troy, New York: Rensselaer Polytechnic Institute; 2003.

Cajochen C. Alerting effects of light. *Sleep Med Rev*. 2007;11:453–64.

Caribé AC, Studart P, Bezerra-Filho S, Brietzke E, Nunes Noto M, Vianna-Sulzbach M, et al. Is religiosity a protective factor against suicidal behavior in bipolar I outpatients? *J Affect Disord*. 2015;186:156–61.

Carrà G, Bartoli F, Crocarno C, Brady KT, Clerici M. Attempted suicide in people with co-occurring bipolar and substance use disorders: systematic review and meta-analysis. *J Affect Disord*. 2014;167:125–35.

Chellappa SL. Individual differences in light sensitivity affect sleep and circadian rhythms. *Sleep*. 2020;44:zsa214. <https://doi.org/10.1093/sleep/zsaa214>.

Chengappa KN, Kupfer DJ, Frank E, Houck PR, Grochocinski VJ, Cluss PA, et al. Relationship of birth cohort and early age at onset of illness in a bipolar disorder case registry. *Am J Psychiatry*. 2003;160:1636–42.

Christodoulou C, Douzenis A, Papadopoulos FC, Papadopoulou A, Bouras G, Gournellis R, et al. Suicide and seasonality. *Acta Psychiatr Scand*. 2012;125:127–46.

Chung RY, Yip BH, Chan SS, Wong SY. Cohort effects of suicide mortality are sex specific in the rapidly developed Hong Kong Chinese population, 1976–2010. *Depress Anxiety*. 2016;33:558–66.

CIA. The World Factbook. 2020. <https://www.cia.gov/the-world-factbook/>. Accessed 3 Feb 2021.

Coimbra DG, Pereira E, Silva PEAC, de Sousa-Rodrigues CF, Barbosa FT, de Siqueira Figueredo D, et al. Do suicide attempts occur more frequently in the spring too? A systematic review and rhythmic analysis. *J Affect Disord*. 2016;196:125–37.

Davis GE, Lowell WE. Evidence that latitude is directly related to variation in suicide rates. *Can J Psychiatry*. 2002;47:572–4.

Dervic K, Carballo JJ, Baca-Garcia E, Galfalvy HC, Mann JJ, Brent DA, et al. Moral or religious objections to suicide may protect against suicidal behavior in bipolar disorder. *J Clin Psychiatry*. 2011;72:1390–6.

Dong M, Lu L, Zhang L, Zhang Q, Ungvari GS, Ng CH, et al. Prevalence of suicide attempts in bipolar disorder: a systematic review and meta-analysis of observational studies. *Epidemiol Psychiatr Sci*. 2019;29:e63.

Eskin M, Baydar N, El-Nayal M, Asad N, Noor IM, Rezaeian M, et al. Associations of religiosity, attitudes towards suicide and religious coping with suicidal ideation and suicide attempts in 11 muslim countries. *Soc Sci Med*. 2020;265:113390.

Fisk AS, Tam SKE, Brown LA, Vyazovskiy VV, Bannerman DM, Peirson SN. Light and cognition: roles for circadian rhythms, sleep, and arousal. *Front Neurol*. 2018;9:56.

Foster RG, Hughes S, Peirson SN. Circadian photoentrainment in mice and humans. *Biology (basel)*. 2020;9:180.

Galvão PVM, Silva HRSE, Silva CMFPD. Temporal distribution of suicide mortality: a systematic review. *J Affect Disord*. 2018;228:132–42.

Geoffroy PA. Clock genes and light signaling alterations in bipolar disorder: when the biological clock is off. *Biol Psychiatry*. 2018;84:775–7.

Geoffroy PA, Palagini L. Biological rhythms and chronotherapeutics in depression. *Prog Neuropsychopharmacol Biol Psychiatry*. 2021;106:110158.

Geoffroy PA, Bellivier F, Scott J, Etain B. Seasonality and bipolar disorder: a systematic review, from admission rates to seasonality of symptoms. *J Affect Disord*. 2014;168:210–23.

Gonzalez R. The relationship between bipolar disorder and biological rhythms. *J Clin Psychiatry*. 2014;75:e323–31.

Gottlieb JF, Benedetti F, Geoffroy PA, Henriksen TEG, Lam RW, Murray G, et al. The chronotherapeutic treatment of bipolar disorders: a systematic review and practice recommendations from the ISBD task force on chronotherapy and chronobiology. *Bipolar Disord*. 2019;21:741–73.

Grant BF, Stinson FS, Dawson DA, Chou SP, Dufour MC, Compton W, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2004;61:807–16.

Gunnell D, Middleton N, Whitley E, Dorling D, Frankel S. Influence of cohort effects on patterns of suicide in England and Wales, 1950–1999. *Br J Psychiatry*. 2003;182:164–70.

Haans A. The natural preference in people's appraisal of light. *J Environ Psychol*. 2014;39:51–61.

Hansson C, Joas E, Pålsson E, Hawton K, Runeson B, Landén M. Risk factors for suicide in bipolar disorder: a cohort study of 12 850 patients. *Acta Psychiatr Scand*. 2018;138:456–63.

Harvey AG. Sleep and circadian rhythms in bipolar disorder: seeking synchrony, harmony, and regulation. *Am J Psychiatry*. 2008;165:820–9.

Hunt GE, Malhi GS, Cleary M, Lai HM, Sitharthan T. Comorbidity of bipolar and substance use disorders in national surveys of general populations, 1990–2015: systematic review and meta-analysis. *J Affect Disord*. 2016;206:321–30.

Jacob L, Haro JM, Koyanagi A. The association of religiosity with suicidal ideation and suicide attempts in the United Kingdom. *Acta Psychiatr Scand*. 2019;139:164–73.

Jee HJ, Cho CH, Lee YJ, Choi N, An H, Lee HJ. Solar radiation increases suicide rate after adjusting for other climate factors in South Korea. *Acta Psychiatr Scand*. 2017;135:219–27.

Jones SG, Bencs RM. Circadian disruption in psychiatric disorders. *Sleep Med Clin*. 2015;10:481–93.

Kalman JL, Papiol S, Forstner AJ, Heilbronner U, Degenhardt F, Strohmaier J, et al. Investigating polygenic burden in age at disease onset in bipolar disorder: findings from an international multicentric study. *Bipolar Disord*. 2019;21:68–75.

Ketchesin KD, Becker-Krail D, McClung CA. Mood-related central and peripheral clocks. *Eur J Neurosci*. 2020;51:326–45.

Knoop M, Stefani O, Bueno B, Matusiak B, Hobday R, Wirz-Justice A, et al. Daylight: what makes the difference? *Light Res Technol*. 2020;52:423–42.

Kummu M, Varis O. The world by latitudes: a global analysis of human population, development level and environment across the north–south axis over the past half century. *Appl Geogr*. 2011;31:495–507.

Kwon JW, Chun H, Cho SI. A closer look at the increase in suicide rates in South Korea from 1986–2005. *BMC Public Health*. 2009;9:72.

- LeGates TA, Fernandez DC, Hattar S. Light as a central modulator of circadian rhythms, sleep and affect. *Nat Rev Neurosci*. 2014;15:443–54.
- Li F, Forbes AB, Turner EL, Preisser JS. Power and sample size requirements for GEE analyses of cluster randomized crossover trials. *Stat Med*. 2019;38:636–49.
- Logan RW, McClung CA. Rhythms of life: circadian disruption and brain disorders across the lifespan. *Nat Rev Neurosci*. 2019;20:49–65.
- Maruani J, Anderson G, Etain B, Lejoux M, Bellivier F, Geoffroy PA. The neurobiology of adaptation to seasons: relevance and correlations in bipolar disorders. *Chronobiol Int*. 2018;35:1335–53.
- McCarthy MJ. Missing a beat: assessment of circadian rhythm abnormalities in bipolar disorder in the genomic era. *Psychiatr Genet*. 2019;29:29–36.
- McClung CA. How might circadian rhythms control mood? Let me count the ways. *Biol Psychiatry*. 2013;74:242–9.
- McGlashan EM, Poudel GR, Vidafar P, Drummond SPA, Cain SW. Imaging individual differences in the response of the human suprachiasmatic area to light. *Front Neurol*. 2018;9:1022.
- Melo MCA, Abreu RLC, Linhares Neto VB, de Bruin PFC, de Bruin VMS. Chronotype and circadian rhythm in bipolar disorder: a systematic review. *Sleep Med Rev*. 2017;34:46–58.
- Morselli PL, Elgie R. GAMIAN-Europe. GAMIAN-Europe/BEAM survey I-global analysis of a patient questionnaire circulated to 3450 members of 12 European advocacy groups operating in the field of mood disorders. *Bipolar Disord*. 2003;5:265–78.
- Müller H, Biermann T, Renk S, Reulbach U, Ströbel A, Kornhuber J, et al. Higher environmental temperature and global radiation are correlated with increasing suicidality—a localized data analysis. *Chronobiol Int*. 2011;28:949–57.
- Münch M, Brändstedt AE, Brown SA, Gjedde A, Kantermann T, Martiny K, et al. The effect of light on humans. In: *Changing perspectives on daylight: Science, technology and culture*. Sponsored supplement to *Science/AAS; Science/AAAS Custom Publishing Office: Washington, DC, USA*, 2017. pp. 16–23.
- Münch M, Wirz-Justice A, Brown SA, Kantermann T, Martiny K, Stefani O, et al. The role of daylight for humans: gaps in current knowledge. *Clocks Sleep*. 2020;2:61–85.
- NASA. NASA Power. Solar Data Overview. 2020. <https://power.larc.nasa.gov/docs/methodology/solar/>. Accessed 3 Feb 2021.
- Nesvåg R, Knudsen GP, Bakken IJ, Høyve A, Ystrom E, Surén P, et al. Substance use disorders in schizophrenia, bipolar disorder, and depressive illness: a registry-based study. *Soc Psychiatry Psychiatr Epidemiol*. 2015;50:1267–76.
- Nock MK, Borges G, Bromet EJ, Cha CB, Kessler RC, Lee S. Suicide and suicidal behavior. *Epidemiol Rev*. 2008;30:133–54.
- Norström T, Rossow I. Alcohol consumption as a risk factor for suicidal behavior: a systematic review of associations at the individual and at the population level. *Arch Suicide Res*. 2016;20:489–506.
- Odagiri Y, Uchida H, Nakano M. Gender differences in age, period, and birth-cohort effects on the suicide mortality rate in Japan, 1985–2006. *Asia Pac J Public Health*. 2011;23:581–7.
- Oladunjoye AQ, Oladunjoye OO, Ayeni OA, Olubiyi O, Fuchs A, Gurski J, et al. Seasonal trends in hospitalization of attempted suicide and self-inflicted injury in United States adults. *Cureus*. 2020;12:e10830.
- Østergaard MLD, Nordentoft M, Hjorthøj C. Associations between substance use disorders and suicide or suicide attempts in people with mental illness: a Danish nation-wide, prospective, register-based study of patients diagnosed with schizophrenia, bipolar disorder, unipolar depression or personality disorder. *Addiction*. 2017;112:1250–9.
- Page A, Milner A, Morrell S, Taylor R. The role of under-employment and unemployment in recent birth cohort effects in Australian suicide. *Soc Sci Med*. 2013;93:155–62.
- Pan W. Akaike's information criterion in generalized estimating equations. *Biometrics*. 2001;57:120–5.
- Papadopoulos FC, Frangakis CE, Skalkidou A, Petridou E, Stevens RG, Trichopoulos D. Exploring lag and duration effect of sunshine in triggering suicide. *J Affect Disord*. 2005;88:287–97.
- Paul S, Brown T. Direct effects of the light environment on daily neuroendocrine control. *J Endocrinol*. 2019 Aug 1;JOE-19-0302.R1.
- Petridou E, Papadopoulos FC, Frangakis CE, Skalkidou A, Trichopoulos D. A role of sunshine in the triggering of suicide. *Epidemiology*. 2002;13:106–9.
- Pew Research Center, Oct. 3, 2017, "Many countries favor specific religions, officially or unofficially". <http://www.pewforum.org/2017/10/03/many-countries-favor-specific-religions-officially-or-unofficially/>. Accessed 3 Feb 2021.
- Phillips JA. A changing epidemiology of suicide? The influence of birth cohorts on suicide rates in the United States. *Soc Sci Med*. 2014;114:151–60.
- Phillips AJK, Vidafar P, Burns AC, McGlashan EM, Anderson C, Rajaratnam SMW, et al. High sensitivity and interindividual variability in the response of the human circadian system to evening light. *Proc Natl Acad Sci USA*. 2019;116:12019–24.
- Plans L, Barrot C, Nieto E, Rios J, Schulze TG, Papiol S, et al. Association between completed suicide and bipolar disorder: a systematic review of the literature. *J Affect Disord*. 2019;242:111–22.
- Pollock NJ, Apok C, Concepcion T, Delgado RA Jr, Rasmus S, Chatwood S, et al. Global goals and suicide prevention in the Circumpolar North. *Indian J Psychiatry*. 2020;62:7–14.
- Pompili M, Gonda X, Serafini G, Innamorati M, Sher L, Amore M, et al. Epidemiology of suicide in bipolar disorders: a systematic review of the literature. *Bipolar Disord*. 2013;15:457–90.
- Postolache TT, Mortensen PB, Tonelli LH, Jiao X, Frangakis C, Soriano JJ, et al. Seasonal spring peaks of suicide in victims with and without prior history of hospitalization for mood disorders. *J Affect Disord*. 2010;121:88–93.
- Prayag AS, Münch M, Aeschbach D, Chellappa SL, Gronfier C. Light modulation of human clocks, wake, and sleep. *Clocks Sleep*. 2019;1:193–208.
- Roshanaei-Moghaddam B, Katon W. Premature mortality from general medical illnesses among persons with bipolar disorder: a review. *Psychiatr Serv*. 2009;60:147–56.
- Ruuhela R, Hiltunen L, Venäläinen A, Pirinen P, Partonen T. Climate impact on suicide rates in Finland from 1971 to 2003. *Int J Biometeorol*. 2009;53:167–75.
- Schaffer A, Isometsä ET, Tondo L, Moreno H, D, Turecki G, Reis C., et al. International Society for Bipolar Disorders Task Force on Suicide: meta-analyses and meta-regression of correlates of suicide attempts and suicide deaths in bipolar disorder. *Bipolar Disord*. 2015;17:1–16.
- Schwartz PJ. Chris Cornell, the black hole sun, and the seasonality of suicide. *Neuropsychobiology*. 2019;78:38–47.
- Stack S, Kposowa AJ. Religion and suicide acceptability: a cross-national analysis. *J Sci Study Relig*. 2011;50:289–306.
- Stackhouse, PW Jr, Zhang T, Westberg D, Barnett AJ, Bristow T, Macpherson B, Hoell JM. NASA POWER (Prediction of Worldwide Energy Resources). Release 8.01 (with GIS Applications) Methodology (Data Parameters, Sources, & Validation). 2018 <https://power.larc.nasa.gov>. Accessed 3 Feb 2021.
- Su MK, Chan PY, Hoffman RS. The seasonality of suicide attempts: a single poison control center perspective. *Clin Toxicol (Phila)*. 2020;58:1034–41.
- Takaesu Y. Circadian rhythm in bipolar disorder: a review of the literature. *Psychiatry Clin Neurosci*. 2018;72:673–82.
- Tidemalm D, Haglund A, Karanti A, Landén M, Runeson B. Attempted suicide in bipolar disorder: risk factors in a cohort of 6086 patients. *PLoS ONE*. 2014;9:e94097.
- Toftdahl NG, Nordentoft M, Hjorthøj C. Prevalence of substance use disorders in psychiatric patients: a nationwide Danish population-based study. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51:129–40.
- Tondo L, Pompili M, Forte A, Baldessarini RJ. Suicide attempts in bipolar disorders: comprehensive review of 101 reports. *Acta Psychiatr Scand*. 2016;133:174–86.
- Tondo L, Vázquez GH, Baldessarini RJ. Prevention of suicidal behavior in bipolar disorder. *Bipolar Disord*. 2021;23:14–23.
- Turner PL, Mainster MA. Circadian photoreception: ageing and the eye's important role in systemic health. *Br J Ophthalmol*. 2008;92:1439–44.
- Twenge JM, Cooper AB, Joiner TE, Duffy ME, Binau SG. Age, period, and cohort trends in mood disorder indicators and suicide-related outcomes in a nationally representative dataset, 2005–2017. *J Abnorm Psychol*. 2019;128:185–99.
- UN. Gender Inequality Index (GII). 2020. <http://hdr.undp.org/en/content/gender-inequality-index-gii>. Accessed 3 Feb 2021.
- VanderWeele TJ, Li S, Tsai AC, Kawachi I. Association between religious service attendance and lower suicide rates among US women. *JAMA Psychiatr*. 2016;73:845–51.

- Walker WH 2nd, Walton JC, DeVries AC, Nelson RJ. Circadian rhythm disruption and mental health. *Transl Psychiatry*. 2020;10:28.
- Wang S, Zhang Z, Yao L, Ding N, Jiang L, Wu Y. Bright light therapy in the treatment of patients with bipolar disorder: a systematic review and meta-analysis. *PLoS ONE*. 2020;15:e0232798.
- Webler FS, Spitschan M, Foster RG, Andersen M, Peirson SN. What is the 'spectral diet' of humans? *Curr Opin Behav Sci*. 2019;30:80–6.
- WHO. Hsuan resources data by country. 2019a. <https://apps.who.int/gho/data/node.main.MHHR?lang=en>. Accessed 3 Feb 2021.
- WHO. WHO Urbanization. 2019b <https://ourworldindata.org/urbanization>. Accessed 3 Feb 2021.
- Wild M. Enlightening global dimming and brightening. *Bull Am Meteor Soc*. 2012;93:27–37.
- Wirz-Justice A, Benedetti F. Perspectives in affective disorders: clocks and sleep. *Eur J Neurosci*. 2020;51:346–65.
- Wirz-Justice A, Skene DJ, Münch M. The relevance of daylight for humans. *Biochem Pharmacol*. 2020;28: 114304. <https://doi.org/10.1016/j.bcp.2020.114304>.
- Woo JM, Okusaga O, Postolache TT. Seasonality of suicidal behavior. *Int J Environ Res Public Health*. 2012;9:531–47.
- World Bank. Gini Index (World Bank estimate). 2020a. <https://data.worldbank.org/indicator/SI.POVGINI>. Accessed 3 Feb 2021.
- World Bank. Individuals using the Internet (% of population). 2020b. <https://data.worldbank.org/indicator/IT.NET.USER.ZS>. Accessed 3 Feb 2021.
- Wu A, Wang JY, Jia CX. Religion and completed suicide: a meta-analysis. *PLoS ONE*. 2015;10:e0131715.
- Young TK, Revich B, Soininen L. Suicide in circumpolar regions: an introduction and overview. *Int J Circumpolar Health*. 2015;74:27349.
- Yu B, Chen X. Age and birth cohort-adjusted rates of suicide mortality among us male and female youths aged 10 to 19 years from 1999 to 2017. *JAMA Netw Open*. 2019;2:e1911383.
- Yuodelis-Flores C, Ries RK. Addiction and suicide: a review. *Am J Addict*. 2015;24:98–104.
- Zeger SL, Liang KY. Longitudinal data analysis for discrete and continuous outcomes. *Biometrics*. 1986;42:121–30.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

---

# Variations in seasonal solar insolation are associated with a history of suicide attempts in bipolar I disorder

## ORIGINALITY REPORT

26%

SIMILARITY INDEX

21%

INTERNET SOURCES

21%

PUBLICATIONS

13%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** [findresearcher.sdu.dk:8443](https://findresearcher.sdu.dk/8443) Internet Source 1%
- 2** [docplayer.net](https://docplayer.net) Internet Source 1%
- 3** Submitted to American Public University System Student Paper 1%
- 4** [orca.cf.ac.uk](https://orca.cf.ac.uk) Internet Source 1%
- 5** Jörn Conell, Rita Bauer, Tasha Glenn, Martin Alda et al. "Online information seeking by patients with bipolar disorder: results from an international multisite survey", International Journal of Bipolar Disorders, 2016 Publication 1%
- 6** [www.mdpi.com](https://www.mdpi.com) Internet Source 1%
- 7** M. Bauer, T. Glenn, M. Alda, R. Bauer, P. Grof, W. Marsh, S. Monteith, R. Munoz, N. Rasgon, 1%

K. Sagduyu, P. C. Whybrow. "Trajectories of adherence to mood stabilizers in patients with bipolar disorder", International Journal of Bipolar Disorders, 2019

Publication

---

8	Submitted to Quinsigamond Community College Student Paper	1 %
9	pubmed.ncbi.nlm.nih.gov Internet Source	1 %
10	academic.oup.com Internet Source	1 %
11	link.springer.com Internet Source	1 %
12	Submitted to Tarleton State University Student Paper	1 %
13	Rita Bauer, Tasha Glenn, Scott Monteith, Peter C. Whybrow, Michael Bauer. "Survey of psychiatrist use of digital technology in clinical practice", International Journal of Bipolar Disorders, 2020 Publication	1 %
14	www.physiciansweekly.com Internet Source	1 %
15	Bauer, Rita, Jörn Conell, Tasha Glenn, Martin Alda, Raffaella Arda, Bernhard T. Baune,	<1 %

Michael Berk, Yuly Bersudsky, Amy Bilderbeck, Alberto Bocchetta, Letizia Bossini, Angela M. Paredes Castro, Eric YW. Cheung, Caterina Chillotti, Sabine Choppin, Maria Del Zompo, Rodrigo Dias, Seetal Dodd, Anne Duffy, Bruno Etain, Andrea Fagiolini, Miryam Fernández Hernandez, Julie Garnham, John Geddes, Jonas Gildebroy, Ana Gonzalez-Pinto, Guy M. Goodwin, Paul Grof, Hirohiko Harima, Stefanie Hassel, Chantal Henry, Diego Hidalgo-Mazzei, Vaisnvy Kapur, Girish Kunigiri, Beny Lafer, Erik R. Larsen, Ute Lewitzka, Rasmus W. Licht, Anne Hvenegaard Lund, Blazej Misiak, Scott Monteith, Rodrigo Munoz, Takako Nakanotani, René E Nielsen, Claire O'Donovan, Yasushi Okamura, Yamima Osher, Patryk Piotrowski, Andreas Reif, Philipp Ritter, Janusz K. Rybakowski, Kemal Sagduyu, Brett Sawchuk, Elon Schwartz, Ângela M. Scippa, Claire Slaney, Ahmad H. Sulaiman, Kirsi Suominen, Aleksandra Suwalska, Peter Tam, Yoshitaka Tatebayashi et al. "Internet use by patients with bipolar disorder: Results from an international multisite survey", *Psychiatry Research*, 2016.

Publication

17	<a href="https://preview-journalbipolardisorders.springeropen.com">preview-journalbipolardisorders.springeropen.com</a> Internet Source	<1 %
18	<a href="https://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	<1 %
19	Scott Monteith, Tasha Glenn, John Geddes, Emanuel Severus, Peter C. Whybrow, Michael Bauer. "Internet of things issues related to psychiatry", <i>International Journal of Bipolar Disorders</i> , 2021 Publication	<1 %
20	<a href="https://journals.sagepub.com">journals.sagepub.com</a> Internet Source	<1 %
21	"Bipolar Depression: Molecular Neurobiology, Clinical Diagnosis, and Pharmacotherapy", Springer Science and Business Media LLC, 2016 Publication	<1 %
22	Submitted to Aspen University Student Paper	<1 %
23	Submitted to Georgia Gwinnett College Student Paper	<1 %
24	<a href="https://hdl.handle.net">hdl.handle.net</a> Internet Source	<1 %
25	<a href="https://www.biorxiv.org">www.biorxiv.org</a> Internet Source	<1 %

26

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

&lt;1 %

27

[neurotree.org](http://neurotree.org)

Internet Source

&lt;1 %

28

[www.pubfacts.com](http://www.pubfacts.com)

Internet Source

&lt;1 %

29

Mirko Manchia, Eduard Vieta, Olav B. Smeland, Cara Altimus et al. "Translating big data to better treatment in bipolar disorder - a manifesto for coordinated action", *European Neuropsychopharmacology*, 2020

Publication

&lt;1 %

30

[brainstimulationclinic.squarespace.com](http://brainstimulationclinic.squarespace.com)

Internet Source

&lt;1 %

31

Understanding Suicide, 2016.

Publication

&lt;1 %

32

Bauer, Michael, Tasha Glenn, Martin Alda, Ole A Andreassen, Raffaella Arda, Frank Bellivier, Michael Berk, Thomas D Bjella, Letizia Bossini, Maria Del Zompo, Seetal Dodd, Andrea Fagiolini, Mark A Frye, Ana Gonzalez-Pinto, Chantal Henry, Flávio Kapczinski, Sebastian Kliwicki, Barbara Kłnig, Mauricio Kunz, Beny Lafer, Carlos Lopez-Jaramillo, Mirko Manchia, Wendy Marsh, Mónica Martínez-Cengotitabengoa, Ingrid Melle, Gunnar

&lt;1 %

Morken, Rodrigo Munoz, Fabiano G Nery, Claire Oâ€™Donovan, Andrea Pfennig, Danilo Quiroz, Natalie Rasgon, Andreas Reif, Janusz Rybakowski, Kemal Sagduyu, Christian Simhandl, Carla Torrent, Eduard Vieta, Mark Zetin, and Peter C Whybrow. "Impact of sunlight on the age of onset of bipolar disorder : Sunlight and age of onset", *Bipolar Disorders*, 2012.

Publication

---

33

[buscador.una.edu.ni](http://buscador.una.edu.ni)

Internet Source

<1 %

---

34

[journals.plos.org](http://journals.plos.org)

Internet Source

<1 %

---

35

Alexandra Schneider, Regina Hampel, Karl-Heinz Ladwig, Jens Baumert, Karoline Lukaschek, Annette Peters, Susanne Breitner. "Impact of meteorological parameters on suicide mortality rates: A case-crossover analysis in Southern Germany (1990–2006)", *Science of The Total Environment*, 2020

Publication

---

36

Pierre A. Geoffroy, Laura Palagini. "Biological rhythms and chronotherapeutics in depression", *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 2020

Publication

---

<1 %

37	Thomson, Daniel, Alyna Turner, Sue Lauder, Margaret E. Gigler, Lesley Berk, Ajeet B. Singh, Julie A. Pasco, Michael Berk, and Louisa Sylvia. "A brief review of exercise, bipolar disorder, and mechanistic pathways", <i>Frontiers in Psychology</i> , 2015. Publication	<1 %
38	<a href="http://www.nature.com">www.nature.com</a> Internet Source	<1 %
39	<a href="http://www.scientistsolutions.com">www.scientistsolutions.com</a> Internet Source	<1 %
40	<a href="http://expertscape.com">expertscape.com</a> Internet Source	<1 %
41	<a href="http://safetylit.org">safetylit.org</a> Internet Source	<1 %
42	<a href="http://www.studypool.com">www.studypool.com</a> Internet Source	<1 %
43	<a href="http://www.ywmedicalcentre.com">www.ywmedicalcentre.com</a> Internet Source	<1 %
44	<a href="http://digitalcommons.chapman.edu">digitalcommons.chapman.edu</a> Internet Source	<1 %
45	<a href="http://www.dmimsu.edu.in">www.dmimsu.edu.in</a> Internet Source	<1 %
46	Submitted to University of Queensland Student Paper	<1 %

47	<a href="http://www.siftdesk.org">www.siftdesk.org</a> Internet Source	<1 %
48	Submitted to Trinity College Dublin Student Paper	<1 %
49	<a href="http://www.medrxiv.org">www.medrxiv.org</a> Internet Source	<1 %
50	Submitted to Rutgers University, New Brunswick Student Paper	<1 %
51	<a href="http://www.kenes.com">www.kenes.com</a> Internet Source	<1 %
52	Submitted to Florida International University Student Paper	<1 %
53	Submitted to University of West Florida Student Paper	<1 %
54	<a href="http://beluga.sub.uni-hamburg.de">beluga.sub.uni-hamburg.de</a> Internet Source	<1 %
55	<a href="http://isbd2012.org">isbd2012.org</a> Internet Source	<1 %
56	Submitted to University of Hertfordshire Student Paper	<1 %
57	<a href="http://royalsocietypublishing.org">royalsocietypublishing.org</a> Internet Source	<1 %
58	<a href="http://www.allbusiness.com">www.allbusiness.com</a>	

Internet Source

<1 %

59

[www.aminer.org](http://www.aminer.org)

Internet Source

<1 %

60

[www.eapmverona2018.com](http://www.eapmverona2018.com)

Internet Source

<1 %

61

Ashlee B. Grierson, Ian B. Hickie, Sharon L. Naismith, Daniel F. Hermens, Elizabeth M. Scott, Jan Scott. "Circadian rhythmicity in emerging mood disorders: state or trait marker?", *International Journal of Bipolar Disorders*, 2016

Publication

<1 %

62

Mythily Subramaniam, Yunjue Zhang, Jue Hua Lau, Janhavi Ajit Vaingankar, Edimansyah Abdin, Siow Ann Chong, Eng Sing Lee. "Patterns of physical activity and health-related quality of life amongst patients with multimorbidity in a multi-ethnic Asian population", *Research Square*, 2019

Publication

<1 %

63

[doku.pub](http://doku.pub)

Internet Source

<1 %

64

[ebin.pub](http://ebin.pub)

Internet Source

<1 %

65

[tel.archives-ouvertes.fr](http://tel.archives-ouvertes.fr)

Internet Source

<1 %

66

[www.iesabroad.org](http://www.iesabroad.org)

Internet Source

<1 %

67

[www.igsli.org](http://www.igsli.org)

Internet Source

<1 %

68

[www.scribd.com](http://www.scribd.com)

Internet Source

<1 %

69

Céline Vetter, P. Morgan Pattison, Kevin Houser, Michael Herf et al. "A Review of Human Physiological Responses to Light: Implications for the Development of Integrative Lighting Solutions", LEUKOS, 2021

Publication

<1 %

70

Submitted to East Tennessee State University

Student Paper

<1 %

71

Jack Wood, Eugenia Wong, Ridwaan Joghee, Aya Balbaa et al. "Upregulation of expression occurs exclusively on microglial contact with plaques ", Cold Spring Harbor Laboratory, 2022

Publication

<1 %

72

Journal of Product & Brand Management, Volume 21, Issue 7 (2012-11-03)

Publication

<1 %

73

[bmcmmededuc.biomedcentral.com](http://bmcmmededuc.biomedcentral.com)

Internet Source

<1 %

74

[www.ijhpm.com](http://www.ijhpm.com)

Internet Source

<1 %

75

[www.nice.org.uk](http://www.nice.org.uk)

Internet Source

<1 %

76

Submitted to Creighton University

Student Paper

<1 %

77

Maria Faurholt-Jepsen, Darius Adam Rohani, Jonas Busk, Maj Vinberg, Jakob Eyvind Bardram, Lars Vedel Kessing. "Voice analyses using smartphone-based data in patients with bipolar disorder, unaffected relatives and healthy control individuals, and during different affective states", International Journal of Bipolar Disorders, 2021

Publication

<1 %

78

Submitted to Oral Roberts University

Student Paper

<1 %

79

Sandra J. Rosenthal, Travis Josephs, Oleg Kovtun, Richard McCarty. "Rate of change in solar insolation is a hidden variable that influences seasonal alterations in bipolar disorder", Brain and Behavior, 2021

Publication

<1 %

80

Submitted to University of Wolverhampton

Student Paper

<1 %

81

Yejin Lee, Dongbin Lee, Hyewon Jung, Yunji Cho, Ji Hyun Baek, Kyung Sue Hong. "The heterogeneous early illness courses of Korean patients with bipolar disorders – replication of the stage model", Research Square Platform LLC, 2022

Publication

<1 %

82

res.mdpi.com

Internet Source

<1 %

83

www.cambridge.org

Internet Source

<1 %

84

Emilio Sacchetti, Alessandro Galluzzo, Paolo Valsecchi. "Oral ziprasidone in the treatment of patients with bipolar disorders: a critical review", Expert Review of Clinical Pharmacology, 2014

Publication

<1 %

85

C. Christodoulou. "Suicide and seasonality : Seasonal variation of suicide", Acta Psychiatrica Scandinavica, 02/2012

Publication

<1 %

86

Cise Kizilirmak, Emanuele Monteleone, Jose M. Garcia-Manteiga, Francesca Brambilla et al. "Small transcriptional differences lead to

<1 %

distinct NF- $\kappa$ B dynamics in quasi-identical cells", Cold Spring Harbor Laboratory, 2021

Publication

---

87

Dick Schijven, Merel C. Postema, Masaki Fukunaga, Junya Matsumoto et al. "Large-scale analysis of structural brain asymmetries in schizophrenia via the ENIGMA consortium", Cold Spring Harbor Laboratory, 2022

Publication

---

<1 %

88

Max Lam, Chen Chia-Yen, Xia Yan, W. David Hill et al. "Identifying Nootropic Drug Targets via Large-Scale Cognitive GWAS and Transcriptomics", Cold Spring Harbor Laboratory, 2020

Publication

---

<1 %

89

Rita Bauer, Tasha Glenn, Sergio Strejilevich, Jörn Conell et al. "Internet use by older adults with bipolar disorder: international survey results", International Journal of Bipolar Disorders, 2018

Publication

---

<1 %

90

Tondo, L., M. Pompili, A. Forte, and R. J. Baldessarini. "Suicide attempts in bipolar disorders: comprehensive review of 101 reports", Acta Psychiatrica Scandinavica, 2015.

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      < 5 words

Exclude bibliography      On