

# RELATIONSHIP BETWEEN BODY MASS INDEX AND EXTRAPYRAMIDAL SYMPTOMS IN ASIAN PATIENTS WITH SCHIZOPHRENIA: THE RESEARCH ON ASIAN PSYCHOTROPIC

*by Andi Jayalangkara*

---

**Submission date:** 15-Mar-2022 08:22AM (UTC+0700)

**Submission ID:** 1784487968

**File name:** ENTS\_WITH\_SCHIZOPHRENIA-\_THE\_RESEARCH\_ON\_ASIAN\_PSYCHOTROPIC.pdf (141.04K)

**Word count:** 8214

**Character count:** 41028

## RELATIONSHIP BETWEEN BODY MASS INDEX AND EXTRAPYRAMIDAL SYMPTOMS IN ASIAN PATIENTS WITH SCHIZOPHRENIA: THE RESEARCH ON ASIAN PSYCHOTROPIC PRESCRIPTION PATTERNS FOR ANTIPSYCHOTICS (REAP-AP)

Seon-Cheol Park<sup>1</sup>, Adarsh Tripathi<sup>2</sup>, Ajit Avasthi<sup>3</sup>, Sandeep Grover<sup>3</sup>, Andi J. Tanra<sup>4</sup>, Takahiro A. Kato<sup>5</sup>, Toshiya Inada<sup>6</sup>, Kok Yoon Chee<sup>7</sup>, Mian-Yoon Chong<sup>8</sup>, Shu-Yu Yang<sup>9</sup>, Sih-Kun Lin<sup>10</sup>, Kang Sim<sup>11</sup>, Yu-Tao Xiang<sup>12</sup>, Afzal Javed<sup>13</sup>, Norman Sartorius<sup>14</sup>, Naotaka Shinfuku<sup>15</sup> & Yong Chon Park<sup>16</sup>

<sup>1</sup>Department of Psychiatry, Inje University Haeundae Paik Hospital, Busan, Republic of Korea

<sup>2</sup>Department of Psychiatry, King George's Medical University, Chowk, Lucknow, India

<sup>3</sup>Department of Psychiatry, Post Graduate Institute of Medical Education and Research (PGIMER), Chandigarh, India

<sup>4</sup>Department of Psychiatry, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

<sup>5</sup>Department of Neuropsychiatry, Graduate School of Medicine, Kyushu University, Fukuoka, Japan

<sup>6</sup>Department of Psychiatry and Psychobiology, Nagoya University, Graduate School of Medicine, Nagoya, Japan

<sup>7</sup>Tunku Abdul Rahman Institute of Neuroscience, Kuala Lumpur Hospital, Kuala Lumpur, Malaysia

<sup>8</sup>Chang Gung Memorial Hospital, Chiayi, and Chang Gung University School of Medicine, Taiwan

<sup>9</sup>Department of Pharmacy, Songde Branch, Tapei City Hospital, Tapei, Taiwan

<sup>10</sup>Psychiatric Center, Taipei City Hospital, Taipei, Taiwan

<sup>11</sup>West Region, Institute of Mental Health and Yong Loo Lin School of Medicine, National University of Singapore,

Singapore, Singapore

<sup>12</sup>Faculty of Health Sciences, Unit of Psychiatry, Institute of Translational Medicine, University of Macau,

Macau SAR, China

<sup>13</sup>Pakistan Psychiatric Research Centre, Fountain House, Lahore, Pakistan

<sup>14</sup>Association for the Improvement of Mental Health Programmes, Geneva, Switzerland

<sup>15</sup>Department of Social Welfare, School of Human Sciences, Seinan Gakuin University, Fukuoka, Japan

<sup>16</sup>Department of Neuropsychiatry, Hanyang University Guri Hospital, Guri, Republic of Korea

received: 15.2.2020;

revised: 5.3.2020;

accepted: 23.5.2020

### SUMMARY

**Background:** Although an inverse relationship between body mass index (BMI) and Parkinson disease (PD) has been repeatedly reported, our knowledge, the relationship between BMI and antipsychotic-induced extrapyramidal symptoms (EPS) has rarely been studied in patients with schizophrenia. Our study aimed to evaluate the relationship between BMI and EPS in patients with schizophrenia.

**Subjects and methods:** Using data from the Research on Asian Psychotropic Prescription Patterns for Antipsychotics (REAP-AP) study, we compared the prevalence of EPS in 1448 schizophrenia patients stratified as underweight, normal range, overweight pre-obese, overweight obese I, overweight obese II, and overweight obese III according to the World Health Organization (WHO) classification system for body weight status, and with underweight, normal range, overweight at risk, overweight obese I, and overweight obese II according to the Asia-Pacific obesity classification.

**Results:** In the first step of the WHO classification system for body weight status, adjusting for the potential effects of confounding factors, the multinomial logistic regression model revealed that underweight was significantly associated with greater rates of bradykinesia and muscle rigidity, and a lower rate of gait disturbance. In the second step of the Asia-Pacific obesity classification, adjusting for the potential effects of confounding factors, the multinomial logistic regression model revealed that underweight was significantly associated with a higher rate of muscle rigidity.

**Conclusion:** Findings of the present study consistently revealed that underweight was associated with a greater rate of muscle rigidity in a stepwise pattern among Asian patients with schizophrenia. Although the mechanism underlying the inverse relationship between BMI and muscle rigidity cannot be sufficiently explained, it is speculated that low BMI may contribute to the development of muscle rigidity regardless of antipsychotic "typicality" and dose in patients with schizophrenia.

**Key words:** antipsychotics - Asian, body mass index (BMI) - extrapyramidal symptoms (EPS) - muscle rigidity - schizophrenia

\* \* \* \* \*

### INTRODUCTION

An inverse relationship between body mass index (BMI) and Parkinson disease (PD) has been reported by previous studies, despite several controversies. Low BMI

has been regarded as a risk factor for PD regardless of the confounding factors including smoking, caffeine and alcohol (Logroscino et al. 2007, Noyce et al. 2017). 5-hydroxytryptamine-1A (5-HT<sub>1A</sub>) receptor agonist can reduce the motor deficits in PD as well as antipsychotic-

induced extrapyramidal symptoms (EPS) in schizophrenia because serotonergic input from the dorsal raphe nucleus can modulate dopaminergic neurons in the substantia nigra. Hence, 5-HT<sub>1A</sub> receptor agonist has been regarded as a promising therapeutic agent for PD and schizophrenia (Haleem 2015, Ohno 2011). Also, from the viewpoint of neuro-inflammation, the microglial activation associated with both PD and EPS can be a cause for prostaglandin release (Arora et al. 2018). Furthermore, because the facilitatory role of dopamine D<sub>2</sub> receptor agonists on movement can be potentiated by blocking the cannabinoid CB<sub>1</sub> receptor, the endogenous cannabinoid system has been suggested as a novel therapeutic target for dopamine-related diseases including PD and psychosis (De Fonseca et al. 2001). It can, therefore, be speculated that common biological underpinnings may be shared by both core parkinsonian symptoms in PD and antipsychotic-induced EPS in schizophrenia. However, to our knowledge, the relationship between BMI and antipsychotic-induced EPS has been rarely studied. Therefore, using data from the Research on Asian Psychotropic Prescription Patterns for Antipsychotics (REAP-AP) study, the largest international collaborative psychiatric survey in Asia (Park et al. 2018), we aimed to clarify the relationship between BMI and EPS in Asian patients with schizophrenia. Among Asian patients with schizophrenia, BMI groups were categorized according to the World Health Organization (WHO) classification system for body weight status (World Health Organization 1998) and the Asia-Pacific obesity classification (International Obesity Task Force 2000). Additionally, we aimed to compare findings regarding the relationship between BMI and EPS according to the different obesity classification systems.

## SUBJECTS AND METHODS

As described elsewhere (Park et al. 2018), in the fourth REAP-AP, a consecutive recruitment of 3,744 patients with schizophrenia was conducted from 71 survey centers in 15 Asian countries including Bangladesh, China, Hong Kong, India, Indonesia, Japan, Korea, Malaysia, Myanmar, Pakistan, Singapore, Sri Lanka, Taiwan, Thailand and Vietnam from March to June 2016. The institutional review boards of Taipei City Hospital, Taipei, Taiwan (receipt number: TCHIRB-10412128-E) and other survey centers approved the study protocols and informed consent forms. All study subjects signed the informed consent form prior to their participation in the study. Using the predefined case report form, the baseline characteristics and pharmacotherapy-related details were collected by the study coordinators, under the supervision of clinical psychiatrists. In our study, the analysis of data from the REAP-AP study was performed, and the subjects who met the following inclusion criteria were selected: (i) a diagnosis of schizo-

phrenia according to the *International Classification of Diseases, 10<sup>th</sup> Revision* (ICD-10) (World Health Organization 1992), (ii) pharmacotherapy with antipsychotics, and (iii) availability of the complete Drug-induced Extrapyramidal Symptom Scale (DIEPSS) data (Inada 2009, Kim et al. 2002). Finally, we included 1448 schizophrenia patients recruited from India, Indonesia, Japan, Taiwan and Malaysia.

In the first step, BMI, defined as weight in kilograms (kg) divided by height in meters squared (m<sup>2</sup>), was used to categorize schizophrenia patients into 6 groups according to the WHO classification system for body weight status (World Health Organization 1998) as follows: underweight (<18.50 kg/m<sup>2</sup>), normal range (18.50–24.99 kg/m<sup>2</sup>), overweight (pre-obese (25.00–29.99 kg/m<sup>2</sup>)), overweight obese I (30.00–34.99 kg/m<sup>2</sup>), overweight obese II (35.00–39.99 kg/m<sup>2</sup>), and overweight obese III (>40.00 kg/m<sup>2</sup>). However, compared with Europeans, Asians who exhibit risks associated with obesity are increased in those with lower BMIs, and Pacific Islanders exhibit lower body fat levels for the same BMI. Therefore, in the second step, schizophrenia patients were categorized into 5 groups according to the Asia-Pacific obesity classification (International Obesity Task Force 2000) as follows: underweight (<18.50 kg/m<sup>2</sup>); normal range (18.50–22.99 kg/m<sup>2</sup>); overweight at risk (23.00–24.99 kg/m<sup>2</sup>); overweight obese I (25.00–29.9 kg/m<sup>2</sup>); and overweight obese II (>30.0 kg/m<sup>2</sup>).

The DIEPSS was used to evaluate the antipsychotic-induced EPS among the study subjects. All the DIEPSS items, which are gait, bradykinesia, sialorrhea, muscle rigidity, tremor, akathisia, dystonia, dyskinesia and overall severity, were rated on a 5-point Likert scale with score 1 (normal) to 4 (severe). The inter-rater reliability, test-retest reliability and concurrent validity were assessed. According to the confirmed metric characteristics, the presence of gait, bradykinesia, sialorrhea, muscle rigidity, tremor, akathisia, dystonia, dyskinesia and overall severity was indicated by a score of 2 or higher in each of the items and the presence of parkinsonism was indicated by a total sum score of 5 or higher in 5 items including gait, bradykinesia, sialorrhea, muscle rigidity and tremor (Inada 2009, Kim et al. 2002).

The Brief Psychiatric Rating Scale (BPRS) (Leucht et al. 2005, Overall & Gorham 1962), the Charlson Comorbidity Index (CCI) (Charlson et al. 1987, Quan et al. 2005) and the Anatomical Therapeutic Chemical (ATC) classification (World Health Organization 2019) were used to evaluate the psychiatric symptoms, quantify the comorbid physical disease and classify the used psychotropic drugs among the study subjects, respectively. The BPRS items - somatic concern, anxiety, emotional withdrawal, conceptual disorganization, feelings of guilt, tension, mannerism and posturing, grandiosity, depressed mood, hostility, suspiciousness, hallucinatory behavior, motor retardation, uncooperativeness, unusual thought

content, blurred affect, excitement and disorientation - were rated on a 7-point Likert scale from scores 1 (not present) to 7 (very severe) (Leucht et al. 2005, Overall & Gorham 1962). The CCI evaluated the comorbid physical diseases based on their weight and was regarded as a predictive factor for long-term outcome. Also, in our study, the CCI level was classified into 3 groups according to 1- and 2-year survival rates: 0-1, 2-3 and 4-5 (Charlson et al. 1987, Quan et al. 2005). Both the BPRS and the CCI were validated and confirmed by the study findings (Charlson et al. 1987, Leucht et al. 2005, Overall & Gorham 1962, Quan et al. 2005). In addition, based on the ATC classification system (World Health Organization 2019), psychotropic drugs were classified into antipsychotics (N05A), anticonvulsants (antiepileptics; N03A), antidepressants (N06A), anxiolytics (N05B and N05C) and antiparkinsonian drugs (N04). According to the definition of Tandon and coworkers (2010), antipsychotic medications were dichotomized into first- or second-generation antipsychotics in an association with "atypicality", which is defined as a favorable antipsychotic effect with a reduced risk of EPS.

In the first step for the WHO obesity classification, using analysis of covariance (ANCOVA) for continuous variables and the chi-squared ( $\chi^2$ ) test for discrete variables, baseline characteristics and psychotic-induced EPS in schizophrenia patients with underweight, normal range, pre-obese, obese I, obese II, and obese III were compared. Multinomial logistic regression analyses were used to adjust for the potential effect of confounding factors in terms of the differences in the proportions of antipsychotic-induced EPS according to the WHO obesity classification. In the second step for the Asia-Pacific obesity classification, using ANCOVA for continuous variables and the  $\chi^2$  test for discrete variables, baseline characteristics and psychotic-induced EPS in schizophrenia patients with underweight, normal range, at risk, obese I, and obese II were compared. Multinomial logistic regression analyses were used to adjust for the potential effect of confounding factors in terms of the differences in the proportions of antipsychotic-induced EPS according to the Asia-Pacific Obesity classification. Statistical significance was set at  $P < 0.05$  (two-tailed) for all the tests. All statistical analyses were conducted using IBM SPSS 24 (IBM Co., Armonk, NY, USA).

## RESULTS

As shown in Table 1, among 1448 Asian patients with schizophrenia, the percentages of underweight, normal range, pre-obese, obese I, obese II, and obese III were 8.4% (n=122), 56.1% (n=812), 25.3% (n=366), 7.4% (n=107), 1.7% (n=25), and 1.1% (n=16), respectively. The percentage representation of the country ( $\chi^2=60.692$ ,  $P < 0.0001$ ), regional classification ( $\chi^2=21.989$ ,  $P=0.015$ ), income group ( $\chi^2=32.548$ ,  $P < 0.0001$ ), enrollment as an outpatient ( $\chi^2=41.957$ ,  $P < 0.0001$ ), duration

of illness ( $\chi^2=60.630$ ,  $P < 0.0001$ ), duration of untreated psychosis ( $\chi^2=20.028$ ,  $P=0.029$ ), antidepressants ( $\chi^2=21.864$ ,  $P < 0.0001$ ), age ( $F=1767.2$ ,  $P < 0.0001$ ) and BPRS total score ( $F=1325.9$ ,  $P < 0.0001$ ) were significantly different among schizophrenia patients with normal range, pre-obese, obese I, obese II, and obese III in terms of the WHO classification system for body weight status. Multinomial logistic regression analyses were used to adjust for the potential effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, duration of untreated psychosis, antidepressants and BPRS total score. Additionally, underweight was defined as the reference category of the BMI classification in the multinomial logistic regression analyses. After adjusting for the effect of confounding factors, the presence of gait disturbance (normal range, adjusted odds ratio aOR=0.626,  $P=0.248$ ; pre-obese, aOR=0.644,  $P=0.335$ ; obese I, aOR=0.395,  $P=0.151$ ; obese II, aOR=3.868,  $P=0.041$ ; obese III, aOR=0.770,  $P=0.821$ ), bradykinesia (normal range, aOR=0.518,  $P=0.035$ ; pre-obese, aOR=0.246,  $P=0.671$ ; obese I, aOR=0.387,  $P=0.054$ ; obese II, aOR=1.451,  $P=0.544$ ; obese III, aOR=1.704,  $P=0.474$ ), and muscle rigidity (normal range, aOR=0.432,  $P=0.004$ ; pre-obese, aOR=0.717,  $P=0.308$ ; obese I, aOR=0.333,  $P=0.031$ ; obese II, aOR=0.596,  $P=0.523$ ; obese III, aOR=0.344,  $P=0.333$ ) were significantly different among the divided patient groups based on the WHO classification system for body weight status.

As shown in Table 2, the percentages of underweight, normal range, at risk, obese I, and obese II were 8.4% (n=122), 37.1% (n=537), 19.0% (n=275), 25.3% (n=366), and 10.2% (n=148), respectively. The percentage representation of the country ( $\chi^2=62.437$ ,  $P < 0.0001$ ), regional classification ( $\chi^2=23.577$ ,  $P=0.003$ ), income group ( $\chi^2=32.841$ ,  $P < 0.0001$ ), enrollment as an outpatient ( $\chi^2=42.164$ ,  $P < 0.0001$ ), duration of illness ( $\chi^2=54.674$ ,  $P < 0.0001$ ), antidepressants ( $\chi^2=21.484$ ,  $P < 0.0001$ ), and age ( $F=3557.8$ ,  $P < 0.0001$ ) were significantly different among schizophrenia patients with underweight, normal range, at risk, obese I, and obese II in terms of the Asia-Pacific obesity classification. Multinomial logistic regression analyses were used to adjust for the potential effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, and antidepressants. Additionally, underweight was defined as the reference category for the BMI classification in multinomial logistic regression analyses. After adjusting for the effect of confounding factors, the presence of muscle rigidity (normal range, aOR=0.569,  $P=0.035$ ; at risk, aOR=0.387,  $P=0.003$ ; obese I, aOR=0.241,  $P=0.715$ ; obese II, aOR=0.430,  $P=0.034$ ) and akathisia (normal range, aOR=0.641,  $P=0.176$ ; at risk, aOR=0.440,  $P=0.034$ ; obese I, aOR=0.643,  $P=0.214$ ; obese II, aOR=0.436,  $P=0.094$ ) were significantly different among the divided patient groups based on the Asia-Pacific obesity classification system.

**Table 1.** Antipsychotic-induced extrapyramidal symptoms divided into groups according to the World Health Organization (WHO) classification system for body weight status in Asian patients with schizophrenia (n=1448)

	Underweight (n=122)		Normal range (n=812)		Overweight: Pre-obese (n=366)		Overweight: Obese I (n=107)		Overweight: Obese II (n=25)		Overweight: Obese III (n=16)		Statistical coefficient	Unadjusted P-value	Adjusted P-value*	
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)				
Age, mean (SD) years	35.4 (13.7)		38.4 (12.9)		39.2 (11.5)		41.5 (12.1)		38.5 (11.3)		36.9 (7.1)		F=1767.2	<0.0001	-	
Male, n (%)	69 (56.6)		504 (62.1)		234 (63.9)		53 (49.5)		13 (52.0)		10 (62.5)		$\chi^2=9.506$	0.091	-	
Country													$\chi^2=60.692$	<0.0001	-	
India, n (%)	36 (29.5)		226 (27.8)		125 (34.2)		31 (29.0)		5 (20.0)		3 (18.8)					
Indonesia, n (%)	52 (42.6)		293 (36.1)		85 (23.2)		18 (16.8)		9 (36.0)		6 (37.5)					
Japan, n (%)	7 (5.7)		83 (10.2)		41 (11.2)		10 (9.3)		4 (16.0)		0 (0.0)					
Malaysia, n (%)	21 (17.2)		105 (12.9)		67 (18.3)		29 (27.1)		5 (20.0)		3 (18.0)					
Taiwan, n (%)	6 (4.9)		105 (12.9)		48 (13.1)		19 (17.8)		2 (0.1)		4 (0.3)					
Regional classification**													$\chi^2=21.989$	0.015	-	
East Asian, n (%)	13 (10.7)		188 (23.2)		89 (24.3)		29 (27.1)		6 (24.0)		4 (25.0)					
Southeast Asian, n (%)	73 (59.8)		398 (49.0)		152 (41.5)		47 (43.9)		14 (56.0)		9 (56.3)					
South Asian, n (%)	36 (29.5)		226 (27.8)		125 (34.2)		31 (29.0)		5 (20.0)		3 (18.8)					
Income group***													$\chi^2=32.548$	<0.0001	-	
High, n (%)	13 (10.7)		188 (23.2)		89 (24.3)		29 (27.1)		6 (24.0)		4 (25.0)					
Upper middle, n (%)	21 (17.2)		105 (12.9)		67 (18.3)		29 (27.1)		5 (20.0)		3 (18.8)					
Lower middle, n (%)	88 (72.1)		519 (63.9)		210 (57.4)		49 (45.8)		14 (56.0)		9 (56.3)					
Employment, n (%)	19 (15.6)		155 (19.1)		79 (21.6)		22 (20.6)		1 (4.0)		3 (18.8)		$\chi^2=6.214$	0.286	-	
Outpatient, n (%)	58 (47.5)		340 (51.7)		238 (65.0)		78 (72.9)		21 (84.0)		10 (62.5)		$\chi^2=41.957$	<0.0001	-	
Duration of illness													$\chi^2=60.630$	<0.0001	-	
<1 year, n (%)	26 (21.3)		158 (19.3)		33 (9.0)		8 (7.5)		2 (8.0)		0 (0.0)					
1-5 years, n (%)	33 (27.0)		177 (21.8)		79 (21.6)		20 (18.7)		6 (24.0)		2 (12.5)					
5-10 years, n (%)	28 (23.0)		136 (16.7)		65 (17.8)		12 (11.2)		7 (28.0)		2 (12.5)					
>10 years, n (%)	35 (28.7)		341 (42.0)		189 (51.6)		67 (62.6)		10 (40.0)		12 (75.0)					
DUP													$\chi^2=20.028$	0.029	-	
<3 months, n (%)	43 (35.2)		318 (39.2)		132 (36.1)		44 (41.1)		16 (64.0)		6 (37.5)					
3-12 months, n (%)	51 (41.8)		232 (28.6)		124 (33.9)		37 (34.6)		4 (16.0)		5 (31.3)					
>1 year, n (%)	28 (23.0)		262 (32.3)		110 (30.1)		26 (24.3)		5 (20.0)		5 (31.3)					

According to the WHO classification system for body weight status, the body mass index was classified into the underweight (<18.50), normal weight (18.50-24.99), overweight: pre-obese (25.00-29.99), overweight: obese I (30.00-34.99), overweight: obese II (35.00-39.99), and overweight: obese III (>40.00).  
 \* Adjusted for the effects of age, regional classification, income group, enrollment in the multinomial logistic regression analyses.  
 \*\* Defined by United Nations classification: East Asia (Japan and Taiwan), South Asia (India) and Southeast Asia (Indonesia and Malaysia).  
 \*\*\* Defined by the World Bank list of economies: high income (Japan and Taiwan), upper middle income (Malaysia) and lower middle income (India and Indonesia).  
 BMI - body mass index; BPRS - Brief Psychiatric Rating Scale; CCI - Charlson comorbidity index; DUP - duration of untreated psychosis; FGA - first-generation antipsychotics; SGA - second-generation antipsychotics; WHO - World Health Organization; a - normal range; b - pre-obese; c - obese I; d - obese II; e - obese III.

Table 1. - Continues 39

	Underweight (n=122)	Normal range (n=812)	Overweight: Pre-obese (n=366)	Overweight: Obese I (n=107)	Overweight: Obese II (n=25)	Overweight: Obese III (n=16)	Statistical coefficient	Unadjusted P-value	Adjusted P-value*
Cigarette smoking							$\chi^2=10.847$	0.370	-
Current, n (%)	29 (24.8)	260 (33.1)	95 (27.1)	28 (26.7)	6 (25.0)	4 (25.0)			
Previous, n (%)	12 (10.3)	59 (7.5)	33 (9.4)	8 (7.6)	3 (12.5)	3 (18.8)			
No, n (%)	76 (65.0)	466 (59.4)	222 (63.4)	69 (65.7)	15 (62.5)	9 (56.3)			
CCI level							$\chi^2=16.645$	0.341	-
0-1, n (%)	101 (82.8)	627 (77.2)	275 (75.1)	73 (68.2)	16 (64.0)	15 (93.8)			
2-3, n (%)	19 (15.6)	162 (20.0)	85 (23.2)	29 (27.1)	8 (32.0)	1 (6.3)			
4-5, n (%)	2 (1.6)	23 (2.8)	6 (1.6)	5 (4.7)	1 (4.0)	0 (0.0)			
BPRS, mean (SD)	35.9 (13.9)	35.4 (13.3)	34.0 (12.6)	31.8 (11.8)	35.4 (13.3)	38.3 (14.1)	F=1325.9	<0.0001	-
Psychotropic drugs							$\chi^2=5.259$	0.385	-
FGA, n (%)	41 (33.6)	283 (34.9)	133 (36.3)	35 (32.7)	13 (52.0)	8 (50.0)			
SGA, n (%)	104 (85.2)	677 (83.4)	307 (83.9)	94 (87.9)	19 (76.0)	11 (68.8)	$\chi^2=5.345$	0.375	-
Antiparkinson, n (%)	49 (40.2)	308 (37.9)	124 (33.9)	48 (44.9)	11 (44.0)	2 (12.5)	$\chi^2=9.675$	0.085	-
Anticonvulsants, n (%)	9 (7.4)	72 (8.9)	35 (9.6)	12 (11.2)	2 (8.0)	2 (12.5)	$\chi^2=1.422$	0.922	-
Anxiolytics, n (%)	41 (33.6)	264 (32.5)	93 (25.4)	33 (30.8)	6 (24.0)	4 (25.0)	$\chi^2=7.317$	0.198	-
Antidepressants, n (%)	9 (7.4)	53 (6.5)	53 (14.5)	14 (13.1)	2 (8.0)	2 (12.5)	$\chi^2=21.864$	<0.0001	-
Gait, n (%)	9 (7.8)	66 (8.1)	28 (7.7)	5 (4.7)	6 (24.0)	2 (12.5)	$\chi^2=10.874$	0.054	0.254 <sup>a</sup> 0.331 <sup>b</sup> 0.151 <sup>c</sup> 0.041 <sup>d</sup> 0.820 <sup>e</sup>
Bradykinesia, n (%)	18 (14.8)	89 (11.0)	46 (12.6)	8 (7.5)	5 (20.0)	3 (18.8)	$\chi^2=6.094$	0.297	0.036 <sup>a</sup> 0.242 <sup>b</sup> 0.054 <sup>c</sup> 0.541 <sup>d</sup> 0.472 <sup>e</sup>
Sialorrhea, n (%)	9 (7.4)	67 (8.3)	27 (7.4)	8 (7.5)	3 (12.0)	2 (12.5)	$\chi^2=1.349$	0.930	0.482 <sup>a</sup> 0.665 <sup>b</sup> 0.639 <sup>c</sup> 0.241 <sup>d</sup> 0.958 <sup>e</sup>

According to the WHO classification system for body weight status, the body mass index was classified into the underweight (<18.50), normal weight (18.50-24.99),

overweight, pre-obese (25.00-29.99), overweight-obese I (30.00-34.99), overweight-obese II (35.00-39.99), and overweight-obese III (>40.00).

\* Adjusted for the effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, DUP, total score on the BPRS, and antidepressants.

Underweight was defined as the reference category of the BMI classification in the multinomial logistic regression analyses.

\*\* Defined by United Nations classification: East Asia (Japan and Taiwan), South Asia (India) and Southeast Asia (Indonesia and Malaysia).

\*\*\* Defined by the World Bank list of economies: high income (Japan and Taiwan), upper middle income (Malaysia) and lower middle income (India and Indonesia).

BMI - body mass index; BPRS - Brief Psychiatric Rating Scale; CCI - Charlson comorbidity index; DUP - duration of untreated psychosis; FGA - first-generation antipsychotics;

SGA - second-generation antipsychotics; WHO - World Health Organization; a - normal range; b - pre-obese; c - obese I; d - obese II; e - obese III.

Table 1. Continues

	Underweight (n=122)	Normal range (n=812)	Overweight: Pre-obese (n=366)	Overweight: Obese I (n=107)	Overweight: Obese II (n=25)	Overweight: Obese III (n=16)	Statistical coefficient	Unadjusted P-value	Adjusted P-value*
Rigidity, n (%)	25 (20.5)	88 (10.8)	47 (27.5)	7 (6.4)	3 (12.0)	1 (6.3)	$\chi^2=13.628$	0.021	0.005 <sup>a</sup> 0.251 <sup>b</sup> 0.032 <sup>c</sup> 0.537 <sup>d</sup> 0.347 <sup>e</sup>
Tremor, n (%)	20 (16.4)	136 (16.7)	72 (19.7)	16 (15.0)	4 (16.0)	3 (18.8)	$\chi^2=2.142$	0.829	0.336 <sup>a</sup> 0.048 <sup>b</sup> 0.433 <sup>c</sup> 0.345 <sup>d</sup> 0.829 <sup>e</sup>
Akathisia, n (%)	15 (12.3)	59 (7.3)	28 (7.7)	5 (4.7)	2 (8.0)	0 (0.0)	$\chi^2=6.634$	0.249	0.156 <sup>a</sup> 0.664 <sup>b</sup> 0.120 <sup>c</sup> 0.776 <sup>d</sup> 0.999 <sup>e</sup>
Dystonia, n (%)	9 (7.4)	37 (4.6)	20 (5.5)	1 (1.5)	1 (4.0)	0 (0.0)	$\chi^2=6.675$	0.246	0.087 <sup>a</sup> 0.361 <sup>b</sup> 0.996 <sup>c</sup> 0.869 <sup>d</sup> 0.999 <sup>e</sup>
Dyskinesia, n (%)	4 (3.3)	43 (5.3)	14 (3.8)	3 (2.8)	2 (8.0)	0 (0.0)	$\chi^2=4.128$	0.531	0.323 <sup>a</sup> 0.896 <sup>b</sup> 0.976 <sup>c</sup> 0.219 <sup>d</sup> 0.999 <sup>e</sup>
Overall, n (%)	23 (18.9)	134 (16.5)	63 (17.2)	15 (14.0)	5 (20.0)	2 (12.5)	$\chi^2=1.449$	0.919	0.554 <sup>a</sup> 0.783 <sup>b</sup> 0.518 <sup>c</sup> 0.598 <sup>d</sup> 0.915 <sup>e</sup>
Parkinsonism, n (%)	25 (34.7)	131 (31.7)	68 (34.3)	16 (33.3)	5 (33.3)	3 (37.5)	$\chi^2=0.638$	0.986	0.411 <sup>a</sup> 0.730 <sup>b</sup> 0.620 <sup>c</sup> 0.556 <sup>d</sup> 0.989 <sup>e</sup>

According to the WHO classification system for body weight status, the body mass index was classified into the underweight ( $\leq 18.50$ ), normal weight (18.50-24.99), overweight: pre-obese (25.00-29.99), overweight: obese I (30.00-34.99), overweight: obese II (35.00-39.99), and overweight: obese III ( $>40.00$ ).  
 \* Adjusted for the effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, DUP, total score on the BPRS, and antidepressants.  
 Underweight was defined as the reference category of the BMI classification in the multinomial logistic regression analyses.  
 \*\* Defined by United Nations classification: East Asia (Japan and Taiwan), South Asia (India) and Southeast Asia (Indonesia and Malaysia).  
 \*\*\* Defined by the World Bank list of economies: high income (Japan and Taiwan), upper middle income (Malaysia) and lower middle income (India and Indonesia).  
 BMI - body mass index; BPRS - Brief Psychiatric Rating Scale; CCI - Charlson comorbidity index; DUP - duration of untreated psychosis; FGA - first-generation antipsychotics; SGA - second-generation antipsychotics; WHO - World Health Organization; a - normal range; b - pre-obese; c - obese I; d - obese II; e - obese III

**Table 2.** Antipsychotic-induced extrapyramidal symptoms divided into groups according to the Asia-Pacific obesity classification in Asian patients with schizophrenia (n=1448)

	Underweight (n=122)	Normal range (n=537)	Overweight: At risk (n=275)	Overweight: Obese I (n=366)	Overweight: Obese II (n=148)	Statistical coefficient	Unadjusted P-value	Adjusted P-value*
Age, mean (SD) years	35.4 (13.7)	38.1 (13.2)	39.0 (12.4)	39.2 (11.5)	40.5 (11.6)	F=3557.8	<0.0001	-
Male, n (%)	69 (56.6)	333 (62.0)	171 (62.2)	234 (63.9)	76 (51.4)	$\chi^2=8.519$	0.074	-
Country						$\chi^2=62.437$	<0.0001	-
India, n (%)	36 (29.5)	138 (25.7)	88 (32.0)	125 (34.2)	39 (26.4)			
Indonesia, n (%)	52 (42.6)	207 (38.5)	86 (31.3)	85 (23.2)	33 (22.3)			
Japan, n (%)	7 (5.7)	64 (11.9)	19 (6.9)	41 (11.2)	14 (9.5)			
Malaysia, n (%)	21 (17.2)	63 (11.7)	42 (15.3)	67 (18.3)	37 (25.0)			
Taiwan, n (%)	6 (4.9)	65 (12.1)	40 (14.5)	48 (13.1)	25 (16.9)			
Regional classification**						$\chi^2=23.577$	0.003	-
East Asian, n (%)	13 (10.7)	129 (24.0)	59 (21.5)	89 (24.3)	39 (26.4)			
Southeast Asian, n (%)	73 (59.8)	270 (50.3)	128 (46.5)	152 (41.5)	70 (47.3)			
South Asian, n (%)	36 (29.5)	138 (25.7)	88 (32.0)	125 (34.2)	39 (26.4)			
Income group***						$\chi^2=32.841$	<0.0001	-
High, n (%)	13 (10.7)	129 (24.0)	59 (21.5)	89 (24.3)	39 (26.4)			
Upper middle, n (%)	21 (17.2)	63 (11.7)	42 (15.3)	67 (18.3)	37 (25.0)			
Lower middle, n (%)	88 (72.1)	345 (64.2)	174 (63.3)	210 (48.6)	72 (48.6)			
Employment, n (%)	19 (15.6)	100 (18.6)	55 (20.0)	79 (21.6)	26 (17.6)	$\chi^2=2.847$	0.584	-
Outpatient, n (%)	58 (47.5)	268 (49.9)	152 (55.3)	238 (65.0)	109 (73.6)	$\chi^2=42.164$	<0.0001	-
Duration of illness						$\chi^2=54.674$	<0.0001	-
< 1 year, n (%)	26 (21.3)	108 (20.1)	50 (18.2)	33 (9.0)	10 (6.8)			
1-5 years, n (%)	33 (27.0)	121 (22.5)	56 (20.4)	79 (21.6)	28 (18.9)			
5-10 years, n (%)	28 (23.0)	85 (15.8)	51 (18.5)	65 (17.8)	21 (14.2)			
> 10 years, n (%)	35 (28.7)	223 (41.5)	118 (42.9)	189 (51.6)	89 (60.1)			
DUP						$\chi^2=14.492$	0.070	-
< 3 months, n (%)	43 (35.2)	210 (39.0)	108 (39.3)	132 (36.1)	66 (44.6)			
3-12 months, n (%)	51 (41.8)	152 (28.3)	80 (29.1)	124 (33.9)	46 (31.1)			
> 1 year, n (%)	28 (23.0)	175 (32.6)	87 (31.6)	110 (30.1)	36 (24.3)			

According to the WHO classification system for body weight status, the body mass index was classified into the underweight (<18.50), normal weight (18.50-24.99), overweight: pre-obese (25.00-29.99), overweight: obese I (30.00-34.99), overweight: obese II (35.00-39.99), and overweight: obese III (>40.00).  
 \* Adjusted for the effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, DUP, total score on the BPRS, and antidepressants.  
 \*\* Defined by United Nations classification: East Asia (Japan and Taiwan), South Asia (India) and Southeast Asia (Indonesia and Malaysia).  
 \*\*\* Defined by the World Bank list of economies: high income (Japan and Taiwan), upper middle income (Malaysia) and lower middle income (India and Indonesia).  
 BMI - body mass index; BPRS - Brief Psychiatric Rating Scale; CCI - Charlson comorbidity index; DUP - duration of untreated psychosis; FGA - first-generation antipsychotics; SGA - second-generation antipsychotics; WHO - World Health Organization; a - normal range; b - at risk; c - obese I; d - obese II.

Table 2. Continues

	Underweight (n=122)	Normal range (n=537)	Overweight: At risk (n=275)	Overweight: Obese I (n=366)	Overweight: Obese II (n=148)	Statistical coefficient	Unadjusted P-value	Adjusted P-value*
Cigarette smoking						$\chi^2=15.014$	0.059	-
Current, n (%)	29 (24.8)	186 (35.9)	74 (27.7)	95 (27.1)	38 (26.2)			
Previous, n (%)	12 (10.3)	41 (7.9)	18 (6.7)	33 (9.4)	14 (9.7)			
No, n (%)	76 (65.0)	291 (56.2)	175 (65.5)	222 (63.4)	93 (64.1)			
CCI level						$\chi^2=18.255$	0.108	-
0-1, n (%)	101 (82.8)	422 (78.6)	205 (74.5)	275 (75.1)	104 (70.3)			
2-3, n (%)	19 (15.6)	96 (17.9)	66 (24.0)	85 (23.2)	38 (25.7)			
4-5, n (%)	2 (1.6)	19 (3.6)	4 (1.5)	6 (1.6)	6 (4.1)			
BPRS, mean (SD)	35.9 (1.4)	35.9 (0.7)	34.5 (1.0)	34.0 (0.8)	33.0 (1.2)	F=1.635	0.163	-
Psychotropic drugs								
FGA, n (%)	41 (33.6)	186 (34.5)	98 (35.6)	133 (36.3)	56 (37.8)	$\chi^2=0.915$	0.922	0.942 <sup>a</sup>
SGA, n (%)	104 (85.2)	446 (83.1)	231 (84.0)	307 (83.9)	124 (83.8)	$\chi^2=0.406$	0.982	0.923 <sup>b</sup>
Antiparkinson, n (%)	49 (40.2)	203 (37.8)	105 (38.2)	124 (33.9)	61 (41.2)	$\chi^2=3.363$	0.499	0.657 <sup>c</sup>
Anticonvulsants, n (%)	9 (7.4)	47 (8.8)	25 (9.1)	35 (9.6)	16 (10.8)	$\chi^2=1.133$	0.889	0.449 <sup>d</sup>
Anxiolytics, n (%)	41 (33.6)	171 (31.8)	93 (33.8)	93 (25.4)	43 (29.1)	$\chi^2=7.065$	0.132	0.335 <sup>a</sup>
Antidepressants, n (%)	9 (7.4)	37 (6.9)	16 (5.8)	53 (14.5)	18 (12.2)	$\chi^2=21.484$	<0.0001	0.118 <sup>b</sup>
Gait, n (%)	41 (33.6)	171 (31.8)	93 (33.8)	93 (25.4)	43 (29.1)	$\chi^2=0.403$	0.982	0.801 <sup>c</sup>
Bradykinesia, n (%)	18 (14.8)	64 (11.9)	25 (9.1)	46 (12.6)	16 (10.8)	$\chi^2=3.325$	0.505	0.551 <sup>d</sup>
Sialorrhea, n (%)	9 (7.4)	48 (8.9)	19 (6.9)	27 (7.4)	13 (8.8)	$\chi^2=1.466$	0.833	0.507 <sup>a</sup>
								0.914 <sup>b</sup>
								0.969 <sup>c</sup>
								0.735 <sup>d</sup>

According to the WHO classification system for body weight status, the body mass index was classified into the underweight (<18.50), normal weight (18.50-24.99), overweight, pre-obese (25.00-29.99), overweight-obese I (30.00-34.99), overweight-obese II (35.00-39.99), and overweight-obese III (>40.00).  
 \* Adjusted for the effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, DUP, total score on the BPRS, and antidepressants.  
 Underweight was defined as the reference category of the BMI classification in the multinomial logistic regression analyses.  
 \*\* Defined by United Nations classification: East Asia (Japan and Taiwan), South Asia (India) and Southeast Asia (Indonesia and Malaysia).  
 \*\*\* Defined by the World Bank list of economies: high income (Japan and Taiwan), upper middle income (Malaysia) and lower middle income (India and Indonesia).  
 BMI - body mass index; BPRS - Brief Psychiatric Rating Scale; CCI - Charlson comorbidity index; DUP - duration of untreated psychosis; FGA - first-generation antipsychotics; SGA - second-generation antipsychotics; WHO - World Health Organization; a - normal range; b - at risk; c - obese I; d - obese II.

Table 2. Continues

	Underweight (n=122)	Normal range (n=537)	Overweight:		Statistical coefficient	Unadjusted P-value	Adjusted P-value*
			At risk (n=275)	Obese I (n=366)			
Rigidity, n (%)	25 (20.5)	65 (12.1)	23 (8.4)	47 (12.8)	$\chi^2=15.017$	0.004	0.035 <sup>a</sup> 0.003 <sup>b</sup> 0.241 <sup>c</sup> 0.034 <sup>d</sup>
Tremor, n (%)	20 (16.4)	95 (17.7)	41 (14.9)	72 (19.7)	$\chi^2=2.980$	0.561	0.635 <sup>a</sup> 0.835 <sup>b</sup> 0.217 <sup>c</sup> 0.811 <sup>d</sup>
Akathisia, n (%)	15 (12.3)	43 (8.0)	16 (5.8)	28 (7.7)	$\chi^2=6.988$	0.137	0.176 <sup>a</sup> 0.034 <sup>b</sup> 0.214 <sup>c</sup> 0.094 <sup>d</sup>
Dystonia, n (%)	9 (7.4)	28 (5.2)	9 (3.3)	20 (5.5)	$\chi^2=7.708$	0.103	0.511 <sup>a</sup> 0.134 <sup>b</sup> 0.721 <sup>c</sup> 0.069 <sup>d</sup>
Dyskinesia, n (%)	4 (3.3)	27 (5.0)	16 (5.8)	14 (3.8)	$\chi^2=2.661$	0.616	0.442 <sup>a</sup> 0.283 <sup>b</sup> 0.607 <sup>c</sup> 0.727 <sup>d</sup>
Overall, n (%)	23 (18.9)	84 (15.6)	50 (18.2)	63 (17.2)	$\chi^2=1.698$	0.791	0.624 <sup>a</sup> 0.926 <sup>b</sup> 0.951 <sup>c</sup> 0.997 <sup>d</sup>
Parkinsonism, n (%)	25 (34.7)	91 (34.1)	40 (27.4)	68 (34.3)	$\chi^2=2.499$	0.646	0.612 <sup>a</sup> 0.246 <sup>b</sup> 0.809 <sup>c</sup> 0.965 <sup>d</sup>

According to the WHO classification system for body weight status, the body mass index was classified into the underweight (<18.50), normal weight (18.50-24.99), overweight: pre-obese (25.00-29.99), overweight: obese I (30.00-34.99), overweight: obese II (35.00-39.99), and overweight: obese III (>40.00).  
 \* Adjusted for the effects of age, regional classification, income group, enrollment as an outpatient, duration of illness, DUP, total score on the BPRS, and antidepressants.  
 Underweight was defined as the reference category of the BMI classification in the multinomial logistic regression analyses.  
 \*\* Defined by United Nations classification: East Asia (Japan and Taiwan), South Asia (India) and Southeast Asia (Indonesia and Malaysia).  
 \*\*\* Defined by the World Bank list of economies: high income (Japan and Taiwan), upper middle income (Malaysia) and lower middle income (India and Indonesia).  
 BMI - body mass index; BPRS - Brief Psychiatric Rating Scale; CCI - Charlson comorbidity index; DUP - duration of untreated psychosis; FGA - first-generation antipsychotics; SGA - second-generation antipsychotics; WHO - World Health Organization; a - normal range; b - at risk; c - obese I; d - obese II.

## DISCUSSION

In summary, in terms of the WHO classification system for body weight status, underweight (rather than normal range, pre-obese, obese I, obese II, or obese III) was associated with greater rates of bradykinesia and muscle rigidity and lower rates of gait disturbance among Asian patients with schizophrenia. Furthermore, in terms of the Asia-specific obesity classification, underweight (rather than normal range, at risk, obese I, or obese II) was associated with greater rates of muscle rigidity and akathisia. Although different obesity classification systems were used, our findings consistently revealed that underweight was associated with a greater rate of muscle rigidity, in a stepwise pattern, among the 1448 Asian patients with schizophrenia.

Previous investigations on the relationship between BMI and muscle rigidity have been rare and insufficient. Since muscle rigidity is regarded as a typical of PD, based on the speculations about the relationship between BMI and PD (Logroscino et al. 2007, Noyce et al. 2017), an inverse relationship between BMI and muscle rigidity can be elucidated. First, the alterations in the autonomic nervous system, which are associated with appetite or metabolism regulation, may increase the vulnerability to muscle rigidity. The appetite and weight regulation is closely implicated in hypothalamus. Also, schizophrenia patients usually have structural and functional disturbances in the hypothalamus (Klomp et al. 2012). Thus, there is a possibility that the hypothalamic disturbances may precede the decrease in BMI and vulnerability to muscle rigidity in patients with schizophrenia. Second, increased body fat may have a protective effect on the vulnerability to EPS in patients with schizophrenia (Cheshire & Wszolek 2005). Namely, the levels of circulating and central insulin, which can be affected by the high level of BMI, may reduce neurodegeneration (Craft and Watson 2004). Third, brain-derived neurotrophic factor (*BDNF*) and regulatory-associated protein of mTOR (*RAPTOR*) have been proposed as the candidate genes for the link between BMI and PD (Logroscino et al. 2007, Noyce et al. 2017). Since *BDNF* is a potential therapeutic agent for neurodegenerative diseases (Nagal & Tuszynski 2011) and *RAPTOR* contributes to the links between the mTOR pathway, macroautophagy, and PD (Bove et al. 2011), these genes may also be associated with the inverse relationship between BMI and muscle rigidity. Fourth, poor nutrition-related weight loss, anemia, and other physical complications might be a contributory factor to the occurrence of muscle rigidity.

Our study has several limitations. First, the REAP-AP was designed as a cross-sectional study. The inferences about temporal causality were limited. Second, physical activity may be inversely associated with both BMI and EPS (Xu et al. 2010). Third, the possibility

that statistical significance may be different from clinical significance cannot be excluded. The potential effects of physical activity on the relationship between BMI and EPS were not adjusted in our study. Despite these limitations, our study sufficiently demonstrates that the BMI is inversely related with muscle rigidity in Asians with schizophrenia. Our findings suggest that low BMI may be a risk factor for motor rigidity irrespective of antipsychotic "typicality" and dose in patients with schizophrenia.

## CONCLUSION

Findings of the present study consistently revealed that underweight was associated with a greater rate of muscle rigidity in a stepwise pattern among Asian patients with schizophrenia. Although the mechanism underlying the inverse relationship between BMI and muscle rigidity cannot be sufficiently explained, it is speculated that low BMI may contribute to the development of muscle rigidity regardless of antipsychotic "typicality" and dose in patients with schizophrenia.

### Acknowledgements:

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (2019R1A2C1090146).

**Conflict of interest:** None to declare.

### Contribution of individual authors:

Seon-Cheol Park: analyzed the data, wrote the first draft of the manuscript, revision of manuscript, agree with results and conclusions and accept final manuscript.

Adarsh Tripathi, Ajit Avasthi, Sandeep Grover, Andi J. Tanra, Takahiro A. Kato, Toshiya Inada, Kok Yoon Chee, Mian-Yoon Chong, Shu-Yu Yang, Sih-Ku Lin, Kang Sim, Yu-Tao Xiang & Afzal Javed: revision of manuscript, agree with results and conclusions and accept final manuscript.

Norman Sartorius, Naotaka Shinfuku & Yong Chon Park: conceived and designed the study, revision of manuscript, agree with results and conclusions and accept final manuscript.

## References

1. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorder, 5th edition*. Washington, D.C.: American Psychiatric Press, 2013
2. Arora D, Mudgal J, Nampoothiri M, Mallik SB, Kinra M, Hall S, Anoopkumar-Dukie S, et al: *Interplay between adenosine receptor antagonist and cyclooxygenase inhi-*

- bitor in haloperidol-induced extrapyramidal effects in mice. *Metab Brain Dis* 2018; 33:1045-1051
3. Bove J, Martí-Áñez-Vicente M, Vila M: Fighting neurodegeneration with rapamycin: mechanistic insights. *Nat Rev Neurosci* 2011; 12:437-452
  4. Charlson ME, Pompei P, Ales KL, MacKenzie CR: A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987; 40:373-383
  5. Cheshire WP Jr, Wszolek ZK: Body mass index is reduced early in Parkinson's disease. *Parkinsonism Relat Disord* 2005; 11:35-38
  6. Craft S, Watson CG: Insulin and neurodegenerative disease: Shared and specific mechanisms. *Lancet Neurol* 2004; 3:169-178
  7. De Fonseca FR, Gorriti MA, Bilbao A, Esuero L, Garcia-Segura LM, Piomelli D, et al: Role of the endogenous cannabinoid system as a modulator of dopamine transmission: Implications for Parkinson's disease and schizophrenia. *Neurotox Res* 2001; 3:23-35
  8. Haleem DJ: 5-HT<sub>1A</sub> receptor-dependent control of nigrostriatal dopamine neurotransmission in the pharmacology of Parkinson's disease and schizophrenia. *Behav Pharmacol* 2015; 26:45-58
  9. International Obesity Task Force. Tokyo, Japan: World Health Organization Western Pacific Region, 2000
  10. Inada T: DIEPSS: A Second-Generation Rating Scale for Antipsychotic-induced Extrapyramidal Symptoms: Drug-induced Extrapyramidal Symptoms Scale. Tokyo, Japan: Seiwa Shoten Publishers, Inc, 2009
  11. Kim JH, Jung HY, Kang UK, Jeong SH, Ahn YM, Byun HJ, et al: Metric characteristics of the drug-induced extrapyramidal symptoms scale: A practical combined rating scale for drug-induced movement disorders. *Mov Disord* 2002; 17:1354-1359
  12. Klomp A, Koolschijn PC, Hulshoff Pol HE, Kahn RS, Haren NE: Hypothalamus and pituitary volume in schizophrenia: a structural MRI study. *Int J Neuropsychopharmacol* 2012; 15:281-288
  13. Leucht S, Kane JM, Kissling W, Hamann J, Etschel E, Engel R: Clinical implications of brief psychiatric rating scale score. *Br J Psychiatry* 2005; 187:366-371
  14. Logroscino G, Sesso HD, Paffenbarger RS, Lee IM: Body mass index and risk of Parkinson's disease: a prospective cohort study. *Am J Epidemiol* 2007; 166:1186-1190
  15. Nagahara AH, Tuszynski MH: Potential therapeutic uses of BDNF in neurological and psychiatric disorders. *Nat Rev Drug Discov* 2011; 10:209-219
  16. Noyce AJ, Kila DA, Hermani G, Nicloas A, Price TR, De Pablo-Fernandez, et al: Estimating the causal influence of body mass index on risk of Parkinson disease: A Mendelian randomization study. *PLoS Med* 2017; 14:e1002314
  17. Ohno Y: Therapeutic role of 5-HT<sub>1A</sub> receptors in the treatment of schizophrenia and Parkinson's disease. *CNS Neurosci Ther* 2011; 17:58-65
  18. Overall JE, Gorham DR: The brief psychiatric rating scale. *Psychol Rep* 1962; 10:779-812
  19. Park YC, Yang S-Y, Chong M-Y, Kanba S, Sartorius N, Shinfuku N, et al: Differences in High Dose Antipsychotic Prescription in Patients with Schizophrenia in Asian Countries/Areas: Findings from the REAP-AP Study. *Psychiatry Investig* 2018; 15:1007-1008
  20. Quan H, Sundararajan V, Halfon P, Fong A, Burnand B, Luthi JC, et al: Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care* 2005; 43:1130-1139
  21. Tandon R, Nasrallah HA, Keshavan MS: Schizophrenia, "just the facts" 5. Treatment and prevention past, present and future. *Schizophr Res* 2010; 122:1-23
  22. World Health Organization: *The ICD-10 Classification of Mental and Behavioral Disorders, Clinical Descriptions and Diagnostic Guidelines*. Geneva: World Health Organization, 1992
  23. World Health Organization: *Obesity: Preventing and Managing the Global Epidemic*. Geneva: World Health Organization, 1998
  24. World Health Organization: *Anatomical Therapeutic Chemical (ATC) classification system*. Available at: [https://www.whocc.no/atc/structure\\_and\\_principles/](https://www.whocc.no/atc/structure_and_principles/) Accessed 01-28 2019
  25. Xu Q, Park Y, Huang X, Hollenbeck A, Blair A, Schatzkin A, Chen H: Physical activities and future risk of Parkinson disease. *Neurology* 2010; 75:341-348

Correspondence:

Seon-Cheol Park, MD, PhD  
Department of Psychiatry, Inje University Haeundae Paik Hospital  
875, Haeun-daero, Haeundae-gu, Busan 48108, Republic of Korea  
E-mail: cogito-ergo-sum@hanmail.net

# RELATIONSHIP BETWEEN BODY MASS INDEX AND EXTRAPYRAMIDAL SYMPTOMS IN ASIAN PATIENTS WITH SCHIZOPHRENIA: THE RESEARCH ON ASIAN PSYCHOTROPIC

## ORIGINALITY REPORT

28%

SIMILARITY INDEX

23%

INTERNET SOURCES

23%

PUBLICATIONS

13%

STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to University of Nottingham Student Paper	2%
2	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	2%
3	<a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	2%
4	<a href="http://knpa.inforang.com">knpa.inforang.com</a> Internet Source	2%
5	<a href="http://www.j3.jstage.jst.go.jp">www.j3.jstage.jst.go.jp</a> Internet Source	2%
6	<a href="http://www.psychiatryinvestigation.org">www.psychiatryinvestigation.org</a> Internet Source	1%
7	<a href="http://epdf.tips">epdf.tips</a> Internet Source	1%
8	Parker, Georgina, David Lie, Dan J. Siskind, Melinda Martin-Khan, Beverly Raphael, David Crompton, and Steve Kisely. "Mental health	1%

implications for older adults after natural disasters – a systematic review and meta-analysis", International Psychogeriatrics, 2016.

Publication

9

[helda.helsinki.fi](https://helda.helsinki.fi)

Internet Source

1 %

10

Sun Jae Jung, Sue Kyung Park, Aesun Shin, Sang-Ah Lee, Ji-Yeob Choi, Yun-Chul Hong, Keun-Young Yoo, Jong-Koo Lee, Daehee Kang. "Body mass index at age 18–20 and later risk of spontaneous abortion in the Health Examinees Study (HEXA)", BMC Pregnancy and Childbirth, 2015

Publication

1 %

11

[www.mdpi.com](https://www.mdpi.com)

Internet Source

1 %

12

Keren Wen, Youjin Zhao, Qiyong Gong, Ziyu Zhu et al. "Cortical thickness abnormalities in patients with first episode psychosis: a meta-analysis of psychoradiologic studies and replication in an independent sample", Psychoradiology, 2021

Publication

1 %

13

[Www.psychiatryinvestigation.org](https://www.psychiatryinvestigation.org)

Internet Source

1 %

14

"Issue Information", The Kaohsiung Journal of Medical Sciences, 2020

Publication

1 %

---

15 Satya Krishna Ramachandran. "Independent Predictors and Outcomes of Unanticipated Early Postoperative Tracheal Intubation after Nonemergent, Noncardiac Surgery :", *Anesthesiology*, 05/2011  
Publication <1 %

---

16 [www.frontiersin.org](http://www.frontiersin.org)  
Internet Source <1 %

---

17 [www.rxfiles.ca](http://www.rxfiles.ca)  
Internet Source <1 %

---

18 Submitted to University of Edinburgh  
Student Paper <1 %

---

19 Seon - Cheol Park, Eun Young Jang, Yu - Tao Xiang, Shigenobu Kanba et al. "Network analysis of the depressive symptom profiles in Asian patients with depressive disorders: Findings from the Research on Asian Psychotropic Prescription Patterns for Antidepressants (REAP - AD)", *Psychiatry and Clinical Neurosciences*, 2020  
Publication <1 %

---

20 Park, S.-C., M.-S. Lee, N. Shinfuku, N. Sartorius, and Y. C. Park. "Gender differences in depressive symptom profiles and patterns of psychotropic drug usage in Asian patients with depression: Findings from the Research on Asian Psychotropic Prescription Patterns <1 %

for Antidepressants study", Australian and  
New Zealand Journal of Psychiatry, 2015.

Publication

---

21 [journals.plos.org](http://journals.plos.org) <1 %  
Internet Source

---

22 [www.scielo.br](http://www.scielo.br) <1 %  
Internet Source

---

23 [kumel.medlib.dsmc.or.kr](http://kumel.medlib.dsmc.or.kr) <1 %  
Internet Source

---

24 [kupdf.net](http://kupdf.net) <1 %  
Internet Source

---

25 [psychiatryinvestigation.org](http://psychiatryinvestigation.org) <1 %  
Internet Source

---

26 Galli, M. M., N. M. Protzman, E. M. Mandelker,  
A. Malhotra, E. Schwartz, and S. A. Brigido.  
"Comparing Tendinous and Ligamentous  
Ankle Pathology in Atraumatic Overweight  
and Nonoverweight Patients: A  
Comprehensive MRI Review", Foot & Ankle  
Specialist, 2014.  
Publication

---

27 [www.doc-cafes.com](http://www.doc-cafes.com) <1 %  
Internet Source

---

28 Shi-Wei Xu, Min Dong, Qinge Zhang, Shu-Yu  
Yang et al. "Clozapine prescription pattern in  
<1 %

patients with schizophrenia in Asia: The REAP survey (2016)", Psychiatry Research, 2019

Publication

29

Su - Min Jeong, Kyungdo Han, Dahye Kim, Sang Youl Rhee, Wooyoung Jang, Dong Wook Shin. "Body mass index, diabetes, and the risk of Parkinson's disease", Movement Disorders, 2019

Publication

<1 %

30

Submitted to Universiti Kebangsaan Malaysia

Student Paper

<1 %

31

M. P. Bishop, George M. Simpson, C. W. Dunnett, Harriet Kiltie. "Efficacy of loxapine in the treatment of paranoid schizophrenia", Psychopharmacology, 1977

Publication

<1 %

32

[bmcendocrdisord.biomedcentral.com](http://bmcendocrdisord.biomedcentral.com)

Internet Source

<1 %

33

[www.nature.com](http://www.nature.com)

Internet Source

<1 %

34

TOSHIYA INADA, GOHEI YAGI. "Current topics in tardive dyskinesia in Japan", Psychiatry and Clinical Neurosciences, 1995

Publication

<1 %

35

[serval.unil.ch](http://serval.unil.ch)

Internet Source

<1 %

36

[www.jstage.jst.go.jp](http://www.jstage.jst.go.jp)

Internet Source

&lt;1 %

37

Pierre Lavaud, Kibby McMahon, Marina Sánchez Rico, Cécile Hanon et al. "Long-term care utilization within older adults with schizophrenia: Associated factors in a multicenter study", *Psychiatry Research*, 2022

Publication

&lt;1 %

38

Seon-Cheol Park. "Network Analysis of Depressive Symptomatology in Elderly Patients with Major Depressive Disorder", *Current Psychiatry Research and Reviews*, 2021

Publication

&lt;1 %

39

Submitted to University of Central Lancashire

Student Paper

&lt;1 %

40

[repository.umac.mo](http://repository.umac.mo)

Internet Source

&lt;1 %

41

[www.nutricionhospitalaria.com](http://www.nutricionhospitalaria.com)

Internet Source

&lt;1 %

42

Georg Juckel, Antje Müller-Schubert, Wolfgang Gaebel, Ulrich Hegerl. "Residual symptoms and P300 in schizophrenic outpatients", *Psychiatry Research*, 1996

Publication

&lt;1 %

43

[mafiadoc.com](http://mafiadoc.com)

Internet Source

<1 %

44

Matthew W Rogers. "The association of BMI and knee pain among persons with radiographic knee osteoarthritis: A cross-sectional study", BMC Musculoskeletal Disorders, 2008

Publication

<1 %

45

Submitted to Open University Malaysia

Student Paper

<1 %

46

[hrmars.com](http://hrmars.com)

Internet Source

<1 %

47

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

48

[www.atsjournals.org](http://www.atsjournals.org)

Internet Source

<1 %

49

[www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

50

Kim, Sung-Wan, Jin-Sang Yoon, Yong Sik Kim, Yong-Min Ahn, Chul-Eung Kim, Hyo-Jin Go, Ik-Seung Chee, Sung-Won Jung, Young-Chul Chung, Young-Don Kim, Soohyun Joe, Jonghun Lee, Young-Joon Kwon, Bo-Hyun Yoon, and Young-Myo Jae. "The effect of paliperidone extended release on subjective well-being and responses in patients with schizophrenia",

<1 %

# Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2012.

Publication

51

[vsip.info](http://vsip.info)

Internet Source

<1 %

52

Hye Soon Park, Jee-Young Park, Hong-Jun Cho. "Attitudes and Reported Practice for Obesity Management in Korea After Introduction of Anti-obesity Agents", Journal of Korean Medical Science, 2005

Publication

<1 %

53

[bmchealthservres.biomedcentral.com](http://bmchealthservres.biomedcentral.com)

Internet Source

<1 %

54

[foreigner.gilhospital.com](http://foreigner.gilhospital.com)

Internet Source

<1 %

55

[www.koreascience.kr](http://www.koreascience.kr)

Internet Source

<1 %

56

"Handbook of Neurotoxicity", Springer Nature, 2014

Publication

<1 %

57

Joonho Choi, Hyung-Jun Yoon, Jae Hong Park, Yukako Nakagami et al. "Network Analysis-Based Disentanglement of the Symptom Heterogeneity in Asian Patients with Schizophrenia: Findings from the Research on Asian Psychotropic Prescription Patterns for

<1 %

# Antipsychotics", Journal of Personalized Medicine

Internet Source

---

58

Chika Kubota, Toshiya Inada, Shih-Ku Lin, Ajit Avasthi et al. "The factor structure of extrapyramidal symptoms evaluated using the DIEPSS in patients with schizophrenia: results from the 2016 REAP AP-4 study", Research Square, 2021

Publication

---

<1 %

59

Ilona Csoti, Wolfgang H. Jost, Heinz Reichmann. "Parkinson's disease between internal medicine and neurology", Journal of Neural Transmission, 2015

Publication

---

<1 %

60

Kamini Rajaratnam, Yu-Tao Xiang, Adarsh Tripathi, Helen F.K. Chiu et al. "Clinical Use of Mood Stabilizers With Antidepressants in Asia", Journal of Clinical Psychopharmacology, 2017

Publication

---

<1 %

61

Masoom M. Abbas, Zheyu Xu, Louis C.S. Tan. "Epidemiology of Parkinson's Disease-East Versus West", Movement Disorders Clinical Practice, 2018

Publication

---

<1 %

62

Seon-Cheol Park, Min-Soo Lee, Sang Woo Hahn, Shigenobu Kanba et al. "Clinical

<1 %

Characteristics and Psychotropic Prescribing Patterns Associated with impaired Concentration in Asians with Depressive Disorders: The REAP-AD Study", The Tohoku Journal of Experimental Medicine, 2017

Publication

63

[hrcak.srce.hr](http://hrcak.srce.hr)

Internet Source

<1 %

64

[www.lib.kobe-u.ac.jp](http://www.lib.kobe-u.ac.jp)

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On