

The concordance of triglyceride glucose index (TyG index) and homeostatic model assessment for insulin resistance (Homa-IR) in non-diabetic subjects of adult Indonesian males

by Andi Makbul Aman

FILE	NT_FOR_INSULIN_RESISTANCE__HOMA-IR__IN_NON-DIABETIC_SUBJECTS.PDF (1.1M)	WORD COUNT	3048
TIME SUBMITTED	11-FEB-2021 03:59AM (UTC+0700)	CHARACTER COUNT	15626
SUBMISSION ID	1506557696		



4

The concordance of triglyceride glucose index (TyG index) and homeostatic model assessment for insulin resistance (Homa-IR) in non-diabetic subjects of adult Indonesian males



Makbul Aman^a, Dewi Resnawita^{b,*}, Haerani Rasyid^c, Hasyim Kasim^c, Syakib Bakri^c,
Husaini Umar^a, Nu'man AS. Daulat^d, Arifin Seweng^e

^a Division of Endocrinology Diabetes and Metabolism, Department of Internal Medicine, Faculty of Medicine, Universitas Hasanuddin, Indonesia

^b Department of Internal Medicine, Faculty of Medicine, Universitas Hasanuddin, Indonesia

^c Division of Nephrology, Department of Internal Medicine, Faculty of Medicine, Universitas Hasanuddin, Indonesia

^d Division of Gastroenterology and Hepatology, Department of Internal Medicine, Faculty of Medicine, Universitas Hasanuddin, Indonesia

^e Department of Biostatistics, Faculty of Public Health, Universitas Hasanuddin, Indonesia

ARTICLE INFO

Keywords:

Insulin resistance
Type 2 diabetes mellitus
Homeostasis model assessment of insulin resistance
Triglyceride glucose index

ABSTRACT

Background: Recently, an examination with the TyG index has developed as an alternative to the gold standard, by which this measurement has a relatively high sensitivity in assessing IR. The measurement with TyG index only requires triglyceride levels and fasting blood glucose and is expected to be an examination instrument used to assess IR available in all health facilities in developing countries, especially in Indonesia. In this study, we analyzed the concordance between the TyG index and HOMA-IR in assessing IR in nondiabetic subjects of adult Indonesian males.

Method: A cross sectional study was conducted at the outpatient installation of Wahidin Sudirohusodo Hospital and Hasanuddin University Hospital during March until June 2019. An examination of FBG, TyG and fasting insulin was performed. The statistical test results were considered significant if the *p* value was < 0.05.

Result: The 88 healthy male subjects, the mean age was 51.15 ± 6.8 years. The ROC curve showed that the cut-off value of the TyG index in predicting HOMA-IR was 4.66, with the sensitivity of 86.2% and specificity of 44.1%. The OR value revealed that subjects with a TyG index of ≥ 4.66 indicated a 5-time greater risk for IR than subjects with a TyG index < 4.66.

Conclusion: There was a significant concordance between the TyG index and the HOMA-IR in the subjects of adult Indonesian males.

1. Background

Insulin resistance (IR) is broadly defined as a subnormal biological response to normal insulin concentration.¹ It precedes the development of type 2 diabetes mellitus and cardiovascular disease. Therefore, it is very important to establish a concise and practical method for identifying individuals at high risk of IR.^{2,3}

The gold standard in IR diagnosis is HEC, but this method is invasive, complex, and expensive. Therefore, it is difficult to apply on daily clinical screening and population-based epidemiological studies.^{2,4,5}

These days, the emerging methods performed for clinical screening are the HOMA-IR, the QUIKI index and the Matsuda index, but each of them is not currently available in all health services.^{4,6} HOMA-IR is and

one of the valid inspection instruments and the most common methods used for IR measurement, but this examination requires plasma insulin levels which, however, are currently not available on demand.^{7,8}

In recent years, an examination with the TyG index has developed. This approach has a fairly high sensitivity in assessing IR. The measurement with TyG index only requires triglyceride levels and fasting glucose. Thus, the TyG index can be used as a test instrument to assess IR in all health facilities in developing countries, especially in Indonesia.^{5,9} However, since

This study aims to investigate the concordance between TyG index and HOMA-IR in assessing IR in nondiabetic patients.

* Corresponding author.

E-mail address: dewewita@gmail.com (D. Resnawita).

<https://doi.org/10.1016/j.cegh.2020.09.003>

Received 2 July 2020; Received in revised form 5 August 2020; Accepted 11 September 2020

Available online 19 September 2020

2213-3984 / © 2020 INDIACLEN. Published by Elsevier, a division of RELX India, Pvt. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Characteristics of the subjects (n = 88).

Variables	Minimum	Maximum	Mean	SD
Age	40	65	51,15	6,83
FBG	73	125	94,92	9,94
TG	63	477	170,69	88,10
Fasting Insulin	2,0	28	8,29	7,27
HOMA-IR	0,36	7,48	2,00	1,82
TyG Index	4,34	5,50	4,79	0,25

2. Method

A cross sectional study was performed at the outpatient installation of Wahidin Sudirohusodo Makassar hospital and Hasanuddin University hospital during March 2019–June 2019. This study received the ethical clearance approval from the Ethics Commission of Biomedical Research in Humans, Faculty of Medicine, Universitas Hasanuddin, Makassar, number of 231/UN4.6.4.5.31/PP36/2019. The approval of medical actions and the signing of informed consent were reported by providing the background, objectives and benefits of the study, as well as blood drawings that the study subjects underwent.

Consecutive sampling was performed in this study. 88 nondiabetic subjects who met the inclusion criteria were included in the study. The inclusion criteria were men, 40–65 years old without diabetes or a history of diabetes without consumption of anti-diabetes drugs, anti-dyslipidemia, anti-hypertension and/or without consumption of the aforementioned drugs in the period of > 1 month healthy (no signs of sickness), and with an approval of involvement in the study by signing the informed consent. The subjects were diagnosed with DM if the FBG was ≥ 126 mg/dL and categorized as IR if the tertile was 3 HOMA-IR. In addition, healthy subjects were those with a good physical, mental and social condition and were not only determined by the absence of illness or weakness. The HOMA-IR score was resulted from $[\text{FBG (mg/dL)} \times \text{insulin } (\mu\text{U/L})/405]$, and TyG index was calculated by $\text{Ln} [\text{TG (mg/dL)} \times \text{FBG (mg/dL)}]/2$.

The data were analyzed by using SPSS ver. 22 with descriptive statistical analysis and Chi square statistical test, Pearson's correlation, ROC curves, kappa and logistic regression. The statistical test results were considered significant if the p value was < 0.05.

Table 2
TyG index and HOMA-IR on ROC curve.

TyG Index	HOMA-IR		Total	p	Sensitivity	Specificity	OR ^a
	IR	Non-IR					
$\geq 4,66$	25	33	58	0,005	86,2%	44,1%	5,0
< 4,66	4	26	30				
Total	29	59	88				

OR: Odds Ratio CI: Confidence Interval.

^a 95% CI (1,52–15,93).

Table 3
TyG index dengan HOMA-IR correlation.

Variable	Statistics	Homa-IR
TyG index	Pearson Correlation	0,436
	Sig. (2-tailed)	0,000
	n	88

3. Results

Table 1 shows the results of the data analysis among 88 healthy male subjects aged 40–65 years. It can be seen that the mean age was 51.15 ± 6.8 years. The characteristics of the study subjects indicated the rate of FBG at 94.92 ± 9.94 , TG at 170.69 ± 88.10 , fasting insulin at 8.29 ± 7.27 , HOMA-IR at 2.0 ± 1.82 , and TyG index at 4.79 ± 0.25 .

The value of HOMA-IR and TyG index was divided based on tertile because there was no cutoff in assessing IR either from HOMA-IR value or TyG index value. Tertile 1.2 of HOMA-IR was < 2.24, while tertile 3 of HOMA-IR was ≥ 2.24 . On the other hand, tertile 1.2 of TyG index was < 4.89 and tertile 3 of TyG index remained at ≥ 4.89 .

Fig. 1 illustrates the results of the ROC curve analysis. It showed that the value of Area Under Curve (AUC) was 0.701 ($p < 0.01$), which indicated that the TyG Index was fairly sensitive in predicting HOMA-IR values with a cutoff value of 4.66.

Table 2 shows that there was a significant concordance between TyG index and HOMA-IR ($p < 0.01$), with a sensitivity of 86.2% and specificity of 44.1%. OR values indicated that subjects with TyG index at ≥ 4.66 were 5 times more at risk of IR than those with TyG index




Fig. 1. ROC curve of TyG index on HOMA-IR.




Fig. 2. The positive correlation between TyG index and HOMA-IR.

at < 4.66.

Tal 19³ reveals that there is a significant positive correlation between the TyG index and the HOMA-IR: the higher the TyG index, the higher the HOMA-IR ($p < 0.001$). The result of the correlation test conf³⁶; the coefficient value at $r = 0.436$. Fig. 2 illustrates the positive correlation between the TyG index and the HOMA-IR.

4. Discussion

The study was not conducted in female subjects due the influ³ce of menstruation on blood glucose. Several studies reported that certain physiological parameters, such as blood glucose is affected by menstruation. In healthy non diabetic women, some researches indicated a worsening in glucose tolerance, as assessed by the oral glucose tolerance test (OGTT), during the luteal phase (secretory phase).¹⁰

This study shows that the average HOMA-IR value remained at 2.0 ± 1.82 . A similar study by Lina Y et al. (2009) in Makassar showed different results with a HOMA-IR value of 1.90 ± 1.10 .¹¹ Another study by Ritawaty et al. (20³⁵) found an average HOMA-IR value of 3.2 ± 3.40 .¹² Gayozo-Diz et al. stated that the value of HOMA-IR differed according to ethnicity, clinical estimation methods,³⁴ the condition of the metabolism of the population.⁷ The average value of the TyG index in this study was 4.679 ± 0.25 . In the study by Wongsari MH et al. (2018) in Makassar, however, the value of the TyG index was 4.59 ± 0.22 .¹³

This study shows that there is a significant concordance between the TyG index and the HOMA-IR index. TyG index with a high value indicates a high HOMA-IR value ($p < 0.01$), with a sensitivity of 86.2%, instead of a low specificity at 44.1% and a cutoff at 4.66. This is in line⁸ th a research by Fahimeh et al. (2014) which also found a significant concordance between the TyG and HOMA-IR indices in the population of obese women in Iran.¹⁴ Another study by Guerrero-R⁶ero et al. (2010 and 2016) in Mexico also showed the sar⁸ result between the TyG index and the HOMA-IR.^{2,15} The significant concordance between the TyG index and the HOMA-IR is useful in recognizing insulin resistance among subjects with var¹⁹ grades of body weight and glucose tolerance. There is a remarkable correlation between the TyG index and several obesity factors including BMI, fasting insulin, and LP. This correlation³³ uch greater than that with HOMA-IR.^{9,16} Irace also reported that the TyG-Index had a more accurate association with the risk factors for carotid atherosclerosis compared to HOMA-IR.¹⁷

32

The TyG index has been validated in several pop¹⁵ons around the world for use as IR screening.^{18,19} This study shows that the TyG index has high sensitivity but low specificity. A research by Simental-Mendia et al. (2008) in a healthy population aged 18–65 years in Mexico also showed the same results.⁶ (15) the other hand, a study by Guerrero-Romero et al. (2010) shows that the TyG index has high sensitivity and specificity.¹⁵ Another study by Bastard et al. (2012) of 163 postmenopausal nondiabetic women suggested that, in a large population, further investigation on TyG index was needed.¹⁴ (16) themore, a cross sectional study of 82 patients in Brazil (2011) showed that the TyG ind² indicated better results than HOMA-IR.^{9,16}

High-sensitivity test, such as the TyG index, has fewer false negatives and is useful in identifying subjects with IR on clinical examination. This is very beneficial because fasting insulin measurement is not necessary in the TyG index assessment, so it can be used as an alternative in assessing IR, especially in the first-level public health facilities² at improve early detection of IR and suggest early lifestyle changes.^{2,6} On the other hand, the TyG index in identifying the proportion of subject² without any disease (specificity) is low, so there are false positives that limit its use for screening purposes.⁶ Therefore, this study suggests that subjects with TyG index of < 4.66 are very likely to have IR. Conversely, those with TyG index of ≥ 4.66 indicate the possibility of IR and still require additional examinations, such as HOMA-IR. According to the OR value, the subjects with TyG index of ≥ 4.66 had a 5-time-greater risk of experiencing IR than those with TyG index of < 4.66.

In con⁶usion, this study shows that there is a significant concordance between the TyG index and the HOMA-IR index in adult Indonesian men. This result suggests that screening for IR in all healthy individuals should be performed, considering that IR can occur at various age levels. Furthermore, TyG index examination can be applied at all health facilities. Further researches with larger population are needed in the following periods of time.

5. Study limitation

The limitation of this study can be identified over the sens⁶ivity and specificity of the TyG index that was compared only with HOMA-IR, which is not the gold standard in diagnosing IR.

Authors' contribution

All persons who meet authorship criteria are listed as authors, and all authors certify that they have participated sufficiently in the work to take public responsibility for the content, including participation in the concept, design, analysis, writing, or revision of the manuscript. Furthermore, each author certifies that this material or similar material has not been and will not be submitted to or published in any other publication.

Category 1.

Conception and design of study: Makbul Aman, Dewi Resnawita, Ha 31 i Rasyid.

Acquisition of data: Dewi Resnawita.

Analysis and Interpretation of Data: Arifin Seweng, Makbul Aman, Dewi Resnawita, Haerani Rasyid.

Category 2.

Drafting the manuscript: Makbul Aman, Dewi Resnawita, Haerani Rasyid. 43 28

Revising the manuscript for important intellectual content: Hasyim Kasim, Syakib Bakri, Husaini Umar, Nu'man AS Daud.

Authors approval

21

All authors read and approved the final version of the manuscript.

Acknowledgement

We 1 would like to thank all subjects who have cooperated with the authors in conducting this research. All of the authors are responsible for the content and writing of the paper.

References

- Mantzoros C. Insulin resistance: definition and clinical spectrum. . Available from: <https://www.uptodate.com/contents/insulin-resistance-definition-and-clinical-spectrum/contributors>; 2018.
- Guerrero-Romero F, Villalobos-Molina R, Jiménez-Flores JR, et al. Fasting triglycerides and glucose index as a diagnostic test for insulin resistance in young adults. *Arch Med Res*. 2016;47(5):382–387.
- Do HD, Lohsoonthorn V, Jiamjarasrangsri W, Lertmaharit S, Williams MA. Prevalence of insulin resistance and its relationship with cardiovascular disease risk factors among Thai adults over 35 years old. *Diabetes Res Clin Pract*. 2010;89(3):303–308.
- Gutch M, Kumar S, Razi SM, Gupta K, Gupta A. Assessment of insulin sensitivity/resistance. *Indian J Endocrinol Metab*. 2015;19(1):160–164.
- Radziuk J. CLINICAL REVIEW 119 insulin sensitivity and its Measurement : structural commonalities among the methods *. 2015;85(12):4426–4433.
- Simental-Mendía LE, Rodríguez-Morán M, Guerrero-Romero F. The product of fasting glucose and triglycerides as surrogate for identifying insulin resistance in apparently healthy subjects. *Metab Syndr Relat Disord*. 2008;6(4):299–304.
- Gayoso-Diz P, Otero-González A, Rodríguez-Alvarez MX, et al. Insulin resistance (HOMA-IR) cut-off values and the metabolic syndrome in a general adult population: effect of gender and age: EPIRCE cross-sectional study. *BMC Endocr Disord*. 2013;13.
- Antuna-Puente B, Disse E, Rabasa-Lhoret R, Laville M, Capeau J, Bastard JP. How can we measure insulin sensitivity/resistance? *Diabetes Metab*. 2011;37(3):179–188.
- Vasques ACJ, Novaes FS, de Oliveira M da S, et al. TyG index performs better than HOMA in a Brazilian population: a hyperglycemic clamp validated study. *Diabetes Res Clin Pract*. 2011;93(3):8–10.
- Bennal A, Kerure S. Glucose handling during menstrual cycle. *Int J Reprod Contraception, Obstet Gynecol*. 2013;2(3):284–287.
- Lina Y, Lawrence GS, Wijaya A, As'ad S. Fibroblast growth factor 21 (FGF21), free fatty acid (FFA), high sensitivity C-reactive protein (hsCRP) and homeostasis model assessment of insulin resistance (HOMA-IR) among Indonesian obese non-diabetic males. *Indones Biomed J*. 2009;1(3):76.
- Ritawaty R, Sukmawati IR, Patellongi I, Sandra F. Pathomechanism of insulin resistance in men with central obesity: correlation of GGT, GPx, hs-CRP and plasma total cysteine. *Indones Biomed J*. 2013;5(2):101.
- Wongsari MH. Kesesuaian triglyceride glucose index (TyG index) dan homeostatic model assessment for insulin resistance (HOMA-IR) pada pria dewasa dengan. *Obesitas Sentral*. 2018;1:102.
- Mohammadabadi F, Vafaiyan Z, Hosseini SM, Aryaie M, Eshghinia S. Assessment of insulin resistance with two methods: HOMA-IR and TyG index in Iranian obese women. *Iran J Diabet. Obes*. 2014;6(1):23–27.
- Guerrero-Romero F, Simental-Mendía LE, González-Ortiz M, et al. The product of triglycerides and glucose, a simple measure of insulin sensitivity. Comparison with the euglycemic-hyperinsulinemic clamp. *J Clin Endocrinol Metab*. 2010;95(7):3347–3351.
- Bastard JP, Lavoie ME, Messier V, Prud'homme D, Rabasa-Lhoret R. Evaluation of two new surrogate indices including parameters not using insulin to assess insulin sensitivity/resistance in non-diabetic postmenopausal women: a MONET group study. *Diabetes Metab*. 2012;38(3):258–263.
- Trace C, Carallo C, Scavelli FB, et al. Markers of insulin resistance and carotid atherosclerosis. A comparison of the homeostasis model assessment and triglyceride glucose index. *Int J Clin Pract*. 2013;67(7):665–672.
- Lee S, Kwon H, Park Y, et al. Predicting the development of diabetes using the product of triglycerides and Glucose : the chungju metabolic disease cohort. *GMC) Study*. 2014;9(2).
- Du T, Yuan G, Zhang M, Zhou X, Sun X, Yu X. *Clinical Usefulness of Lipid Ratios , Visceral Adiposity Indicators , and the Triglycerides and Glucose Index as Risk Markers of Insulin Resistance*. vols. 1–10. 2014; 2014.

The concordance of triglyceride glucose index (TyG index) and homeostatic model assessment for insulin resistance (Homa-IR) in non-diabetic subjects of adult Indonesian males

ORIGINALITY REPORT

%29
SIMILARITY INDEX

%19
INTERNET SOURCES

%24
PUBLICATIONS

%7
STUDENT PAPERS

PRIMARY SOURCES

1 www.mdpi.com **%2**
Internet Source

2 Luis E. Simental-Mendía. "The Product of Fasting Glucose and Triglycerides As Surrogate for Identifying Insulin Resistance in Apparently Healthy Subjects", Metabolic Syndrome and Related Disorders, 12/2008 **%2**
Publication

3 Bennal, Amruta, and Sudha Kerure. "Glucose handling during menstrual cycle", International Journal of Reproduction Contraception Obstetrics and Gynecology, 2013. **%1**
Publication

4 www.ijem.in **%1**
Internet Source

5 Submitted to Georgia Gwinnett College **%1**
Student Paper

Qu, Chunmei, Xiaoxin Zhou, Gangyi Yang, Ling

6

Li, Hua Liu, and Zerong Liang. "The natural logarithm of zinc- α 2-glycoprotein/HOMA-IR is a better predictor of insulin sensitivity than the product of triglycerides and glucose and the other lipid ratios", Cytokine, 2016.

Publication

% 1

7

www.kcdcphrp.org

Internet Source

% 1

8

www.science.gov

Internet Source

% 1

9

Eirini Dikaiakou, Elpis Athina Vlachopapadopoulou, Stavroula A. Paschou, Fani Athanasouli et al. "Triglycerides-glucose (TyG) index is a sensitive marker of insulin resistance in Greek children and adolescents", Endocrine, 2020

Publication

% 1

10

Isaac Bador Kamal Lettor, Paschal Awingura Apanga, Maxwell Tii Kumbeni, Ramatu Akunvane, Robert Akparibo. "Knowledge and perceptions on COVID-19 among Senior High School students in Ghana: a cross-sectional study", Cold Spring Harbor Laboratory, 2020

Publication

% 1

11

"EDUCATION", APLAR Journal of Rheumatology, 2006

Publication

% 1

12

hdl.handle.net

Internet Source

% 1

13

irep.iium.edu.my

Internet Source

% 1

14

Atsushi Nakagomi, Yuko Sunami, Yohei Kawasaki, Takehiko Fujisawa, Yoshio Kobayashi. "Sex difference in the association between surrogate markers of insulin resistance and arterial stiffness", *Journal of Diabetes and its Complications*, 2020

Publication

% 1

15

dmsjournal.biomedcentral.com

Internet Source

% 1

16

Wen Guo, Jing Lu, Pei Qin, Xiaona Li, Wenfang Zhu, Juan Wu, Nianzhen Xu, Qun Zhang. "The triglyceride-glucose index is associated with the severity of hepatic steatosis and the presence of liver fibrosis in non-alcoholic fatty liver disease: a cross-sectional study in Chinese adults", *Lipids in Health and Disease*, 2020

Publication

% 1

17

zombiedoc.com

Internet Source

% 1

18

pt.scribd.com

Internet Source

% 1

João Carlos Locateli, Wendell Arthur Lopes,

19

Caroline Ferraz Simões, Gustavo Henrique de Oliveira et al. "Triglyceride/glucose index is a reliable alternative marker for insulin resistance in South American overweight and obese children and adolescents", Journal of Pediatric Endocrinology and Metabolism, 2019

Publication

% 1

20

epub.ub.uni-muenchen.de

Internet Source

% 1

21

link.springer.com

Internet Source

% 1

22

Submitted to Grand Canyon University

Student Paper

% 1

23

Erna Amin, Bambang Rahardjo, Kusworini. "Aromatherapy oils of lavender (*Lavandula angustifolia*) inhalance reduced norepinephrine levels of woman with postpartum blues based on Edinburgh postpartum depression scale", AIP Publishing, 2020

Publication

% 1

24

B Kang, Y Yang, E Y Lee, H K Yang, H-S Kim, S-Y Lim, J-H Lee, S-S Lee, B-K Suh, K-H Yoon. "Triglycerides/glucose index is a useful surrogate marker of insulin resistance among adolescents", International Journal of Obesity, 2017

Publication

<% 1

25 Kim, Sang-Hoon, Jae-Youn Moon, Yeong Min Lim, Kyung Ho Kim, Woo-In Yang, Jung-Hoon Sung, Seung Min Yoo, In Jai Kim, Sang-Wook Lim, Dong-Hun Cha, and Seung-Yun Cho. "Association of insulin resistance and coronary artery remodeling: an intravascular ultrasound study", Cardiovascular Diabetology, 2015. Publication <% 1

26 Hau D. Do, Vitool Lohsoonthorn, Wiroj Jiamjarasrangi, Somrat Lertmaharit, Michelle A. Williams. "Prevalence of insulin resistance and its relationship with cardiovascular disease risk factors among Thai adults over 35 years old", Diabetes Research and Clinical Practice, 2010 Publication <% 1

27 www.onlinejacc.org Internet Source <% 1

28 onlinelibrary.wiley.com Internet Source <% 1

29 ir.ymlib.yonsei.ac.kr Internet Source <% 1

30 synapse.koreamed.org Internet Source <% 1

31 Ariel Berger, Kevin Bozic, Brett Stacey, John Edelsberg, Alesia Sadosky, Gerry Oster. <% 1

"Patterns of pharmacotherapy and health care utilization and costs prior to total hip or total knee replacement in patients with osteoarthritis", *Arthritis & Rheumatism*, 2011

Publication

32

Asma Kheirollahi, Maryam Teimouri, Mehrdad Karimi, Nariman Moradi, Asie Sadeghi, akram vatannejad. "Evaluation of Lipid Ratios and Triglyceride-Glucose (TyG) Index as Risk Markers of Insulin Resistance in Iranian Polycystic Ovary Syndrome (PCOS) Woman", *Research Square*, 2020

Publication

33

Ki-Bum Won, Byoung Kwon Lee, Hyung-Bok Park, Ran Heo et al. "Quantitative assessment of coronary plaque volume change related to triglyceride glucose index: The Progression of Atherosclerotic Plaque Determined by Computed Tomographic Angiography Imaging (PARADIGM) registry", *Cardiovascular Diabetology*, 2020

Publication

34

Yue Zhang, Xiaosong Ding, Bing Hua, Qingbo Liu, Hui Gao, Hui Chen, Xue-Qiao Zhao, Weiping Li, Hongwei Li. "High Triglyceride-Glucose Index is Associated with Poor Cardiovascular Outcomes in Non-Diabetic ACS Patients with LDL-C Below 1.8mmol/L",

<% 1

<% 1

<% 1

35 journals.plos.org <% 1
Internet Source

36 jkms.org <% 1
Internet Source

37 journals.viamedica.pl <% 1
Internet Source

38 Pham Viet Thai, Hoang Anh Tien, Huynh Van Minh, Paul Valensi. "Triglyceride Glucose Index for the Detection of Asymptomatic Coronary Artery Stenosis in Patients With Type 2 Diabetes", Research Square, 2020
Publication

39 jdmdonline.biomedcentral.com <% 1
Internet Source

40 Trinovita Andraini, Sophie Yolanda. "Prevention of insulin resistance with *Hibiscus sabdariffa* Linn. extract in high-fructose fed rat", Medical Journal of Indonesia, 2015
Publication

41 heanoti.com <% 1
Internet Source

42 Alice Divina Melo de Brito, Helen Hermiana Miranda Hermsdorff, Mariana De Santis <% 1

Filgueiras, Sarah Aparecida Vieira-Ribeiro et al.
"TAG–glucose (TyG) index in childhood: an estimate of cut-off points and the relation to cardiometabolic risk in 4- to 9-year-old children",
Public Health Nutrition, 2020

Publication

43

Minghui Li, Aihua Zhan, Xiao Huang, Lihua Hu, Wei Zhou, Tao Wang, Lingjuan Zhu, Huihui Bao, Xiaoshu Cheng. "Positive association between triglyceride glucose index and arterial stiffness in hypertensive patients: the China H-type Hypertension Registry Study", Research Square, 2020

Publication

<% 1

44

"Posters", Diabetic Medicine, 2006

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES

< 5 WORDS