

Expression Of Interleukin-6 Levels In Elderly Sarcopenia

by Andi Makbu; Aman

FILE	BUL_EXPRESSION_OF_INTERLEUKIN-6_LEVELS_IN_ELDERLY_SARCOPENIA.PDF (292.71K)		
TIME SUBMITTED	10-FEB-2021 05:04AM (UTC+0700)	WORD COUNT	3972
SUBMISSION ID	1505750510	CHARACTER COUNT	20424

Expression Of Interleukin-6 Levels In Elderly Sarcopenia

Resliany Rachim¹, Agus Sudarso¹, Shanti Pricillia Makagiansar¹, Syakib Bakri¹,
Haerani Rasyid¹, A Makbul Aman¹, Hasyim Kasim¹, and Arifin Seweng²

¹Department of Internal Medicine, Medical Faculty, Hasanuddin University, Makassar, Indonesia

²Department of Biostatistics, Public health Faculty, Hasanuddin University, Makassar, Indonesia

Correspondence: Resliany Rachim, Department of Internal Medicine, Hasanuddin University's Faculty of Medicine, Perintis Kemerdekaan Km 10, Tamalanrea, Makassar, South Sulawesi, Indonesia.

Abstract: Background/Aim: Sarcopenia is a syndrome characterized by progressive loss of muscle mass and strength with a risk of adverse clinical outcomes such as physical disability, reduced quality of life, and death. Oxidative stress and inflammatory processes are known as factors that trigger sarcopenia by releasing the catabolic stimuli of IL-6. Based on this we study the dynamics of interleukin (IL-6) in elderly sarcopenia in Wahidin Sudirohusodo Hospital Makassar. **Method:** This study was an analytic cross-sectional design conducted at the Geriatric Polyclinic of Wahidin Sudirohusodo Hospital, Makassar in March-June 2020. The assessed variables were degree of sarcopenia (probable sarcopenia, sarcopenia, dan severe sarcopenia) based on The European Working Group on Sarcopenia in Older People 2 (EWGSOP2) criteria, age, sex, number of comorbids, and IL-6 levels examination. Statistical analysis was performed by descriptive statistical calculations and frequency distribution as well as the Independent-t statistical test, and Anova test. **Results:** Studied from 82 subjects (51 females, 62.2%) with the mean of 68.95 years of age. The prevalence of probable sarcopenia was 40.2%, sarcopenia 8.5%, severe sarcopenia 6.1%, and normal 45.1%. IL-6 levels were found to be significantly increased according to the severity of sarcopenia (normal, 52.81 ng/L; probable sarcopenia, 67.47 ug/L; sarcopenia, 135.36 ng/L; and severe sarcopenia, 287.99 ng/L). Based on age, IL-6 levels increased significantly with age (60-74 years, 63.28 ng/L; ≥75 years, 139.35 ng/L). Based on the number of comorbids, IL-6 levels increased significantly according to the number of comorbids (1 comorbids, 52.86 ng/L; ≥4 comorbids 120.84 ng/L). **Conclusion:** In the elderly subject, IL-6 levels increases according to the severity of sarcopenia. IL-6 levels also increases with age, and the number of comorbids.

Keywords: Sarcopenia, IL-6, Elderly.

1. INTRODUCTION

Population aging has taken place rapidly, especially in developing countries in the first decades of this century. In Indonesia, in 2010 the proportion of the population over 65 years of age was 5% and by 2035 it will increase to 10.6%. According to the population census in 2010, the number of elderly people (aged) in South Sulawesi was 5.4% and in 2035 it is estimated that it will be almost double as compared to 2010, which is 9.9%. The life

expectancy of the Indonesian population (men and women) has increased from 70.1 years in 2010-2015 period to 72.2 years in 2030-2035 period (Badan Pusat Statistik, 2013).

The increase in life expectancy and well-being of the population in line with the increase in the aging population in Indonesia has led to an increasing trend in various health problems (Badan Pusat Statistik, 2013). Changes that occur in aging are an irreversible and inevitable process characterized by changes in the morphology, functionality and biochemistry of the body, including the musculoskeletal system, and the most obvious changes include loss of muscle mass, and reduced muscle strength (Colón et al., 2018). Sarcopenia is a syndrome characterized by progressive loss of muscle mass and muscle strength with adverse clinical outcomes such as physical disability, decreased quality of life, and death. There are several mechanisms involved in the occurrence of sarcopenia including loss of alpha motor neuron axons, lack of physical activity, hormonal changes, impaired nutritional intake, oxidative stress, genetics, and inflammatory processes. There is evidence that pro-inflammatory mediators such as interleukin-6 (IL-6) interact with hormonal changes that occur in the aging process, thereby causing a decrease in physical activity (Aryana et al., 2018). Studies report that levels of pro-inflammatory cytokines such as IL-6 are elevated in elderly sarcopenia (Rong et al., 2018).

Based on the information above, this study aims to determine the expression of IL-6 levels in elderly sarcopenia. This research is still rarely conducted and published in Indonesia. The results of this study are expected to serve as future references for preventive and curative interventions in sarcopenia.

2. METHOD

2.1 Research Subject

This study was an analytic cross-sectional design conducted at the Geriatric Polyclinic of Wahidin Sudirohusodo Hospital, Makassar in March-June 2020. The assessed variables were degree of sarcopenia (probable sarcopenia, sarcopenia, dan severe sarcopenia) based on The European Working Group on Sarcopenia in Older People 2 (EWGSOP2) criteria, age, sex, number of comorbidities, and IL-6 levels examination. The elderly patient was explained about aims of the study and obtained consent.

2.2 Inclusion Criteria and Exclusion Criteria

The inclusion criteria include subjects >60 years old, willing to take part in the study, can walk alone or stand for 5 minutes without other assistive devices. The exclusion criteria include subjects with malignant disease, acute arthritis or acute infection, experience cognitive impairment.

2.3 Data Collection

Data collection was carried out by interview method, anthropometric measurement, handgrip (HG) strength dynamometer examination, muscle mass examination using Bioelectrical Impedance Analysis (BIA), physical performance examination with Time Up And Go test (TUG), and serum IL-6 examination.

2.4 Anthropometry

Anthropometric examination is carried out by measuring body weight (BW) and height (H) based on knee height. Measurement of height (cm) in the elderly using knee height, then converted through the calculation of the formula for Chumlea's knee height.

2.5 Sarcopenia Parameters

Based on the recommendations of EWGSOP2, the parameters of sarcopenia measured in this study were muscle mass, muscle strength, and physical performance. Measurement of muscle mass using the BIA tool by measuring the total skeletal muscle mass (muscle mass of the

upper leg, lower leg) divided by the square of height in meters and expressed in units of kg/m^2 . Examination of muscle strength was performed with an HG strength dynamometer, expressed in kilograms (kg), and physical performance checks were measured using the TUG instrument in seconds.

2.6 Diagnostic Criteria

The diagnostic criteria are based on the diagnostic criteria for sarcopenia established by the EWGSOP2. Muscle mass is said to be low if the value of BIA $<7.0 \text{ kg}/\text{m}^2$ for male subjects and the value of BIA $<5.5 \text{ kg}/\text{m}^2$ for female subjects. Muscle strength is said to be low if the value of HG strength dynamometer $<27 \text{ kg}$ for male subjects and the value of HG strength dynamometer $<16 \text{ kg}$ for female subjects. Physical performance was assessed by the TUG instrument, and said to be low if it was 20 seconds.

2.7 IL-6 Levels

Measurement of levels or concentrations of IL-6 with the ELISA test. IL-6 levels are checked from blood serum. About 3 mL of the subject's venous blood was drawn by the research assistant. Blood in serum was centrifuged at 2000-3000 RPM for 20 minutes and stored at -80°C . IL-6 examination using quantikine HS human IL-6 immunoassay (catalog list no E0090Hu) was measured by the ELISA method in ng/L units. This measurement is carried out once without repetition. The sensitivity of the detected concentration was 1.03 ng/L, and the cut-off value in normal people ranged up to 7 ng/L.

2.8 Statistic Analysis

Statistical analysis was carried out electronically using the SPSS version 22 device. The statistical analysis carried out was descriptive statistical calculations and frequency distribution as well as the Independent-t statistical test, and Anova test. The result of the test is statistically significant if the p value <0.05 .

2.9 Ethical License

Ethical permission has been obtained from the Health Research Ethics Committee of the Hasanuddin University Medical Faculty / Wahidin Sudirohusodo Hospital with letter number: LB.02.01/2.2/4071/2020.

3. RESULTS

3.1 Characteristics Research Subjects

Data analysis was carried out on 82 research subjects, obtained 31 male (37.8%) and 51 female (62.2%). Mean age was 68.95 (6.34) years. The number of subjects in the age range of 60-74 years was 76.8%, and aged ≥ 75 years was 23.2%. Based on the degree of sarcopenia, 33 people (40.2%) had probable sarcopenia, 7 (8.5%) of sarcopenia, 5 (6.1%) of severe sarcopenia, and 37 (45.1%) normal). Based on parameters of sarcopenia, 12 people (14.6%) had low muscle mass, 44 people (53.7%) low muscle strength, and 30 (36.6%) low physical performance. The number of comorbid obtained in subjects with a number of 1-3 comorbid was 52 people (63.4%), and with a total of ≥ 4 comorbid as many as 30 people (36.6%). The mean value of IL-6 obtained in this study was 80.9 (62.7) as seen in Table 1.

Table 1. Characteristics of Research Subjects (n=82)

Variables		n	%
Sex	Male	31	37.8
	Female	51	62.2
Age	60-74 years	63	76.8
	≥ 75 years	19	23.2

Degree of Sarcopenia	Normal	37	45.1
	Probable Sarcopenia	33	40.2
	Sarcopenia	7	8.5
	Severe Sarcopenia	5	6.1
Muscle Mass	Normal	70	85.4
	Low	12	14.6
Muscle Strength	Normal	38	46.3
	Low	44	53.7
Physical Performance	Normal	52	63.4
	Low	30	36.6
Number of Comorbids	1-3	52	63.4
	≥ 4	30	36.6
Interleukin-6 Levels		80.9 (62,7)	

3.2 Description of Interleukin-6 Levels based on Sex, Age, Degree of Sarcopenia, and Number of Comorbids

Interleukin-6 levels were found to be higher in women (84.04) than in men (75.74), but the statistical test results showed that the difference was not significant ($p>0.05$) as shown in Table 2.

Interleukin-6 levels were found to be higher at age ≥ 75 years (139.35) than at age 60-74 years (63.28) and there was a visible pattern of increasing IL-6 levels with increasing age. The results of statistical tests showed that this difference was significant ($p<0.01$) as shown in Table 2.

Levels of IL-6 were found to be the highest in severe sarcopenia (287.99) and lowest in normal (52.81) and this result was statistically significant ($p<0.001$) as shown in Table 2.

Table 2. Levels of Interleukin-6 based on Sex, Age, Degree of Sarcopenia, and Number of Comorbids

Variables	n	Mean	Standard Deviation	p
Sex				
Male	31	75.74	52.16	0.565
Female	51	84.04	68.70	
Age				
60-74 years	63	63.28	29.39	0.004
≥ 75 years	19	139.35	100.24	
Degree of Sarcopenia				
Normal	37	52.81	9.81	<0.001
Probable sarcopenia	33	67.47	29.11	
Sarcopenia	7	135.36	42.15	
Severe Sarcopenia	5	287.99	47.37	
Number of Comorbids				
1-3	52	57.86	15.85	0.001

≥4	30	120.84	89.18	

Interleukin-6 levels were found to be significantly higher in the comorbid count ≥ 4 (120.84) compared to the comorbid counts 1-3 (57.86) and this result was statistically significant ($p < 0.01$) as shown in Table 2.

4. DISCUSSION

This study included 82 research subjects with female (62.2%) and male (37.8%) subjects. The study of Riviati et al. found 60.2% women and 39.8% men (Riviati et al., 2017).

The age of the research subjects is 60 years and over. The mean age was 68.95 (6.34) years. The number of subjects in the 60-74 years age range was 6.8%, and at ≥ 75 years old was 23.2%. Research from Aryana et al. obtained a mean age of 69.52 ± 8.71 years. (Aryana et al., 2018). Meanwhile, the mean age retrieved from the research subjects by Patel et al. was 67 ± 2.6 years (Patel et al., 2013).

The degree of sarcopenia obtained in this study was found to have probable sarcopenia by 40.2%, sarcopenia 8.5%, severe sarcopenia 6.1%, and the remaining 45.1% were normal. In the study of Sobestiansky et al. of the 287 research subjects, it was found that probable sarcopenia was 73% and sarcopenia was 20% (Sobestiansky et al., 2019). Meanwhile, in the study of Pang et al. from 536 study subjects, it was found that probable sarcopenia was 15.3%, sarcopenia was 15.5%, and severe sarcopenia was 10.1% (Pang et al., 2020). From the results of this study, there was quite a lot of probable sarcopenia (40.2%), and this suggests that it turns out that quite a lot of subjects are at risk of sarcopenia, therefore, it is necessary to immediately conduct an assessment to find the causative factors and initiate effective early interventions, so that the incidence of sarcopenia can be prevented. This is consistent with the algorithm from EWGSOP2 for the earliest possible diagnosis and intervention (Cruz-Jentoft et al., 2019).

Of the 82 research subjects, 70 people (85.4%), subjects with normal muscle mass and 12 (14.6%), subjects with low muscle mass. Research by Bian et al. of the 441 study subjects, 82% of subjects with normal muscle mass and 18% of subjects with low muscle mass (Bian et al., 2017). As many as 38 people (46.3%), subjects with normal muscle strength and 44 people (53.7%), subjects with low muscle strength (36). The research of Riviati et al. at the age of more than 75 years, there were 56.5% subjects with low muscle strength and 43.5% subjects with normal muscle strength (Riviati et al., 2017). A total of 52 people (63.4%), subjects with normal physical performance and 30 people (36.6%), subjects with low physical performance. Research by Guedes et al. was found that subjects with low physical performance were 34.3%, while subjects with normal physical performance were 65.7% (Guedes et al., 2019).

5 total of 52 people (63.4%), subjects had a comorbid number of 1-3. From the research of Bahat et al., The average number of comorbid was 2,6 (Bahat et al., 2013). Research by Aryana et al. was done by excluding comorbid or done without any comorbid. This is because this study used a clinical-based setting where the subject came to the polyclinic with various comorbid diseases, while the research of Aryana et al. was community-based (Aryana et al., 2018). The most comorbid diseases in this study were hypertension (85.4%) and Diabetes Mellitus (70.7%). The research of Riviati et al. obtained comorbid hypertension (78.7%) and diabetes mellitus (44.3%) (Riviati et al., 2017). In this study the 27 was no analysis of the subjects with comorbid diseases. In the aging process, there is an increase in

8-hydroxy-2-deoxyguanosine (8-OH-dG) which is a free radical marker. It is also found in hypertension and diabetes mellitus (Nemtsova et al., 2018).

The mean IL-6 level in this study was 80.9 (62.7). Research by Bian et al., the mean IL-6 level in the group with decreased muscle mass was 49.77 (22.14) and in the group with no decreased muscle mass was 39.72 (29.53) (Bian et al., 2017). The increase in IL-6 levels was influenced by various factors, such as age, gender, comorbid diseases (Maggio et al., 2006).

Description of Interleukin-6 Levels by Sex

In this study, it was found that IL-6 levels were higher in female than male, but this difference was not statistically significant ($p > 0.05$). Research by Aureli et al. found that the IL-6 levels were significantly higher in women than in men (Aureli et al., 2014). In this study, it was found that there was no significant relationship between sex and IL-6 levels. These results differ from some literature whereby sex association with IL-6 levels suggests that decreased production and levels of circulating sex hormones cause a mild pro-inflammatory state in the elderly (Maggio et al., 2006). One in-vitro study found that estrogen increases the synthesis of pro-inflammatory cytokines in cells that resemble macrophages (O'Connor et al., 2007).

Description of Interleukin-6 levels based on Age

In this study, it was found that IL-6 levels were higher at ≥ 75 years of age and lower at the age of 60-74 years and there was a statistically significant pattern of increasing IL-6 levels with increasing age ($p < 0.01$). In the study by Kuznicka et al. it was also found that there are significantly higher IL-6 levels at older age, and this occurred in the entire study group ($p < 0.001$) (Kuźniak et al., 2016). Age-related increase in IL-6 levels were due to overproduction of oxygen-free radicals, which stimulate IL-6 production (Maggio et al., 2006).

Description of Interleukin-6 levels based on Degree of sarcopenia

In this study, it was found statistically significant ($p < 0.001$) that IL-6 levels increased according to the severity of sarcopenia. So the more severe the degree of sarcopenia, the higher the IL-6 levels. Based on the IWG/SOP2 definition, probable sarcopenia is a decrease in muscle strength, sarcopenia is a decrease in muscle strength and a decrease in muscle mass, and severe sarcopenia is a decrease in muscle strength, a decrease in muscle mass and a decrease in physical performance. High levels of IL-6 result in decreased muscle mass, muscle strength and physical performance (Aryana et al., 2018; Visser et al., 2002; Grosicki et al., 2020).

Description of Interleukin-6 levels based on Number of Comorbidities

In this study, it was found statistically significant ($p < 0.01$) that IL-6 levels increased according with the number of comorbidities. The physiological role of IL-6 has been mostly studied in the context of the acute phase response, although there is growing evidence that IL-6 plays a central role in the pathogenesis of chronic disease. In chronic inflammation, IL-6 has a detrimental role that favours mononuclear cell accumulation at the site of injury (Maggio et al., 2006).

The inflammatory reaction is only one of the factors causing sarcopenia. Other factors include nutritional problems and decreased hormone levels in the body. Sarcopenia will be one of the main causes of disability and decreased quality of life in the elderly. In recent years, although many experts have explored this in various aspects, the pathological mechanism remains unclear and there are no standard diagnostic criteria. To determine the effective method to prevent and treat this disease, it is necessary to conduct a more large-scale clinical study (Bian et al., 2017). Another limitation of our study is the clinical-based setting where patients who come to the polyclinic with various comorbidities, and hope that

further community-based research can be carried out within healthy individuals in the future with few or no comorbidities so as not to bias IL-6 levels.

5. CONCLUSION

In the elderly subject, IL-6 levels increases according to the severity of sarcopenia. IL-6 levels also increases with age, and the number of comorbidities.

Source of Fundling

This study used private funds.

Ethical Clearance

Ethical permission has been obtained from the Health Research Ethics Committee of the Hasanuddin University Medical Faculty / Wahidin Sudirohusodo Hospital with letter number: LB.02.01/2.2/4071/2020.

Competing Interests Statement

No potential conflict of interest relevant to be declared.

6. REFERENCES

- [1] Aryana S, Lestari AW, Putrawan IB, Purnami NKR, Astika IN, Kuswardhani RAT. The relationship between IL-6 and CRP with Sarcopenia in indigenous elderly population at Pedawa Village, Buleleng, Bali, Indonesia. *Heal Sci J Indones*. 2018;9(1):37–44.
- [2] Aureli A, Sebastiani P, Del Beato T, Marimpietri AE, Graziani A, Sechi E, et al. Involvement of IL-6 and IL-1 receptor antagonist on intellectual disability. *Immunol Lett*. 2014;162(1):124–31.
- [3] Badan Pusat Statistik Indonesia. Proyeksi Penduduk Indonesia. Indonesia Population Projection 2010-2035. Badan Pusat Statistik Indonesia. 2013. 978–979 p.
- [4] Bahat G, Tufan F, Bahat Z, Aydin Y, Tufan A, Akpinar TS, et al. Assessments of functional status, comorbidities, polypharmacy, nutritional status and sarcopenia in Turkish community-dwelling male elderly. *Aging Male*. 2013;16(2):67–72.
- [5] Bian AL, Hu HY, Rong YD, Wang J, Wang JX, Zhou XZ. A study on relationship between elderly sarcopenia and inflammatory factors IL-6 and TNF- α . *Eur J Med Res*. 2017;22(1):4–11.
- [6] Colón CJP, Molina-Vicenty IL, Frontera-Rodríguez M, García-Ferré A, Rivera BP, Cintrón-Vélez G, et al. Muscle and Bone Mass Loss in the Elderly Population: Advances in diagnosis and treatment. *J Biomed [Internet]*. 2018;3(12):40–9. Available from: <http://www.jbiomed.com/v03p0040.htm>
- [7] Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: Revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48(1):16–31.
- [8] Grosicki GJ, Barrett BB, Englund DA, Liu C, Travison TG, Cederholm T, et al. Circulating Interleukin-6 Is Associated with Skeletal Muscle Strength, Quality, and Functional Adaptation with Exercise Training in Mobility-Limited Older Adults. *J frailty aging*. 2020;9(1):57–63.
- [9] Guedes R de C, Dias RC, Neri AL, Ferriolli E, Lourenço RA, Lustosa LP. Declínio da velocidade da marcha e desfechos de saúde em idosos: dados da Rede Fibra. *Fisioter e Pesqui*. 2019;26(3):304–10.
- [10] Maggio, Marcello, Guralnik Jack M, Longo L FL. Interleukin-6 in Aging and Chronic

- Disease: A Magnificent Pathway. *J Gerontol A Biol Sci Med Sci* [Internet]. 2006;61(6):575–84. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf>
- [11] Nemtsova V, Bilovol O, Ilchenko I, Shalimova A. Age-associated features of oxidative stress as marker of vascular aging in comorbid course of hypertension and type 2 diabetes mellitus. *Vessel Plus*. 2018;2(9):27.
- [12] O'Connor MF, Motivala SJ, Valladares EM, Olmstead R, Irwin MR. Sex differences in monocyte expression of IL-6: Role of autonomic mechanisms. *Am J Physiol - Regul Integr Comp Physiol*. 2007;293(1):145–51.
- [13] Pang BWJ, Wee SL, Lau LK, Jabbar KA, Seah WT, Ng DHM, et al. Prevalence and Associated Factors of Sarcopenia in Singaporean Adults—The Yishun Study. *J Am Med Dir Assoc*. 2020.
- [14] Patel HP, Syddall HE, Jameson K, Robinson S, Denison H, Roberts HC, et al. Prevalence of sarcopenia in community-dwelling older people in the UK using the European Working Group on Sarcopenia in Older People (EWGSOP) definition: Findings from the Hertfordshire Cohort Study (HCS). *Age Ageing*. 2013;42(3):378–84.
- [15] Puzianowska-Kuźnicka M, Owczarż M, Wiczorowska-Tobis K, Nadrowski P, Chudek J, Slusarczyk P, et al. Interleukin-6 and C-reactive protein, successful aging, and mortality: The PolSenior study. *Immun Ageing*. 2016;13(1):1–12.
- [16] Riviaty N, Setiati S, Laksmi PW, Abdullah M. Factors Related with Handgrip Strength in Elderly Patients. *Acta Med Indones*. 2017;49(3):215–9.
- [17] Rong YD, Bian AL, Hu HY, Ma Y, Zhou XZ. Study on relationship between elderly sarcopenia and inflammatory cytokine IL-6, anti-inflammatory cytokine IL-10. *BMC Geriatr*. 2018;18(1):1–6.
- [18] Sobestiansky S, Michaelsson K, Cederholm T. Sarcopenia prevalence and associations with mortality and hospitalisation by various sarcopenia definitions in 85-89 year old community-dwelling men: A report from the ULSAM study. *BMC Geriatr*. 2019;19(1):1–13.
- [19] Visser M, Pahor M, Taaffe DR, Goodpaster BH, Simonsick EM, Newman AB, et al. Relationship of interleukin-6 and tumor necrosis factor- α with muscle mass and muscle strength in elderly men and women: The health ABC study. *Journals Gerontol - Ser A Biol Sci Med Sci*. 2002;57(5):326–32.

Expression Of Interleukin-6 Levels In Elderly Sarcopenia

ORIGINALITY REPORT

%**23**

SIMILARITY INDEX

%**14**

INTERNET SOURCES

%**19**

PUBLICATIONS

%**6**

STUDENT PAPERS

PRIMARY SOURCES

- 1** academic.oup.com Internet Source % **1**
- 2** I Gusti Putu Suka Aryana, I Nyoman Astika, Raden Ayu Tuty Kuswardhani, Ida Bagus Putu Putrawan et al. "High Myostatin Serum Related with High Prevalence of Sarcopenia Among Elderly Population in Pedawa Village, Bali, Indonesia", The Indonesian Biomedical Journal, 2019 Publication % **1**
- 3** ejournal2.litbang.kemkes.go.id Internet Source % **1**
- 4** "Handbook of Immunosenescence", Springer Science and Business Media LLC, 2019 Publication % **1**
- 5** www.tandfonline.com Internet Source % **1**
- 6** Gimena dos Santos, Mikhail A. Kutuzov, Karen M. Ridge. "The inflammasome in lung diseases", American Journal of Physiology-Lung Cellular % **1**

7	wjso.biomedcentral.com Internet Source	% 1
8	www.science.gov Internet Source	% 1
9	Submitted to University of Oklahoma Student Paper	% 1
10	eprint.ncl.ac.uk Internet Source	% 1
11	Submitted to Louisiana Tech University Student Paper	% 1
12	pericles.pericles-prod.literatumonline.com Internet Source	% 1
13	Submitted to Udayana University Student Paper	% 1
14	www.id-press.eu Internet Source	% 1
15	F Aloisi. "Peripheral tissue release of interleukin-6 in patients with chronic kidney diseases: Effects of end-stage renal disease and microinflammatory state", <i>Kidney International</i> , 07/2006 Publication	<% 1

16

www.pubmedcentral.nih.gov

Internet Source

<% 1

17

S. P. Marso. "Increase in interleukin-6 following arterial injury is related to insulin resistance, the ?174G?C polymorphism and complex plaque morphology", International Journal of Immunogenetics, 10/2006

Publication

<% 1

18

Mary-Frances O'Connor, Sarosh J. Motivala, Edwin M. Valladares, Richard Olmstead, Michael R. Irwin. "Sex differences in monocyte expression of IL-6: role of autonomic mechanisms", American Journal of Physiology-Regulatory, Integrative and Comparative Physiology, 2007

Publication

<% 1

19

Young Koog Cheon. "Diagnostic Utility of Interleukin-6 (IL-6) for Primary Bile Duct Cancer and Changes in Serum IL-6 Levels Following Photodynamic Therapy", The American Journal of Gastroenterology, 7/7/2007

Publication

<% 1

20

dinajmc.org

Internet Source

<% 1

21

Submitted to Leeds Beckett University

Student Paper

<% 1

22 Submitted to State Islamic University of Alauddin Makassar
Student Paper <% 1

23 bmcgeriatr.biomedcentral.com
Internet Source <% 1

24 joacp.org
Internet Source <% 1

25 Submitted to University of Brighton
Student Paper <% 1

26 www.dovepress.com
Internet Source <% 1

27 www.intechopen.com
Internet Source <% 1

28 journals.lww.com
Internet Source <% 1

29 Stevent Sumantri, Iris Rengganis, Purwita Wijaya Laksmi, Rudy Hidayat, Sukanto Koesnoe, Hamzah Shatri. "The impact of low muscle function on health-related quality of life in Indonesian women with systemic lupus erythematosus", *Lupus*, 2021
Publication <% 1

30 "Abstracts of the 16th International E-Congress of the European Geriatric Medicine Society", *European Geriatric Medicine*, 2020 <% 1

31

Lan Liu, Dian-Yuan Lu, Jian-Rong Cai, Li Zhang. "The value of oral contrast ultrasonography in the diagnosis of gastric cancer in elderly patients", *World Journal of Surgical Oncology*, 2018

Publication

<% 1

32

Helene Payette. "Insulin-Like Growth Factor-1 and Interleukin 6 Predict Sarcopenia in Very Old Community-Living Men and Women: The Framingham Heart Study", *Journal of the American Geriatrics Society*, 9/2003

Publication

<% 1

33

nutritionandmetabolism.biomedcentral.com

Internet Source

<% 1

34

Annika Loftenius Gunilla Sandborgh-. "ACUTE EXPOSURE TO MERCURY FROM AMALGAM: NO SHORT-TIME EFFECT ON THE PERIPHERAL BLOOD LYMPHOCYTES IN HEALTHY INDIVIDUALS", *Journal of Toxicology and Environmental Health, Part A*, 1998

Publication

<% 1

35

Ömer Elma, Sevilay Tumkaya Yilmaz, Tom Deliens, Peter Clarys, Jo Nijs, Iris Coppieters, Andrea Polli, Anneleen Malfliet. "Chronic Musculoskeletal Pain and Nutrition: Where Are

<% 1

We and Where Are We Heading To?", PM&R, 2020

Publication

36

Sema Uçak, Rıdvan Sivritepe, Oğuzhan Kara, Ecem Sevim et al. "Association between sarcopenia and erectile dysfunction in males with type II diabetes mellitus", The Aging Male, 2018

Publication

<% 1

37

ircmj.com

Internet Source

<% 1

38

onlinelibrary.wiley.com

Internet Source

<% 1

39

Isela Carrera, Hector A. Moreno, Sergio Sierra, Alexandre Campos, Marcela Munera, Carlos A. Cifuentes. "Chapter 4 Technologies for Therapy and Assistance of Lower Limb Disabilities: Sit to Stand and Walking", Springer Science and Business Media LLC, 2020

Publication

<% 1

40

"Muscle Atrophy", Springer Science and Business Media LLC, 2018

Publication

<% 1

41

David Barry. "Sarcopenia", Elsevier BV, 2020

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5
WORDS