

Yuyun_Contrast_Induced_Nephropathy.pdf

by Uleng Bahrun

FILE	YUYUN_CONTRAST_INDUCED_NEPHROPHATY.PDF (155.08K)		
TIME SUBMITTED	19-DEC-2019 04:52AM (UTC+0700)	WORD COUNT	2052
SUBMISSION ID	1236725195	CHARACTER COUNT	11966



14

**Analysis of Risk Factors Contrast Induced Nephropathy
in Coronary Artery Disease Patients Obtained
Percutaneous Coronary Intervention at Wahidin
Sudirohusodo Hospital in Makassar, South Sulawesi**

Yuyun Widaningsih^{a*}, Rosdiana Natsir^b, Mansyur Arif^c, Uleung Bahrund^d,
Hasyim Kasim^e, Fitriani Mangerangi^f, Nursin Abdul Kadir^g, Rini Anggraeni^h,
Nadyah Harunaⁱ, Mochammad Hatta^j, Andi Nilawati Usman^k

^{a,c,d,f,g}Department of Clinical Pathology, Hasanuddin University, Makassar, Indonesia

^bDepartment of Biochemistry, Hasanuddin University, Makassar, Indonesia

^eInternal Medicine Department, Hasanuddin University, Makassar, Indonesia

^hDepartment of Hospital Management, Hasanuddin University, Makassar, Indonesia

ⁱMedicine Faculty, UIN Alauddin University, Indonesia

^jDepartment of Immunology and Biomolecular, Hasanuddin University, Makassar, Indonesia

^kHalal Center, Public Health Hasanuddin University

^aEmail: yuyun.widaningsih@yahoo.com

Abstract

Incidence of Contrast Induced Nephropathy (CIN) after percutaneous coronary intervention (PCI) is a determinant factor of Acute Kidney Injury (AKI). Risk factors evidence can strengthen the prevention of CIN before PCI intervention performed. This study aimed to identify risk factors for the incidence of CIN in Coronary Artery Disease (CAD) patients obtained PCI. Evidence of the risk factors can strengthen the prevention of CIN before PCI intervention performed. This study aimed to identify risk factors for the incidence of CIN in patients with CAD obtained PCI.

* Corresponding author.

It was a quantitative research with retrospective design, population were coronary artery disease (CAD) patients and sample size were 25 patients. Bivariate analysis using Fisher exact statistical test showed that variable associated significantly with the incidence of CIN only diabetes mellitus (OR = 21.000, P = 0.007, 95% CI = 1922-299392) while the variable gender, age, smoking, acute myocardial infarction and hypertension did not have a significant relationship ($p > 0.05$). Predictor for incidence CIN based on binary regression analysis was diabetes mellitus (PR = 21.000, P = 0.007, 95% CI = 1922-299392) while variable gender, age, smoking, acute myocardial infarction and hypertension did not have a significant relationship ($p > 0.05$). Result of this study reinforces the need prevention for patients with risk factors before PCI intervention especially diabetes mellitus.

Keywords: Contrast Induced Nephropathy; Percutaneous Coronary Intervention; Diabetes Mellitus.

1. Introduction

Administration of contrast agents in coronary artery disease patients when percutaneous coronary intervention (PCI) is a risk factor for the occurrence of contrast induced nephropathy (CIN), patient characteristics such as age and history of disease is also risk factor for the occurrence of CIN [1-3]. Incidence of CIN may develop to acute kidney injury and was significantly associated with morbidity of hospitalized patients [4].

Some studies revealed that CIN incidence in women increased morbidity and mortality. Age, gender and incidence of diabetes mellitus are predictor of the incidence of CIN after PCI [5-7]. The aging and cardiovascular disease itself is also a determinant of CIN [8]. Diabetes mellitus is a predisposing factor for CIN via several mechanisms such as an increase in the tubules transport activity and formation of reactive oxygen species [9]. Increased Risk of CIN and bad prognostic in patients with diabetes mellitus type 2 Become a consideration for preparing prevention before administration of contrast agents [10, 11].

Knowledge of risk factors can be useful for the prevention and treatment of CIN incidence. Provision of contrast media should be postponed in patients with congestive heart failure until haemodynamic status can be resolved and delayed 24 hours for patients with myocardial infarction. Important consideration also for patients who have risk factors to be considered in getting hypo-osmolar contrast media [12].

Research about CIN in Makassar, South Sulawesi is infrequent, so this study aims to identify risk factors for the incidence of CIN in patients with cardiovascular obtained PCI in wahidinsudirohusodo hospital in Makassar, South Sulawesi.

2. Material and Methods

This study was retrospective that investigated variables suspected could be a predictor for cardiovascular CIN in patients undergoing PCI at Wahidin Sudirohusodo hospital, Makassar, South Sulawesi. Risk factor variables was obtained from medical records of patients. The population were all Coronary Artery Disease (CAD) patients which assigned PCI, sample size were 25 patients obtained by accidental sampling.

Serum creatinine patients were examined 24-48 hours after PCI then categorized into two groups; patients

experienced CIN and patients did not experience CIN. This criterion was based on increasing of serum creatinine > 0.5 mg /dl or 25% improvement from baseline in 24-48 hours after PCI. It was according to Kidney Disease Improving Global Outcomes (KDIGO) of the year 2012.

Using bivariate statistical analysis with Fisher Exact test followed by a multivariate analysis, binary logistic regression. Data then displayed with table contained frequency and percent along with odds ratio (OR) value, probability value(p) and confidence interval (CI) of 95%, declared significant if the p value ≤0.05.

3. Result

Bivariate analysis showed variable associated significantly with incidence of CIN only diabetes mellitus (PR = 21.000, P = 0.007, 95% CI = 1.922-229.392) while variables gender, age, smoking, acute myocardial infarction and hypertension did not have a significant relationship (p≥0.05) (Table 1). Variables met the criteria (p <0.25), namely age, smoking habits and history of diabetes mellitus included in multivariate analysis (Table 2).

Table 1: Risk of Contrast Induced Nephropathy

		Incidence of Nephropathy		Contrast Induced		OR	P	95% CI
		Yes (n=10)	No (n=15)	Yes	No			
		n	%	n	%			
Gender	Female	1	4.00	1	4.00	1.556	0.650*	0.086-28.145
	Male	9	36.00	14	56.00			
Age (year)	<60	5	20.00	13	52.00	6.500	0.075*	0.937-45.106
	≥60	5	20.00	2	8.00			
Smoking	Yes	3	12.00	2	8.00	0.359	0.248*	0.048-2.689
	No	7	28.00	13	52.00			
Myocard Infarction Acute	Yes	2	8.00	6	24.00	0.583	0.402*	0.058-2.414
	No	8	32.00	9	32.00			
Congestive Heart Failure	Yes	2	8.00	2	8.00	1.625	0.532*	0.190-13.933
	No	8	32.00	13	52.00			
Diabetes Mellitus	Yes	6	24.00	1	4.00	21.000	0.007*	1.922-229.392
	No	4	16.00	14	56.00			
Hypertension	Yes	8	32.00	12	48.00	1.000	0.687*	0.135-7.932
	No	3	8.00	2	12.00			

* Fisher Exact Test

n=number, OR=Odd Ratio, P=probability, CI=confidence interval

Binary logistic regression analysis in table 2 showed that variable was being significant predictors for incidence of CIN after PCI was history of diabetes mellitus (OR = 4.199, P = 0.040, 95% CI=1.125-202.026) while age

and smoking habits were not predictor of CIN incidence ($p > 0.05$).

Table 2: Multivariate Regression Analysis

Variable	OR	P	95%CI
Age	1.184	0.277	0.361-35.106
Smoking	0.002	0.969	0.053-16.848
Diabetes Mellitus	4.199	0.040 ^a	1.125-202.026 ^a

OR=Odd Ratio,P=probability, CI=confidence interval

^asignificant ($p \leq 0.05$)

4. Discussion

Study results revealed that variables associated significantly and be a predictor for incidence of CIN only diabetes mellitus disease. Patients who had insulin resistance and diabetes mellitus were more susceptible to CIN compared with non-diabetic patients both PCI and non PCI, [13, 14]. Kidneys of diabetic patients are particularly vulnerable to hypoxia and oxidative stress after administration of contrast agent, hypoxia induced formation of reactive oxygen species (ROS) and it has an important role in the pathogenesis of CIN [9]. ROS could lead to injury of vascular endothelial tubules directly and as a trigger of renal parenchymal hypoxia induced endothelial dysfunction and dysregulation of tubular transport [15]. Giving contrast media (CM) increase production of ROS and renal oxidative stress which mediates damage cell membrane and induced apoptosis and necrosis [16]. Administration low or iso-osmolar contrast media, hydration pre- and post-contrast hydration with 1500 ml of 0.9% saline solution infusion (N-acetylcysteine (NAC) associated with sodium bicarbonate (Na₂HCO₃) infusion) could be alternative for patients with history of diabetes mellitus or renal dysfunction [17-21].

This study was consistent with the results of global finding that diabetes mellitus was predictor for incidence of CIN. It provided recommendations for medical services, especially in Makassar, South Sulawesi, Indonesia that patients with a history of diabetes mellitus would need precaution before PCI. Further research should explore predictor variables for CIN with larger sample size and more variable in order to provide prevention recommendations for patients undergoing PCI.

5. Conclusion

History of diabetes mellitus was predictor variable for patients with coronary artery disease (CAD) undergoing PCI, it reinforces requirement for prevention in patients with risk factors before PCI intervention.

Acknowledgement

The authors would like to acknowledge to my husband, Amir Ilyas for his support during this study.

Competing Interest

The authors declare that they have no competing interests.

References

- [1]. Nough, H., et al., Incidence and Main Determinants of Contrast-Induced Nephropathy following Coronary Angiography or Subsequent Balloon Angioplasty. *Cardiorenal Medicine*, 2013. **3**(2): p. 128-135.
- [2]. McCullough, P.A. and K.R. Sandberg, Epidemiology of contrast-induced nephropathy. *Rev Cardiovasc Med*, 2003. **4 Suppl 5**: p. S3-9.
- [3]. Erley, C., [Iodinated contrast agent-induced nephropathy]. *Radiologe*, 2007. **47**(9): p. 761-7.
- [4]. Tumlin, J., et al., Pathophysiology of contrast-induced nephropathy. *Am J Cardiol*, 2006. **98**(6a): p. 14k-20k.
- [5]. Lucreziotti, S., et al., Female gender and contrast-induced nephropathy in primary percutaneous intervention for ST-segment elevation myocardial infarction. *Int J Cardiol*, 2014. **174**(1): p. 37-42.
- [6]. Iakovou, I., et al., Impact of gender on the incidence and outcome of contrast-induced nephropathy after percutaneous coronary intervention. *J Invasive Cardiol*, 2003. **15**(1): p. 18-22.
- [7]. Chong, E., et al., Risk factors and clinical outcomes for contrast-induced nephropathy after percutaneous coronary intervention in patients with normal serum creatinine. *Ann Acad Med Singapore*, 2010. **39**(5): p. 374-80.
- [8]. Zaraca, F., C.J. Wiedermann, and H. Ebner, Contrast media-induced nephropathy in patients undergoing angiography prior to or during vascular surgery: a systematic review. *Minerva Chir*, 2011. **66**(6): p. 553-60.
- [9]. Heyman, S.N., et al., Why is diabetes mellitus a risk factor for contrast-induced nephropathy? *Biomed Res Int*, 2013. **2013**: p. 123589.
- [10]. Zaytseva, N.V., et al., Contrast-induced nephropathy in patients with type 2 diabetes during coronary angiography: risk-factors and prognostic value. *Diabetes Res Clin Pract*, 2009. **86 Suppl 1**: p. S63-9.
- [11]. Shamkhalova, M., et al., [Contrast-induced nephropathy in coronarography of patients with type 2 diabetes mellitus: risk factors, prognostic significance, prophylactic approaches]. *Ter Arkh*, 2009. **81**(8): p. 36-42.
- [12]. Goldenberg, I. and S. Matetzky, Nephropathy induced by contrast media: pathogenesis, risk factors and preventive strategies. *Cmaj*, 2005. **172**(11): p. 1461-71.
- [13]. Rahman, M.M., et al., Contrast induced nephropathy in diabetic and non-diabetic patients during coronary angiogram and angioplasty. *Mymensingh Med J*, 2010. **19**(3): p. 372-6.
- [14]. Li, Y., et al., Insulin Resistance Increases the Risk of Contrast-Induced Nephropathy in Patients Undergoing Elective Coronary Intervention. *Angiology*, 2016. **67**(2): p. 139-45.
- [15]. Pisani, A., et al., Role of reactive oxygen species in pathogenesis of radiocontrast-induced nephropathy. *Biomed Res Int*, 2013. **2013**: p. 868321.
- [16]. Cetin, M., et al., Ionic high-osmolar contrast medium causes oxidant stress in kidney tissue: partial protective role of ascorbic acid. *Ren Fail*, 2008. **30**(5): p. 567-72.

- [17]. Gupta, R.K. and T.J. Bang, Prevention of Contrast-Induced Nephropathy (CIN) in Interventional Radiology Practice. *Semin Intervent Radiol*, 2010. **27**(4): p. 348-59.
- [18]. Alessandri, N., et al., Prevention of acute renal failure post-contrast imaging in cardiology: a randomized study. *Eur Rev Med Pharmacol Sci*, 2013. **17 Suppl 1**: p. 13-21.
- [19]. Lee, S.W., et al., Preventive strategies of renal insufficiency in patients with diabetes undergoing intervention or arteriography (the PREVENT Trial). *Am J Cardiol*, 2011. **107**(10): p. 1447-52.
- [20]. Subramaniam, R.M., et al., AHRQ Comparative Effectiveness Reviews, in Contrast-Induced Nephropathy: Comparative Effectiveness of Preventive Measures. 2016, Agency for Healthcare Research and Quality (US): Rockville (MD).
- [21]. Owen, R.J., et al., Canadian Association of Radiologists consensus guidelines for the prevention of contrast-induced nephropathy: update 2012. *Can Assoc Radiol J*, 2014. **65**(2): p. 96-105.

ORIGINALITY REPORT

% **16**
SIMILARITY INDEX

% **7**
INTERNET SOURCES

% **14**
PUBLICATIONS

% **9**
STUDENT PAPERS

PRIMARY SOURCES

- 1** "The challenge of subsea tunnelling", International Journal of Rock Mechanics and Mining Sciences & Geomechanics Abstracts, 1994
Publication % **1**
- 2** Yoshitaka Isaka, Hiromitsu Hayashi, Kazutaka Aonuma, Masaru Horio et al. "Guideline on the Use of Iodinated Contrast Media in Patients With Kidney Disease 2018", Circulation Journal, 2019
Publication % **1**
- 3** www.europeanreview.org
Internet Source % **1**
- 4** www.cmaj.ca
Internet Source % **1**
- 5** agroaid-bd.org
Internet Source % **1**
- 6** Vahideh Yavari, Mohammad Ali Ostovan, Javad Kojuri, Raha Afshariani, Alireza Hamidian, Jamshid Roozbeh, Maryam Pakfetrat. "The

preventive effect of pentoxifylline on contrast-induced nephropathy: a randomized clinical trial", International Urology and Nephrology, 2013

Publication

7

Ji, Ling, XiaoFeng Su, Wei Qin, XuHua Mi, Fei Liu, XiaoHong Tang, Zi Li, and LiChuan Yang. "A Novel Risk Score of Contrast-induced Nephropathy after Percutaneous Coronary Intervention : A new risk score model to assess risk of CIN.", Nephrology, 2015.

Publication

% 1

8

Submitted to Midwestern University

Student Paper

% 1

9

Charalampos Mamoulakis, Konstantinos Tsarouhas, Irimi Fragkiadoulaki, Ioannis Heretis et al. "Contrast-induced nephropathy: Basic concepts, pathophysiological implications and prevention strategies", Pharmacology & Therapeutics, 2017

Publication

% 1

10

Sahar A. Khaleel, Nahed A. Raslan, Amany A. Alzokaky, Mohamed G. Ewees et al. "Contrast media (meglumine diatrizoate) aggravates renal inflammation, oxidative DNA damage and apoptosis in diabetic rats which is restored by sulforaphane through Nrf2/HO-1 reactivation",

% 1

Chemico-Biological Interactions, 2019

Publication

11

ShaoBin Duan. "The Protective Role of Telmisartan Against Nephrotoxicity Induced by X-ray Contrast Media in Rat Model", Acta Radiologica, 2009

Publication

% 1

12

Abhay Kumar Chaudhary, Vijay Pathak, Shekhar Kunal, Shubhra Shukla, Pooja Pathak. "CHA2DS2-VASc score as a novel predictor for contrast-induced nephropathy after percutaneous coronary intervention in acute coronary syndrome", Indian Heart Journal, 2019

Publication

% 1

13

Submitted to University of Auckland

Student Paper

% 1

14

www.karger.com

Internet Source

% 1

15

Textbook of Cardiovascular Intervention, 2014.

Publication

% 1

16

Submitted to University of Teesside

Student Paper

% 1

17

Montree Tungjai, Singhadeth Sukantamala, Pimchanok Malasaem, Nathupakorn Dechsupa, Suchart Kothan. "An Evaluation of the Antioxidant Properties of Iodinated Radiographic

% 1

Contrast Media: An in Vitro Study", Toxicology Reports, 2018

Publication

18

Submitted to Monash University

Student Paper

% 1

19

link.springer.com

Internet Source

% 1

20

www.dovepress.com

Internet Source

% 1

21

Zaytseva, N.V.. "Contrast-induced nephropathy in patients with type 2 diabetes during coronary angiography: Risk-factors and prognostic value", Diabetes Research and Clinical Practice, 200912

Publication

% 1

22

Kota Komiyama, Takashi Ashikaga, Dai Inagaki, Tomonori Miyabe et al. "Sodium Bicarbonate-Ascorbic Acid Combination for Prevention of Contrast-Induced Nephropathy in Chronic Kidney Disease Patients Undergoing Catheterization", Circulation Journal, 2017

Publication

% 1

23

Submitted to Marquette University

Student Paper

<% 1

24

Ishibashi, Y., M. Yamauchi, H. Musha, T. Mikami, K. Kawasaki, and F. Miyake. "Impact of

<% 1

Contrast-Induced Nephropathy and Cardiovascular Events by Serum Cystatin C in Renal Insufficiency Patients Undergoing Cardiac Catheterization", *Angiology*, 2010.

Publication

25

"Guidelines on the Use of Iodinated Contrast Media in Patients With Kidney Disease 2012", *Circulation Journal*, 2013.

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5 WORDS