

Erfina_Effectiveness_of_a_Multi modal_Nursing_Intervention.57 .pdf *by*

Submission date: 06-Dec-2022 01:58PM (UTC+0700)

Submission ID: 1973006803

File name: Erfina_Effectiveness_of_a_Multimodal_Nursing_Intervention.57.pdf (397.22K)

Word count: 6349

Character count: 33946

modalities in managing symptoms experienced by patients with cancer is increasingly important to providing holistic care. **Implication for Practice:** Gynecology nurses can lead the implementation of MNI to decrease patient fatigue and depression and to increase sleep quality.

Cancer is the first or second leading cause of death globally among patients less than 70 years old in 112 of 183 countries, with an estimated around 10.0 million in 2020 or 1 in 6 deaths being caused by cancer.¹ Gynecological cancers are the fourth most common cancers in women internationally.² Gynecological cancers include cancers of the cervix, ovary, fallopian tube, vagina, and vulva.² Cervical cancer has the highest prevalence in Indonesia and is the most common type requiring hospitalization (12.8%).³

Patients with gynecological cancer have physical and psychosocial concerns, as well as a need for information and psychological support during and after their cancer treatment.⁴ Side effects result from toxic effects of cancer treatments on the body's systems.⁵ Studies have reported physical changes including fatigue, pain, sleep disorders, menopausal symptoms, kidney disorders, liver disorders, infertility, neuropathy, and skin damage.⁶ Treatments can also influence psychological changes, including the development of depression and uncertainty, and negative perceptions can lead to quality of life problems for people with cancer.⁷

The use of nonpharmacological modalities in the management of symptoms experienced by patients with cancer plays an increasingly important contribution in providing holistic patient care, including as co-analgesic.⁸ Research on nonpharmacological modalities in cancer management is, therefore, important. A variety of interventions have been developed to deal with physical effects of fatigue (eg, physical exercise/activity, resistance exercise, aerobic exercise, yoga, progressive muscle relaxation [PMR]),^{9,10} and to provide psychoeducational support (eg, health education, psychosocial support, spiritual group therapy).^{11,12} However, previous studies have often assessed physical and psychological responses, especially fatigue and psychological responses, separately.

A multimodal, nonpharmacological intervention involves integration of components such as physical programs, restorative approaches, or supportive-expressive or cognitive-behavioral psychosocial interventions, which have been identified as having the benefit of addressing multiple symptoms and as being acceptable to people with cancer.¹³ However, limited studies have reported

the effects of multimodal, nonpharmacological interventions on the physical and psychological status of women with gynecological cancer.¹³ Oncology nurses need to understand multimodal nursing interventions (MNIs) to provide a positive effect on fatigue, quality of sleep, and depression in women with gynecologic cancer. The aim of this study was to examine the effect of an MNI on quality of sleep, fatigue, and level of depression among Indonesian women with gynecological cancer.

■ Conceptual Framework

Multimodal nursing intervention is based on a nursing theory adapted from the work of Roy¹⁴ (Figure 1). According to Roy, the goal of nursing is to increase individual adaptation to a stimulus that a person faces; the adaptation can assist them to obtain health, improve quality of life, or a peaceful death. Its application in the nursing process is a problem-solving method for identifying stimuli and assessing functions in the adaptation model.¹⁴ Multimodal nursing intervention is a combination of PMR intervention and psychological counseling. Progressive muscle relaxation is a well-known mind-body intervention where movement directs the individual's attention to skeletal muscles with the goal of relaxing the whole body. Progressive muscle relaxation has been used by patients with cancer to manage severe symptom burden.¹⁵ Counseling psychology is a psychological treatment designed to improve adjustment to one's daily environment and increase the sense of well-being. Such counseling has fostered patients' coping with side effects of ongoing medical treatments, including surgery, chemotherapy or radiotherapy.¹⁶

■ Methods

Design and Participants

A quasi-experimental design approach was adopted using an equivalent control group pretest-posttest design. A convenience sampling




Figure 1 ■ Theoretical model of MNI based on nursing theory adaptation from the work of Roy.¹⁴

method⁸ was used to recruit eligible participants from a tertiary hospital in South Sulawesi, the national referral hospital with the highest number of hospitalized patients with cervical cancer. Inclusion criteria were women 18 years or older with first- to third-stage gynecological cancer. Patients with cardiovascular disease, chronic kidney disease, acute or mobility limitations, and receiving antidepressant therapy were excluded. To prevent subject pool contamination, patients in the experimental and control groups were selected from the population using cluster method. The cluster method was created according to the patient's room number; patients who had an odd room number were allocated to the experimental group, and those in even number rooms were allocated to the control group. All rooms included 6 beds, and room separation for the experimental and control groups was intended to reduce research bias or contamination. In Indonesian culture, patients are accompanied by 1 family member during hospitalization.

Sample size calculation with different mean hypothesis testing was conducted and determined that 50 participants should be included in both groups to achieve 80% power, with 2-sided α level of .05, using the score of SD according to Charalambous et al,⁹ and 10% anticipated dropout. This sample was then divided into 2 groups, with 25 participants in each group. The control group (n = 25) received the routine care offered by the hospital. The intervention group (n = 25) received an MNI combining physical and psychological interventions adapted from the Nursing Intervention Classification (NIC).¹⁷ The physical intervention involved PMR, and the psychological intervention entailed using psychological counseling. Two patients declined to participate in the study, whereas the other 4 patients were ineligible to take part in this study because of their fourth-stage cancer.

7 Data Collection

Data collection was carried out between August and December 2018. The sociodemographic, quality of sleep, fatigue, and depression questionnaires⁷ were collected by the research team. Potential participants⁷ that met the inclusion criteria were recruited by a team member and if interested were provided information about the study including the aims, processes, confidentiality, and their right to withdraw at any time. The participants who agreed to participate provided written informed consent. After a minimum of 3 days of hospital admission in each group, the pretest data were collected. The posttest data were gathered after 2 weeks of the intervention period for the intervention group and after 2 weeks of hospital routine care for the control group.

INTERVENTION GROUP

The MNI program was performed 1 to 2 weeks after hospital admission and included 2 combination therapies: PMR and psychological counseling. Progressive muscle relaxation is a nursing intervention from the NIC defined as facilitating tensing and releasing of successive muscle groups and impacting different responses. Psychological counseling is a nursing intervention including therapeutic communication⁴⁵ through counseling and has been widely used to minimize stress, improve quality of life, reduce symptoms of depression, and provide support during diagnosis and recovery.¹⁸ The procedure of PMR involved using a sitting or lying position

and relaxing from head to foot for 10 major muscle groups. Participants¹² received a module on PMR for guidance in performing the intervention, and the researchers then instructed the intervention group on how to choose a relaxed²¹ position. This procedure was performed every day for 2 weeks, 3 times per day, and for 20 minutes per session. The instructor provided guidance every day in the morning for 2 weeks, and the participants did the exercises themselves with a module PMR as an exercise guidance. A weekly schedule was developed for each participant and included 3 supervised sessions weekly. A research team member, qualified in PMR, performed all PMR and provided these individually to each participant.

The second intervention entailed psychological counseling. This intervention was offered by 2 mental health nurses who had educational backgrounds in psychiatric nursing. The intervention was performed twice weekly over 2 weeks and for 30 minutes per session. This nursing intervention focused on patient information and psychosocial needs, and patient expression of emotional feelings to help modulate distress and improve his or her ability to deal with psychological challenges. Psychosocial counseling interventions included asking and listening, giving compliments, limited advice, evaluation of understanding, and therapeutic communication.

CONTROL GROUP

The control group received the routine care offered by the hospital. The nurses provided routine interventions such as medication administration and monitoring of vital signs, but with no specific or standard intervention to overcome physical and/or psychological responses.

The protocol for the study is presented in Figure 2.

INSTRUMENTS

The study used a sociodemographic questionnaire, quality of sleep scale, fatigue scale, and depression scale. The sociodemographic questionnaire⁵⁴ included age, educational level, marital status, type of cancer, stage of cancer,¹⁶ and type of medical therapy.

Quality of sleep was assessed by the Pittsburgh Sleep Quality Index (PSQI), a standard self-reporting tool.¹⁹ The PSQI form contains 19 items in 7 components: subjective sleep quality (question 9), sleep latency (questions 2 and 5a), sleep duration (question 4), sleep efficiency (questions 1, 3, and 4), sleep disturbance (questions 5b-5j), use of sleep medication (question 6), and daytime dysfunction (questions 7 and 8). In scoring the PSQI, 7 component scores are derived, each scored 0 (no difficulty) to 3 (severe difficulty). The component scores are summed to produce a global score⁵⁰ (range, 0–21). Higher scores indicate poor quality of sleep, and lower scores indicate good quality of sleep. The PSQI Indonesian version was adapted from the original PSQI with Cronbach's α coefficient of .74.²⁵

Fatigue was assessed using the Piper Fatigue Scale (PFS).²¹ The PFS consists of 22 items with a numerical scale range between 0 and 10 and 5 open-ended questions.¹¹ A high score indicates high levels of fatigue. The PFS when translated into Bahasa Indonesia had a Cronbach's α coefficient of .98.⁶ The open-ended questions in the PFS instrument consist of 5 questions: length of fatigue feeling, contributing factors of fatigue, how to relieve fatigue feeling, additional information related fatigue, and other symptoms.




Figure 2 ■ Flowchart of study protocol.

Depression levels are measured by the Beck Depression Inventory-II (BDI-II). Ratings from the 21 items were summed to calculate a total score, which can range from 0 to 63. Scores of 0 to 13 are suggestive of minimal depression, scores of 14 to 19 are indicative of mild depression, scores of 20 to 28 are indicative of moderate depression, and scores of 29 or greater are suggestive of severe depression.²² The Indonesian BDI-II reliability, as reported by Cronbach's α coefficient, in this study was high ($\alpha = .90$).²³

DATA ANALYSIS

The sociodemographic data were analyzed using frequencies for categorical variables and means and SDs for interval-level variables. The outcomes of this study were analyzed using independent t test, chi-square, Mann-Whitney U test, Wilcoxon, marginal homogeneity, Kolmogorov-Smirnov, and a regression analysis, along with power and effect size. A value of .05 was considered statistically significant. All analyses were performed using SPSS 20.0 for Windows. The responses to the open-ended questions of the PFS instrument were analyzed by first grouping the same or highly similar words of the participants' responses, categorizing the responses, and then coding the contents of each category and quantifying the codes with frequencies.

ETHICS

The research received ethical review and approvals from the relevant institutional review board of the University (admission

number: 639/H4.8.4.5.31/PP36-KOMETIK/2018). The study was performed following the Helsinki Declaration. Participants took part voluntarily after receiving an explanation of the aims and requirements of the research, and written informed consent was obtained. Participants were assured that they could withdraw from the study at any time without stating a reason.


■ Results

Sample

Data were collected over a 5-month recruitment process from 50 women with gynecologic cancer. The mean age of the women in the intervention group was 46.77 ± 7.86 years, whereas the mean age of the women in the control group was 48.08 ± 8.3 years. Most of the patients in both groups were married (intervention, 84.0%; control, 88%). More than half in both groups were diagnosed with cervical cancer: 52% in the intervention and 60% in the control group. The majority of participants were diagnosed with stage III cancer (intervention, 56%; control, 40%) and received chemotherapy (intervention, 60%; control, 52%). No significant differences were found in any of the sociodemographic characteristics between the intervention and control groups at baseline (Table 1).

THE BETWEEN-GROUP DIFFERENCES

Comparing the difference in PSQI scores pre- and posttest, both groups yielded meaningful results. The total PSQI score of the

 **Table 1 • Characteristics of the Respondent**

	Intervention Group (n = 25)		Control Group (n = 25)		χ^2/t	P
	Mean (SD)		Mean (SD)			
Age	46.77 (7.86)		48.08 (8.38)		-0.613a	.54
Educational level	n	%	n	%	0.865	.649
Primary school	11	44	8	32		
High school	10	40	13	52		
University degree	4	16	4	16		
Marital status					0.000	1.000
Married	21	84.0	22	88		
Single	4	16.0	3	12		
Type of cancer					2.143	.343
CA cervix	13	52	15	60		
CA ovary	10	40	10	40		
CA endometrium	2	8	0	0		
Stage of cancer					1.282	.258
I	0	0	0	0		
II	11	44	15	60		
III	14	56	10	40		
Type of therapy					0.325	.569
Chemotherapy	15	60	13	52		
Other adjuvant treatment	10	40	12	48		


Abbreviation: CA, cancer.

 $P < .05$.

posttest indicated a statistically significant difference between the 2 groups ($P < .05$). No statistically significant difference was observed in the total pretest PSQI score in both groups ($P > .05$). The intervention group showed a difference in PSQI score pre- and posttest of 1.60 ± 1.323 , and the control group showed a difference PSQI score on pre- and posttest of 0.12 ± 1.56 , which is statistically significant ($P < .05$) (Table 2). No statistically significant difference was observed in the total PFS pretest scores between the intervention and control groups (119.04 ± 36.16 and 102.24 ± 52.47 ; $P > .05$) or in the total posttest PFS scores between the 2 groups (76.52 ± 24.43 and 93.24 ± 51.12 ; $P > .05$). No statistically significant difference was observed in the BDI-II pretest mean scores between the intervention and control groups ($P > .05$). The BDI-II posttest mean scores showed no statistically significant difference between the 2 groups ($P > .05$) (Table 3).

Table 2 also describes the difference in PFS scores pre- and posttest in the intervention and control groups. The differences of PFS scores pre- and posttest within both groups showed a significant difference ($P < .05$), with the observed differences to be 42.52 ± 19.74 and 9.00 ± 14.52 (Table 2).

Table 4 describes the responses to the 5 open-ended questions from the PFS instrument. All participants in the intervention ($n = 25$) and control groups ($n = 25$) completed the PFS open-ended questions. The duration of fatigue experienced by participants in the intervention group was 2 to 12 months as reported in the pretest ($n = 17$, 68%), and declined to a few days as reported in the posttest ($n = 15$, 60%). Most participants in the control group reported a duration of fatigue of 2 to 12 months in the pretest ($n = 15$, 60%) and of 2 to 12 months in the posttest ($n = 15$, 60%). Contributing factors to fatigue were pain, symptoms of disease, lack of sleep, loss of appetite, headache, breathlessness, ascites, and chemotherapy. Pain was the most frequently identified contributing factor for fatigue in the pretest in the intervention group ($n = 14$, 56%) and the control group ($n = 18$, 72%). In the posttest, pain remained the most frequent contributing factor for fatigue in the intervention group ($n = 9$, 36%) and the control group ($n = 14$, 14%). Participants reported efforts to relieve their fatigue by taking a rest and sleeping, praying, lying down, distraction, and massage. The combination of taking a rest and sleeping was the main endeavor to relieve feelings of

 **Table 2 • Comparison of Pretest, Posttest, and Difference Score of the Intervention and Control Groups in Quality of Sleep and Fatigue (N = 50)**

Groups	PSQI Score			P	PFS Score			P
	Pretest	Posttest	Difference		Pretest	Posttest	Difference	
	Mean \pm SD	Mean \pm SD	Mean \pm SD		Mean \pm SD	Mean \pm SD	Mean \pm SD	
Intervention (n = 25)	4.92 \pm 1.75	3.32 \pm 1.28	1.60 \pm 1.323	.000	119.04 \pm 36.16	76.52 \pm 24.43	42.52 \pm 19.74	.000
Control (n = 25)	5.04 \pm 2.34	4.92 \pm 1.61	0.12 \pm 1.56	.000	102.24 \pm 52.47	93.24 \pm 51.12	9.00 \pm 14.52	.000
P	.898	.001	.001		.204	.112	.000	

Boldface indicated $P < .05$.

Table 3 • Comparison of Pretest, Posttest, and Change Level of Depression by the Beck Depression Inventory-II of the Intervention and Control Groups

Groups	Pretest			Posttest			P	Change of Level of Depression	
	Minimal	Mild	Moderate	Minimal	Mild	Moderate		Decrease	No Decrease
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)		n (%)	n (%)
Intervention (n = 25)	10 (40)	9 (36)	6 (24)	20 (80)	1 (4)	4 (16)	.008	10 (40)	15 (60)
Control (n = 25)	14 (56)	8 (32)	3 (12)	18 (72)	5 (20)	2 (8)	.059	6 (24)	19 (76)
P		.906			1.000				.225

Boldface indicated $P < .05$.

fatigue for the intervention groups in the pretest (n = 15, 60%) and posttest (n = 15, 60%). Lying down was the most frequent endeavor to relieve feelings of fatigue for the control groups in

the pretest (n = 12, 48%) and posttest (n = 10, 40%). Most of the participants in the intervention group did not report other symptoms at the time of the pretest (n = 14; 56%) but did report

Table 4 • Open-Ended Questions From the Piper Fatigue Scale

No. Questions	Open-Ended Questions	Response	Intervention Group		Control Group	
			Pre, n (%)	Post, n (%)	Pre, n (%)	Post, n (%)
1	Length of fatigue feeling	Days	2 (8)	15 (60)	1 (4)	4 (16)
		Weeks	2 (8)	3 (12)	2 (8)	0 (0)
		A month	2 (8)	1 (4)	1 (4)	0 (0)
		2–12 months	17 (68)	4 (16)	15 (60)	15 (60)
		>12 months	2 (8)	2 (8)	6 (24)	6 (24)
		Total	25 (100)	25 (100)	25 (100)	25 (100)
24	Contributing factors of fatigue	Pain	14 (56)	9 (36)	18 (72)	14 (56)
		Symptom of disease	3 (12)	2 (8)	3 (12)	5 (20)
		Lack of sleep	1 (4)	8 (32)	1 (4)	5 (20)
		Loss appetite	1 (4)	1 (4)	0 (0)	0 (0)
		headache	1 (4)	1 (4)	0 (0)	0 (0)
		Ascites	4 (16)	3 (12)	1 (4)	0 (0)
		breathless	1 (4)	1 (4)	0 (0)	0 (0)
		chemotherapy	0 (0)	0 (0)	1 (4)	1 (4)
		Total	25 (100)	25 (100)	25 (100)	25 (100)
		25	How to relieve fatigue feeling	Take a rest and sleep	15 (60)	15 (60)
Sleep	2 (8)			2 (8)	2 (8)	1 (4)
Pray	1 (4)			1 (4)	1 (4)	1 (4)
Lie down	7 (28)			7 (28)	12 (48)	10 (40)
Massage	0 (0)			0 (0)	2 (8)	3 (12)
Distraction	0 (0)			0 (0)	2 (8)	1 (4)
Total	25 (100)			25 (100)	25 (100)	25 (100)
26	Additional description about the fatigue feeling	None	25 (100)	25 (100)	23 (92)	24 (96)
		Lack of sleep	0 (0)	0 (0)	1 (4)	0 (0)
		pregnancy	0 (0)	0 (0)	1 (4)	1 (4)
27	Other symptoms	Total	25 (100)	25 (100)	25 (100)	25 (100)
		Yes	11 (44)	6 (24)	8 (32)	8 (32)
27	Description of symptoms	Total	25 (100)	25 (100)	25 (100)	25 (100)
		Lack of sleep	1 (4)	0 (0)	0 (0)	0 (0)
		Breathless in activity	1 (4)	0 (0)	0 (0)	0 (0)
		Breathless	4 (16)	0 (0)	5 (20)	3 (12)
		Headache	1 (4)	0 (0)	0 (0)	0 (0)
		Medical equipment	1 (4)	0 (0)	0 (0)	0 (0)
		Heartburn	2 (8)	2 (8)	0 (0)	0 (0)
		Edema	0 (0)	2 (8)	2 (8)	1 (4)
		Vomiting	0 (0)	1 (4)	0 (0)	1 (4)
		Cough	1 (4)	1 (4)	1 (4)	3 (12)
		Total	11 (44)	6 (24)	8 (32)	8 (32)

an increase in symptoms at the time of the posttest (n = 19; 76%), whereas in the control group, there were no difference in the frequencies between the pre- and posttest (n = 17; 68%).

The power and effect size of the MNI program were evaluated using linear regression analysis as shown in Table 5. The MNI program for quality of sleep had strong power (0.95) and medium effect size (0.25). The MNI program for fatigue had strong power (1) and large effect size (0.93). The MNI program for depression had strong power (0.93) and small effect size (0.23).

THE WITHIN-GROUP DIFFERENCES

The comparison of pretest, posttest, and difference scores within groups in the PSQI scores are shown in Table 2. Statistically significant differences were observed in the pre- and posttest of the PSQI score in the intervention group, which were 4.92 ± 1.75 and 3.32 ± 1.28 , respectively ($P < .05$), and the control group, which were 5.04 ± 2.34 and 4.92 ± 1.61 , respectively ($P < .05$). The PFS scores within group were statistically significantly different in the intervention and control groups; the intervention group mean scores changed from 119.04 ± 36.16 to 102.24 ± 52.47 , with $P < .05$, and the mean PFS scores in the control group changed from 76.52 ± 24.43 to 93.24 ± 51.12 , with $P < .05$ (Table 2).

The differences of the pre- and posttest within group of BDI-II score was statistically significantly different ($P < .05$). The BDI-II mean scores from pre- to posttest in the control group showed no statistically significant difference ($P > .05$) (Table 3).

Discussion

The purpose of this study was to investigate the effects of a nurse-led MNI intervention on sleep quality, fatigue, and level of depression in patients with gynecological cancer. According to the present study results, statistically significant improvement was observed in both sleep quality and fatigue by participants in the intervention group. The positive change is similar to previous studies that used PMR to improve sleep quality in patients with chronic obstructive pulmonary disease²⁴ and in patients with COVID-19.²⁵ Evidence has shown that nonpharmacological interventions contribute to improvement of sleep quality among adults in treatment for cancer who are not in remission.²⁶ Progressive muscle relaxation training

could help increase physical and mental relaxation²⁵; thus, MNI that includes PMR and counseling could improve sleep quality.

Fatigue is a distressing and persistent symptom for women with gynecological cancer and survivors.²⁷ This study revealed that the MNI program had a positive reduction effect on fatigue. Findings from this study are consistent with other studies that have assessed the efficacy of patient-control cognitive-behavioral interventions, including counseling, for patients with lung, prostate, colorectal, or gynecologic cancer who reported significant reductions in severity of pain and fatigue.²⁸ A study by Seyedi Chegeni et al,²⁴ findings indicated that PMR reduced fatigue and improved sleep quality in patients with stages 3-4 chronic obstructive pulmonary disease. Nonpharmacological interventions for the management of cancer-related fatigue have varied, including exercise/physical activity, where structured and repetitive body movement has been reported to decrease cancer-related fatigue.²⁹

The study findings showed that the difference mean scores pre- and posttest for fatigue of the participants in the intervention group were significantly decreased compared with the control group in this study (42.52 ± 19.74). Furthermore, this study reported decreased duration of feeling fatigued after the intervention, and most of the participants revealed pain and lack of sleep as factors contributing to fatigue. When reviewing all study outcomes with a holistic perspective, a significant decrease in fatigue scores in our study could be related to the effects of MNI, including PMR and counseling, which involve energy restoration in the body and increase the positive psychological state. It is possible that regular practice of PMR contributes to an increase in the oxygen concentration of muscle cells and a positive influence on patients' psychological responses.¹⁵

Although there were significant differences in the level of depression from before to after the intervention in the intervention group, there was no significant change in depression in the control group. A previous study revealed that psychological treatment is not likely to show a unified effect size. The patient's clinical state and treatment side effects were moderating factors. That study highlights a significant, small, positive overall effect size of MNI among adults with gynecological cancer.¹⁶

In this current study, the MNI program for depression had a positive impact but small effect size. One study concluded that providing psychological treatments should be considered as

Table 5 • Predictive Power and Effect Size of the MNI for Change in Difference Score Pre and PostTest of Quality of Sleep, Fatigue, and Depression

Variable	Quality of Sleep			Fatigue			Depression		
	B	SE	β	B	SE	β	B	SE	β
Constant	3.080	0.648		76.040	7.749		9.440	1.959	
MNI	-1.480	0.410	-0.462	-33.52	4.901	-0.703	-4.320	1.239	-0.450
R	0.462			0.703			0.450		
R ²	0.197			0.483			0.186		
F	0.214			0.494			0.202		
P	.001			.000			.001		
DW	2.109			1.238			1.736		
Effect size	0.25			0.93			0.23		
Power	0.95			1.0			0.93		

Abbreviations: DW, The Durbin Watson; MNI, multimodal nursing intervention.

crucial for the patient's health in cancer contexts.¹⁶ Another study concluded that active nursing interventions are necessary to reduce cancer-related fatigue and improve emotional function, sleep quality, and thereby quality of life among women with gynecological cancer.³⁰ Patients with cancer who have received planned nursing interventions have shown much lower rates of psychological symptoms such as anxiety, depression, and significant improvement of emotional well-being compared with those who received routine care offered by hospital staff.³¹ There is a need to decrease depression levels that could negatively affect the overall quality of life.

Limitations

There are limitations of this study that require acknowledgment. Women with gynecological cancer from only a single location were included, the sample size was small, and results may not be generalizable to other areas or populations. However, the study was conducted in a major hospital in the east of Indonesia, which may represent the population of eastern Indonesia. In addition, we performed only a single time assessment after the intervention. Further research with a larger scale and longer duration is recommended to examine effects of MNI on physical and psychological responses among patients with gynecological cancer.

Clinical Implications

The present study showed that MNI is a valuable nursing intervention for increasing sleep quality and reducing fatigue. A small effect size on level of depression in patients with gynecologic cancer was also noted. The best available evidence indicates that MNI is a safe nursing intervention. The MNI, together with other nonpharmacologic therapies, is anticipated to improve quality of life, especially in women with gynecologic cancer who need support in these areas. The use of MNI for these patients can be implemented by nurses.

Conclusion

Multimodal nursing intervention including PMR and counseling is an NIC that can be readily performed by nurses for women with gynecologic cancer. The results provide some evidence that combining physical and psychological interventions could decrease fatigue, increase sleep quality, and exert a small effect on levels of depression. Health services can consider training programs for clinical nurses to provide skills in MNI intervention. This would enable clinical nursing to teach PMR and provide psychological counseling to improve quality of life in patients with gynecologic cancer.

References

- Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2021;71(3):209–249.
- World Health Organization. Cancer. <https://www.who.int/news-room/fact-sheets/detail/cancer>. Published 2018. Accessed December 24, 2018.
- Pusat Data dan Informasi Kesehatan KKR. Stop kanker: Situasi penyakit kanker. infodatin-Kanker. <https://www.kemkes.go.id/article/view/15021800011/situasi-penyakit-kanker.html>. Published 2015. Accessed December 24, 2018.

- Walton LM, Reeve J, Brown PM, Farquhar CM. Gynaecologic cancer patients' needs and experiences of supportive health services in New Zealand. *Psychooncology*. 2010;208(March 2009, 208):201.
- McCorkle R, Dowd M, Ercolano E, et al. Effects of a nursing intervention on quality of life outcomes in post-surgical women with gynecological cancers. *Psychooncology*. 2009;18(1):62–70.
- Mulhaeriah, Afyanti Y, Achmad EK, Sangkala MS. Effectiveness of relaxation breathing exercise on fatigue in gynecological cancer patients undergoing chemotherapy. *Int J Nurs Sci*. 2018;5(4):331–335.
- Wu TY, Lin C. Developing and evaluating an individually tailored intervention to increase mammography adherence among Chinese American women. *Cancer Nurs*. 2015;38(1):40–49.
- Naga BSHB. Evidence-based standards for cancer pain management. 2016; 3(7):41–46.
- Charalambous A, Giannakopoulou M, Bozas E, Marcou Y, Kitsios P, Paikousis L. Guided imagery and progressive muscle relaxation as a cluster of symptoms management intervention in patients receiving chemotherapy: a randomized control trial. *PLoS One*. 2016;11(6):e0156911.
- Bartoszczyk DA, Gilbertson-White S. Interventions for nurse-related barriers in cancer pain management. *Oncol Nurs Forum*. 2015;42(6):634–641.
- Adam R, Bond C, Murchie P. Educational interventions for cancer pain. A systematic review of systematic reviews with nested narrative review of randomized controlled trials. *Patient Educ Couns*. 2015;98(3):269–282.
- Hochstenbach LM, Courtens AM, Zwakhalen SM, van Kleef M, de Witte LP. Self-management support intervention to control cancer pain in the outpatient setting: a randomized controlled trial study protocol. *BMC Cancer*. 2015;15:416.
- Bennett S, Pigott A, Beller EM, Haines T, Meredith P, Delaney C. Educational interventions for the management of cancer-related fatigue in adults. *Cochrane Database Syst Rev*. 2016;11(11):CD008144. doi:10.1002/14651858.CD008144.pub2. www.cochranelibrary.com.
- Alligood MR. *Nursing Theory Utilization & Application*. 5th ed. (Alexopoulos Y, ed.). Maryland Heights, MO: Mosby/Elsevier; 2014. doi:10.5005/jp/books/13072_29.
- Gok Metin Z, Karadas C, Izgu N, Ozdemir L, Demirci U. Effects of progressive muscle relaxation and mindfulness meditation on fatigue, coping styles, and quality of life in early breast cancer patients: An assessor blinded, three-arm, randomized controlled trial. *Eur J Oncol Nurs*. 2019;42:116–125.
- de la Torre-Luque A, Gambara H, López E, Cruzado JA. Psychological treatments to improve quality of life in cancer contexts: a meta-analysis. *Int J Clin Health Psychol*. 2016;16(2):211–219.
- Moorhead SA, McCloskey JC, Bulechek GM. Nursing interventions classification. A comparison with the Omaha System and the Home Healthcare Classification. *J Nurs Adm*. 1993;23(10):23–29.
- Galway K, Black A, Cantwell M, Cardwell CR, Mills M, Donnelly M. Psychosocial interventions to improve quality of life and emotional wellbeing for recently diagnosed cancer patients. *Cochrane Database Syst Rev*. 2012; 11(11):Article CD007064. doi:10.1002/14651858.CD007064.pub2.
- Buyssse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193–213. doi:https://doi.org/10.1016/0165-1781(89)90047-4.
- Fandiani YM, Wantiyah W, Juliningrum PP. The effect of Dzikir therapy on sleep quality of college students at School of Nursing University of Jember. *NurseLine J*. 2017;2(1):52.
- Reeve BB, Stover AM, Alfano CM, et al. The Piper Fatigue Scale-12 (PFS-12): psychometric findings and item reduction in a cohort of breast cancer survivors. *Breast Cancer Res Treat*. 2012;136(1):9–20.
- Beck AT, Steer RA, Carbin MG. Psychometric properties of the Beck Depression Inventory: twenty-five years of evaluation. *Clin Psychol Rev*. 1988;8(1):77–100.
- Ginting H, Näring G, Van Der Veld WM, Srisayekti W, Becker ES. Validating the Beck Depression Inventory-II in Indonesia's general population and coronary heart disease patients. *Int J Clin Heal Psychol*. 2013;13(3):235–242.
- Seyedi Chegeni P, Gholami M, Azargoon A, Hossein Pour AH, Birjandi M, Norollahi H. The effect of progressive muscle relaxation on the management

of fatigue and quality of sleep in patients with chronic obstructive pulmonary disease: a randomized controlled clinical trial. *Complement Ther Clin Pract*. 2018;31:64–70.

25. Liu K, Chen Y, Wu D, Lin R, Wang Z, Pan L. Effects of progressive muscle relaxation on anxiety and sleep quality in patients with COVID-19. *Complement Ther Clin Pract*. 2020;39:101132.
26. Fatmawati, Budworth L. The effectiveness of non-pharmacological interventions in improvement of sleep quality among non-remissive cancer patients: a systematic review of randomized trials. *Commun Sci Technol*. 2016;1(2):70–77.
27. Wang XS, Woodruff JF. Cancer-related and treatment-related fatigue. *Gynecol Oncol*. 2015;136(3):446–452.
28. Kwekkeboom KL, Abbott-Anderson K, Cherwin C, Roiland R, Serlin RC, Ward SE. Pilot randomized controlled trial of a patient-controlled cognitive-behavioral intervention for the pain, fatigue, and sleep disturbance symptom cluster in cancer. *J Pain Symptom Manage*. 2012;44(6):810–822.
29. Yeo TP, Cannaday S. Cancer-related fatigue: impact on patient quality of life and management approaches. *Nurs Res Rev*. 2015;5:65.
30. Zhou L, Tian Y, Yu J, Chen T, An F. Effect of at-home cognitive behavior therapy combined with nursing on revised Piper Fatigue Scale, Pittsburgh Sleep Quality Index, Self-Rating Anxiety Scale and Self-Rating Depression Scale of ovarian cancer patients after chemotherapy. *Int J Clin Exp Med*. 2020;13(6):4227–4234.
31. Sivabalan T, Upasani SV. Effectiveness of nursing interventions on physical and psychological outcome among cancer patients undergoing chemotherapy. *J Krishna Inst Med Sci Univ*. 2016;5(2):57–68.

ORIGINALITY REPORT

20%

SIMILARITY INDEX

%

INTERNET SOURCES

20%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Cong Li, Zhao-Yu Li, Qian Lu, Yu-Jie Zhou, Xue-Ying Qin, Ai-Wen Wu, Dong Pang. "The Effectiveness of a Self-management Program of Bowel Dysfunction in Patients With Mid and Low Rectal Cancer After Sphincter-Preserving Surgery", *Cancer Nursing*, 2022 2%

Publication
- 2** Zehra Gok Metin, Canan Karadas, Nur Izgu, Leyla Ozdemir, Umut Demirci. "Effects of progressive muscle relaxation and mindfulness meditation on fatigue, coping styles, and quality of life in early breast cancer patients: An assessor blinded, three-arm, randomized controlled trial", *European Journal of Oncology Nursing*, 2019 2%

Publication
- 3** Samantha J. Mayo, Sarah Brennenstuhl, Prabdeep Panesar, Ashley Leak Bryant. "Patterns of Concerns Among Hematological Cancer Survivors", *Cancer Nursing*, 2022 1%

Publication

4

Gareth R. Dutton, Karen B. Grothe, Glenn N. Jones, Dori Whitehead, Kathleen Kendra, Phillip J. Brantley. "Use of the Beck Depression Inventory-II with African American primary care patients", *General Hospital Psychiatry*, 2004

Publication

1 %

5

Evidence-based Anticancer Complementary and Alternative Medicine, 2013.

Publication

1 %

6

Yesol Yang, William Romine, Tolu O. Oyesanya, Hyeyoung K. Park, Cristina C. Hendrix. "The Association of Self-esteem With Caregiving Demands, Coping, Burden, and Health Among Caregivers of Breast Cancer Patients", *Cancer Nursing*, 2021

Publication

1 %

7

Suni Hariati, Lisa McKenna, Lely Lusmilasari, Sonia Reisenhofer, Retno Sutomo, Andi Dwi Bahagia Febriani, Dian Sidik Arsyad. "Translation, Adaptation and Psychometric Validation of the Indonesian Version of the Readiness for Hospital Discharge Scale for Parents of Low Birth Weight Infants", *Journal of Pediatric Nursing*, 2020

Publication

1 %

8

Suni Hariati, Retno Sutomo, Lisa McKenna, Sonia Reisenhofer, Lely Lusmilasari, Andi D. B.

1 %

Febriani. "Indonesian mothers' beliefs on caring practices at home for preterm babies after hospital discharge: A qualitative study", *Journal for Specialists in Pediatric Nursing*, 2021

Publication

9

Ranin Ballan, Simon J. Durrant, Robert Stickgold, Alexandra Morgan, Dara S. Manoach, Yafit Gabay. "A failure of sleep-dependent consolidation of visuoperceptual procedural learning in young adults with ADHD", *Translational Psychiatry*, 2022

Publication

10

Hyang Rang Park, Hee Sun Kang, Soo Hyun Kim, Savitri Singh-Carlson. "Effect of a Smart Pill Bottle Reminder Intervention on Medication Adherence, Self-efficacy, and Depression in Breast Cancer Survivors", *Cancer Nursing*, 2021

Publication

11

Mulhaeriah, Yati Afiyanti, Engkus Kusdinar Achmad, Moh Syafar Sangkala. "Effectiveness of Relaxation Breathing Exercise on fatigue in gynecological cancer patients undergoing chemotherapy", *International Journal of Nursing Sciences*, 2018

Publication

1 %

1 %

1 %

12

Kai Liu, Ying Chen, Duozhi Wu, Ruzheng Lin, Zaisheng Wang, Liqing Pan. "Effects of progressive muscle relaxation on anxiety and sleep quality in patients with COVID-19", *Complementary Therapies in Clinical Practice*, 2020

Publication

<1 %

13

Mahsa Musavi, Simin Jahani, Marziyeh Asadzaker, Elham Maraghi, Sasan Razmjoo. "The Effect of Pain Self-Management Education on Pain Severity and Quality of Life in Metastatic Cancer Patients", *Asia-Pacific Journal of Oncology Nursing*, 2021

Publication

<1 %

14

Nemat Ismail Abdel Aziz Ismail, Wafaa Taha Ibrahim Elgzar. "The Effect of Progressive Muscle Relaxation on Post Cesarean Section Pain, Quality of Sleep and Physical Activities Limitation", *International Journal of Studies in Nursing*, 2018

Publication

<1 %

15

Zhiyou Peng, Lin Li, Yuan Chen, Zhiying Feng, Xiangming Fang. "WeChat app-based reinforced education improves the quality of opioid titration treatment of cancer-related pain in outpatients: a randomized control study", *BMC Cancer*, 2020

Publication

<1 %

16

Anita Kjeverud, Kristin Østlie, Anne-Kristine Schanke, Caryl Gay, Magne Thoresen, Anners Lerdal. "Trajectories of fatigue among stroke patients from the acute phase to 18 months post-injury: A latent class analysis", PLOS ONE, 2020

Publication

<1 %

17

Berger, Ann M., Sandra A. Mitchell, Paul B. Jacobsen, and William F. Pirl. "Screening, evaluation, and management of cancer-related fatigue: Ready for implementation to practice? : Cancer-Related Fatigue", CA A Cancer Journal for Clinicians, 2015.

Publication

<1 %

18

Kwekkeboom, Kristine L., Kristen Abbott-Anderson, Catherine Cherwin, Rachel Roiland, Ronald C. Serlin, and Sandra E. Ward. "Pilot Randomized Controlled Trial of a Patient-Controlled Cognitive-Behavioral Intervention for the Pain, Fatigue, and Sleep Disturbance Symptom Cluster in Cancer", Journal of Pain and Symptom Management, 2012.

Publication

<1 %

19

Kusrini S. Kadar, Nurul Fadhilah Gani, Erfina Erfina, Suni Hariati. "Self-care management and health outcomes among Indonesian pregnant women", Enfermería Clínica, 2020

Publication

<1 %

20

Lai, Xiaobin, Frances Kam Yuet Wong, Carenx Wai Yee Leung, Lai Ha Lee, Jessica Shuk Yin Wong, Yim Fan Lo, and Shirley Siu Yin Ching. "Development and Assessment of the Feasibility of a Nurse-Led Care Program for Cancer Patients in a Chemotherapy Day Center : Results of the Pilot Study", *Cancer Nursing*, 2014.

Publication

<1 %

21

Mishra, Shiraz I, Roberta W Scherer, Paula M Geigle, Debra R Berlanstein, Ozlem Topaloglu, Carolyn C Gotay, Claire Snyder, and Shiraz I Mishra. "Exercise interventions on health-related quality of life for cancer survivors", *Cochrane Database of Systematic Reviews*, 2012.

Publication

<1 %

22

Reva Gündoğan, Emine Kaplan Serin. "The Effects of Progressive Relaxation Exercises on COVID-19-Related Fear, Anxiety, and Sleep Quality in Emergency Aid Station Employees", *Holistic Nursing Practice*, 2022

Publication

<1 %

23

Sally Bennett. "Educational interventions for the management of cancer-related fatigue in adults", *Cochrane Database of Systematic Reviews* Protocols, 10/07/2009

Publication

<1 %

24

"The MASCC Textbook of Cancer Supportive Care and Survivorship", Springer Science and Business Media LLC, 2018

Publication

<1 %

25

Teresa Corbett, Jane C Walsh, AnnMarie Groarke, Rona Moss-Morris, Brian E McGuire. "Protocol for a pilot randomised controlled trial of an online intervention for post-treatment cancer survivors with persistent fatigue", BMJ Open, 2016

Publication

<1 %

26

Xinqiong Zhang, Qin Wang, Xiaomin Zhang, Xiaoting Wu, Qiuping Wang, Jingfang Hong. "A Cognitive-Behavioral Intervention for the Symptom Clusters of Chinese Patients With Gastrointestinal Tract Cancer Undergoing Chemotherapy", Cancer Nursing, 2019

Publication

<1 %

27

Karen Kayser. "The Nature of Cancer and Its Psychosocial Challenges", Helping Couples Cope with Women's Cancers, 2008

Publication

<1 %

28

Grace B. Campbell, Sarah M. Belcher, Young Ji Lee, Madeleine Courtney-Brooks et al. "Intensive Daily Symptom and Function Monitoring Is Feasible and Acceptable to Women Undergoing First-Line Chemotherapy

<1 %

29

Abdallah Abu Khait, Mark Lazenby.

"Psychosocial-spiritual interventions among Muslims undergoing treatment for cancer: an integrative review", BMC Palliative Care, 2021

Publication

<1 %

30

Andersen, Christina, Mikael RÃ¸rth, Bent Ejlersen, Maria Stage, Tom MÃ¸ller, Julie Midtgaard, Morten Quist, Kira Bloomquist, and Lis Adamsen. "The effects of a six-week supervised multimodal exercise intervention during chemotherapy on cancer-related fatigue", European Journal of Oncology Nursing, 2012.

Publication

<1 %

31

MohammadJavad Ghazanfari, Samad Karkhah, Amir EmamiZeydi, Hamed Mortazavi, Amir Tabatabaee, Mohsen Adib-Hajbaghery. "A Systematic Review of Potentially Effective Nonpharmacological Interventions for Reducing Fatigue among Iranian Patients Who Receive Hemodialysis", Complementary Medicine Research, 2022

Publication

<1 %

32

Mosleh Kareem, Diyar Hussein Taher. "Effect of Relaxation Training Techniques on Anxiety

<1 %

and Depression among Breast Cancer Patients", Open Access Macedonian Journal of Medical Sciences, 2022

Publication

33

Qianqian Zhu, Lulu Xu, Yexin Chen, Dake Shi, Hong Huang, Yong Cai. "Sleep Quality and Mental Health among Asymptomatic COVID-19 Carriers from Fangcang Shelter Hospitals : The Moderating Role of Social Support", Research Square Platform LLC, 2022

<1 %

Publication

34

"Handbook of Quality of Life in Cancer", Springer Science and Business Media LLC, 2022

<1 %

Publication

35

Andrea Christoforou, Marietta L. van der Linden, Pelagia Koufaki. "Short-term effectiveness of a community-implemented falls prevention referral service", Disability and Rehabilitation, 2017

<1 %

Publication

36

Devianti Tandiallo, Mardiana Ahmad, Syafruddin Syarif, Nasrudin A. Mappaware, Prihantono Prihantono, Burhanuddin Bahar. "Web-Based Partograph on Early Detection of Emergency Cases and Referral Processes", Global Journal of Health Science, 2019

<1 %

Publication

37

Maanya Rajasree Katta, Sai Sudha Valisekka, Pahel Agarwal, Maha Hameed et al. "Non-pharmacological integrative therapies for chronic cancer pain", Journal of Oncology Pharmacy Practice, 2022

Publication

<1 %

38

Yubin Lee, Yoorim Hyun, Myoungsoon You, Heeyoung Lee, Jin-Ok Han, Suin Seo. "The effect of resource loss on depression and peritraumatic distress during the early period of the COVID-19: considering the pandemic-situational and social context", Research Square Platform LLC, 2022

Publication

<1 %

39

Advances in Diagnosis and Management of Ovarian Cancer, 2014.

Publication

<1 %

40

Chich-Hsiu Hung. "Predictors of Postpartum Women's Health Status", Journal of Nursing Scholarship, 12/2004

Publication

<1 %

41

Ran Jinnin, Yasumasa Okamoto, Koki Takagaki, Yoshiko Nishiyama et al. "Detailed course of depressive symptoms and risk for developing depression in late adolescents with subthreshold depression: a cohort study", Neuropsychiatric Disease and Treatment, 2016

<1 %

42 Wenhua Yu, Qian Lu, Yuhan Lu, Renxiu Guo, Xiaoting Hou, Hong Yang. "Relationship Between Anticipatory Grief and Communication Avoidance in Chinese Caregivers of Patients With Advanced Cancer", Cancer Nursing, 2021

Publication

43 Winnie K. W. So, Bernard M. H. Law, Dorothy N. S. Chan, Weijie Xing, Carmen W. H. Chan, Alexandra L. McCarthy. "The Effect of Nonpharmacological Interventions on Managing Symptom Clusters Among Cancer Patients", Cancer Nursing, 2020

Publication

44 Young Ji Lee, Albert Park, Mary Roberge, Heidi Donovan. "What Can Social Media Tell Us About Patient Symptoms", Cancer Nursing, 2022

Publication

45 Cramp, Fiona, James Byron-Daniel, and Fiona Cramp. "Exercise for the management of cancer-related fatigue in adults", Cochrane Database of Systematic Reviews Reviews, 2012.

Publication

46 Marketa Ciharova, Hynek Cígler, Veronika Dostálová, Gabriela Šivicová, Ondrej Bezdicek.

<1 %

"Beck depression inventory, second edition, Czech version: demographic correlates, factor structure and comparison with foreign data", International Journal of Psychiatry in Clinical Practice, 2020

Publication

47

Oksoo Kim, Heeja Jung. "Prediction model for abnormal eating behaviour among hospital nurses: A structural equation modelling approach", International Journal of Nursing Practice, 2021

Publication

<1 %

48

Sermin Timur Taşhan, Yeşim Aksoy Derya, Tuba Uçar, Gülçin Nacar, Behice Erci. "Theory-based training to promote breast cancer screening among women with breast cancer worries: randomized controlled trial", Sao Paulo Medical Journal, 2020

Publication

<1 %

49

Davood Hekmatpou, Azadeh Nasiri, Fatholah Mohaghegh. "Investigating the Effect of Self-Care Training on Life Expectancy and Quality of Life in Patients with Gastrointestinal Cancer under Radiotherapy", Asia-Pacific Journal of Oncology Nursing, 2019

Publication

<1 %

50

Jacobien Marije Kieffer. "Item-order effects in the Oral Health Impact Profile (OHIP)",

<1 %

51

Kairen McCloy, Ciara Hughes, Dr Lynn Dunwoody, Dr Joanne Marley, Dr Jackie Gracey. "Effects of mindfulness - based interventions on fatigue and psychological wellbeing in women with cancer: A systematic review and meta - analysis of randomised control trials", Psycho-Oncology, 2022

Publication

<1 %

Exclude quotes On

Exclude matches < 4 words

Exclude bibliography On