

# 19.\_Agussalim\_Bukhari- Najdah2.pdf

*by*

---

**Submission date:** 13-Jan-2023 02:00PM (UTC+0700)

**Submission ID:** 1992178987

**File name:** 19.\_Agussalim\_Bukhari-Najdah2.pdf (809.33K)

**Word count:** 10905

**Character count:** 56714



Contents lists available at ScienceDirect

Gene Reports

journal homepage: [www.elsevier.com/locate/genrep](http://www.elsevier.com/locate/genrep)



22

## Expression of *Vitamin D Receptor (VDR)* gene and *VDR* polymorphism rs11574113 in pulmonary tuberculosis patients and their household contacts

Najdah Hidayah<sup>a</sup>, Irawaty Djaharuddin<sup>b,c</sup>, Ahyar Ahmad<sup>d</sup>, Agussalim Bukhari<sup>e</sup>, Ilhamjaya Patellongi<sup>f</sup>, Nur Ahmad Tabri<sup>b,c</sup>, Rosana Agus<sup>g</sup>, Subair Subair<sup>a,h</sup>, Irda Handayani<sup>i</sup>, Andi Tenriola<sup>a</sup>, Handayani Halik<sup>a,h</sup>, Muhammad Nasrum Massi<sup>j,\*</sup>

<sup>a</sup> Postgraduate (Doctoral) Program, Faculty of Medicine, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>b</sup> Department of Pulmonology and Respiratory Diseases, Faculty of Medicine, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>c</sup> Dr. Wahidin Sudirohusodo Hospital, Makassar 90245, South Sulawesi, Indonesia

<sup>d</sup> Department of Chemistry, Mathematics and Natural Science Faculty, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>e</sup> Department of Clinical Nutrition, Faculty of Medicine, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>f</sup> Department of Microbiology, Faculty of Medicine, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>g</sup> Department of Biology, Mathematics and Natural Science Faculty, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>h</sup> Rezeky University, Makassar 90234, South Sulawesi, Indonesia

<sup>i</sup> Department of Clinical Pathology, Faculty of Medicine, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

<sup>j</sup> Department of Clinical Microbiology, Faculty of Medicine, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia

### ARTICLE INFO

#### Keywords:

Gene expression  
IGRA  
qPCR  
rs11574113  
Tuberculosis  
VDR

### ABSTRACT

80

The vitamin D receptor mediates the effect of vitamin D, which has received substantial attention in its involvement in reducing the risk of tuberculosis (TB). This study sought to see if *VDR* gene expression and *VDR* SNP rs11574113 was linked to a person's likelihood of developing tuberculosis. New Pulmonary Tuberculosis (PTB) *VDR* gene expression and Household Contact (HC) samples *VDR* gene expressions were examined by the quantitative real-time PCR (qPCR) method, while *VDR* SNP rs11574113 was identified by PCR followed by sequencing. Latent TB infection in household contacts was screened using an Interferon-Gamma Release Assay (IGRA). Eighty-three PTB, 77 HCs (45 positive-IGRA and 28 negative-IGRA) were involved in this study. Among the 45 positive-IGRA HC, 28 (62.2%) were spouses of index cases. IGRA results showed that 28 out of 37 (75.7%) samples of all spouses recruited have positive IGRA results. *VDR* gene expression of PTB was 0.2 and 0.25 downregulated compared to negative-IGRA and positive-IGRA HC, respectively ( $p = 0.014$  and  $p = 0.017$ ). *VDR* gene expression of positive-IGRA was also 0.788 downregulated compared to negative-IGRA HC but did not reach statistical significance ( $p = 0.945$ ). The high expression of the *VDR* gene may be influenced by variant of *VDR* SNP rs11574113. G/G genotype and G/C genotype were associated with higher *VDR* gene expression compared to C/C genotype ( $OR^a = 9.754$  95% CI 1.482–64.195,  $p = 0.018$  and  $OR^a = 9.723$  95% CI 1.416–66.751,  $p = 0.021$ , respectively). However, rs11574113 was not found to be associated with TB incidence and IGRA positivity. Approximately 20% of the people with PTB had higher levels of *VDR* gene expression than those who were not ( $OR^a = 0.736$  95% CI 0.542–0.998,  $p = 0.049$ ). *VDR* gene expression may be associated with the susceptibility of tuberculosis. This study also shows an immense rate of TB infection among spouses of PTB patients based on IGRA.

**Abbreviations:** Mtb, *Mycobacterium tuberculosis*; *VDR*, *Vitamin D Receptor*; TB, tuberculosis; PTB, pulmonary tuberculosis; IGRA, Interferon-Gamma Release Assay; HC, household contact; *GAPDH*, glyceraldehyde-3 phosphate dehydrogenase; qPCR, quantitative real-time PCR; Ct, cycle threshold; TNF, tumor necrosis factor; CYP27B1, *cytochrome p450 27B1*; UTR, untranslated region; SNP, single nucleotide polymorphism.

\* Corresponding author at: Department of Clinical Microbiology, Faculty of Medicine, Universitas Hasanuddin, Jl. Perintis Kemerdekaan KM.10 Tamalanrea, Makassar 90245, South Sulawesi, Indonesia.

E-mail addresses: [nasrumm@unhas.ac.id](mailto:nasrumm@unhas.ac.id), [nasrum.unhas@gmail.com](mailto:nasrum.unhas@gmail.com), [nasrumm2000@yahoo.com](mailto:nasrumm2000@yahoo.com) (M.N. Massi).

<https://doi.org/10.1016/j.genrep.2022.101581>

Received 7 November 2021; Received in revised form 20 February 2022; Accepted 25 February 2022

Available online 28 February 2022

2452-0144/© 2022 Elsevier Inc. All rights reserved.

## 1. Introduction

With about 10 million infections and 1.4 million deaths, tuberculosis is a major global public health concern. Indonesia is among the eight countries that contribute to two-thirds of TB cases worldwide and ranks second in the highest TB incidence (8.5%), with case notifications increasing by about 69% from 2015 to 2019 (World Health Organization, 2020). TB elimination can be achieved not only by effective active TB therapy but also by early case detection to stop transmission through various strategies for preventing new infections and neutralizing the existing dormant infections. Without tackling reservoirs of latent infection, the WHO's "End TB" strategy is difficult to be accomplished (Dye et al., 2013). Around one-fourth of the population worldwide has been infected by *Mycobacterium tuberculosis* (Mtb), the tuberculosis-provoking bacteria. Around 5–10% of these latent cases show symptoms of active TB (Sterling et al., 2020).

In TB contact investigations, household contacts are most likely infected by people with active TB (index case) (Lee, 2016; Fox et al., 2013). Several studies of tuberculosis household contact showed that some contacts remained negative based on TB infection tests, even after prolonged exposure (Fox et al., 2013; Reichler et al., 2018). These observations raise the question of whether some people might be able to withstand the Mtb infection or quickly clear the infection. Learning more about who these people are and what immunological or genetic factors allow them to fight infection may help develop better biomedical prevention and treatment tools, including a TB vaccine.

Some studies have suggested a link between the genetic make-up of the host and the prevalence of tuberculosis (Qidwai et al., 2012; van Tong et al., 2017), one of which is the *Vitamin D Receptor* (VDR) gene (Xu and Shen, 2019; Cao et al., 2016; Huang et al., 2015). The VDR gene is of great interest since it mediates the effects of vitamin D, which has been known to modulate the activity of monocytes-macrophages in the body, which are important in innate defense against infectious pathogens such as Mtb (Henrique et al., 2017; Ashenafi et al., 2018). In numerous studies, vitamin D shortage is associated with TB susceptibility (Tessema et al., 2019; Huang et al., 2017; Junaid et al., 2016). However, some reports did not find a significant discrepancy in vitamin D quantity in TB and non-TB individuals (Ashenafi et al., 2018; Koo et al., 2012; Yuvaraj et al., 2016; Sarin et al., 2016). In addition, randomized controlled trials have not found a significant effect on vitamin D3 supplementation to increase the rate of acid-fast bacilli conversion in TB patients (Ganmaa et al., 2017; Tukvadze et al., 2015; Bekele et al., 2018) or reduce the risk of TB infection (Ganmaa et al., 2020). Intervention with vitamin D3 appears to be modulated by genetic variants in VDR (Ganmaa et al., 2017). Consequently, the action of vitamin D may be contingent on the genotype and activity of the VDR (Sutaria et al., 2014).

The binding of VDR with calcitriol, vitamin D active form, induces the synthesis of cathelicidin (LL37). LL37 plays an essential role in mycobactericidal activity (Coussens et al., 2015; Belyaeva et al., 2017; Sonawane, 2011) and is involved in the autophagy process by enhancing the fusion of mycobacterial phagosomes with lysosomes (Chung et al., 2020; Choi et al., 2012; Paik et al., 2019). The binding of the calcitriol to the intracellular VDR also modulates approximately 900 genes involved in several physiological processes in both natural and adaptive immunity (Bizzaro et al., 2017). VDR helps to regulate the adaptive immune system by preventing over-response by suppressing lymphocyte proliferation and inhibiting the generation of pro-inflammatory cytokines (Sutaria et al., 2014). The researchers identified genetic variants of the VDR gene as being linked to a risk of TB based on the idea that genetic variations may cause anomalies and influence the immunological function of the VDR gene (Sutaria et al., 2013).

Research so far has been investigating the association between VDR polymorphisms with the development of active TB with varied results (Junaid et al., 2016; Medapati et al., 2017; Salimi et al., 2015; Mahmoud and Ali, 2014; Silva-Ramírez et al., 2019). There is still a lack of studies

about VDR gene expression in those infected with Mtb. The purpose of this study was to investigate the association between gene expression of VDR and vulnerability to TB. In this study, we analyze the gene expression of VDR in TB patients compared to controls who were known to be exposed to the Mtb bacilli, in this case, household contacts (HC). This study also aimed to compare VDR gene expression between positive and negative IGRA (Interferon-Gamma Release Assay) HC. IGRA was performed in household contacts to determine the response of cellular immunity (T cells) to TB infection (Trajman et al., 2013; Eom et al., 2018). In addition, the 3' UTR (untranslated region) VDR polymorphism, rs11574113, was also identified to assess its association with VDR gene expression as well as susceptibility to TB. The SNP (single nucleotide polymorphism) rs11574113 G > C is located at intron 8 and has been investigated in only a few studies regarding its association with tuberculosis (Andraos, 2011; Hu et al., 2016).

## 2. Material and methods

### 2.1. Research design and sample recruitment

The study design was cross-sectional, involving PTB patients and their HC as participants. We recruited the PTB patients from The Community Lung Health Center or Balai Besar Kesehatan Paru Masyarakat (BBKPM) Makassar, South Sulawesi, Indonesia, which is one of the main TB referral health centers in South Sulawesi Province. PTB patients included in this study were pulmonary TB patients aged more than 15 years old that diagnosed with active TB for the first time based on clinical manifestations, chest X-ray, and positive microscopic smears, which were further confirmed through TB culture. The exclusion criteria were HIV-positive patients which were examined by using SD Bioline. In this study, the household contacts were contacts aged 15 and above who resided for at least six months in the same residence as TB patients (Halliday et al., 2017), had no clinical symptoms of TB, and never had anti-tuberculosis drug therapy. All participants have signed the consent form.

We collected blood samples from PTB and HC. Before RNA extraction, the blood samples were kept at  $-80^{\circ}\text{C}$ . In order to test for latent TB infection in HC, QuantiFERON Gold Plus TB Test (Qiagen, Germany) was utilized, according to the manufacturer guide (Qiagen, 2016). The positive sputum samples of PTB patients were decontaminated and a culture process proceeded as a gold standard of TB diagnosis in the Hasanuddin University Medical Research Center Laboratory in Makassar, Indonesia.

This research has been approved by The Committee of Research Ethics, Faculty of Medicine, Universitas Hasanuddin, Makassar, South Sulawesi, Indonesia (No. 583/H4.8.4.5.31/PP36-KOMETIK/2018). All study participants gave their written consent.

### 2.2. VDR gene expression by real-time PCR

#### 2.2.1. RNA extraction and complementary DNA synthesis

Total RNA was extracted from whole blood samples, using Tiangen RNAprep Pure Blood Kit for purification of total RNA from human whole blood Cat. No. 4992238 (Tiangen, Biotech, Beijing, China). Concentrated RNA was rinsed with 50  $\mu\text{l}$  RNase free  $\text{H}_2\text{O}$  and utilized to complementary DNA (cDNA).

cDNA has been synthesized by using the iScript<sup>TM</sup> cDNA Synthesis Kit Master Cat. No. 178890 (Bio-Rad, California, USA). The reaction mixture consisted of 5  $\mu\text{g}$  of the total RNA, 1  $\mu\text{l}$  iScript Reverse Transcriptase, 4  $\mu\text{l}$  5 $\times$  the iScript reaction mixture, and nuclease-free water up to 25  $\mu\text{l}$ . The reaction mixture was subsequently incubated in a thermal cycler (Bio-Rad, USA), with 5-minute priming at  $25^{\circ}\text{C}$ , 20 min of reverse transcription (RT), and  $95^{\circ}\text{C}$  for a minute of RT inactivation. Until utilized for real-time PCR, cDNA was kept at  $-20^{\circ}\text{C}$ .

### 2.2.2. VDR gene expression by real-time quantitative polymerase chain reaction

The determination of the strength of gene expression (upregulation or downregulation) of VDR used SsoFast™ EvaGreen® Supermix Cat. No. 172–5200 (Bio-Rad, California, USA), and *glyceraldehyde-3 phosphate dehydrogenase (GAPDH)* as the housekeeping gene with quantitative real-time PCR (qPCR) method. Prior to amplification, the master mix was prepared by mixing 5 µl of SsoFast™ EvaGreen® Supermix, 1 µl of forward primer, 1 µl of reverse primer (each 10 nM primary concentration), the volume of cDNA corresponds to the cDNA concentration of 150 ng/µl, and the volume of nuclease-free water adjusted to reach 3 µl of the cDNA and nuclease-free water mixture. The primers used in this study were F: 5'-CGC ATG ATT GCC ATA CTG CTG G-3' and R: 5'-CCA CCA TCA TTC ACA CGA ACT GG-3' (Salehi-tabar et al., 2018) to amplify the VDR gene (NCBI (The National Center for Biotechnology Information) Reference sequence: NM\_000376) and F: 5'-CCT GCA CCA CCA ACT GCC TTA-3' and R: 5'-GGC CAT CCA CAG TCT TCT GA-3' to amplify GAPDH gene. The real-time PCR engine used was the CFX96 Touch Real-time PCR Detection System (Bio-Rad, California, USA) which is connected to the CFX Manager™ Software # 1845000 for Windows, with qPCR cycles: initial denaturation at 95 °C for 30 s, followed by 45 cycles of denaturation at 95 °C for 5 s, annealing at 57 °C for 30 s, and plate reading. Melt-curve was generated by increasing the temperature from 65.0 to 95.0 °C with increment 0.5 °C/5 s, followed by plate reading.

All analyses were performed by a real-time PCR method using Bio-Rad CFX™ Manager Software (version 3.1, Bio-Rad Lab. Inc., CA, USA). The level of gene expression was determined by the appearance of the amplicon signal on the graph. The signal detected in the small number of cycles indicates a higher level of gene expression. After the amplification process, a dissociation curve was obtained, which then analyzed for its relative expression using a method that compares the Ct (cycle threshold) value of the VDR gene ( $C_{t_{\text{target genes}}}$ ) with the selected reference value, i.e. the expression level of the GAPDH ( $C_{t_{\text{ref}}}$ ), by calculating:

$$\Delta Ct = C_{t_{\text{target genes}}} - C_{t_{\text{ref}}}$$

Remarks:

$\Delta Ct$ : The distinction between the VDR Ct and the GAPDH Ct value.

$C_{t_{\text{target genes}}}$ : Ct value of VDR gene.

$C_{t_{\text{ref}}}$ : Ct value of GAPDH gene.

Then a comparison was obtained with the formula of the gene expression:

$$\text{Relative VDR gene expression} = 2^{-\Delta\Delta Ct}$$

Remarks:

Relative VDR gene expression: VDR gene expression of the target to control group ratio.

$\Delta\Delta Ct$ : The difference between  $\Delta Ct$  of control and target group (Livak and Schmittgen, 2001).

### 2.3. VDR polymorphism (SNP rs11574113)

#### 2.3.1. DNA extraction

A total of 34 µl of blood samples (plasma and buffy coat) were put into a sterile 1.5 ml microcentrifuge tube. 20 µl of Proteinase K was added subsequently homogenized by pipetting, then incubated at 60 °C for 5 min. DNA extraction was performed using the gSYNCTM DNA Extraction Kit (Geneaid, Taiwan) according to the kit protocol. Extraction results were stored at -80 °C until used as a PCR template.

#### 2.3.2. DNA amplification and sequencing

Primers used to amplify a 500 bp of the 3' UTR of VDR (SNP rs11574113) were F 5'-CAG AGC ATG GAC AGG GAG CAA-3' and R 5'-ACT TCG AGC ACA AGG GGC GTT AG-3'. The PCR reaction mixture

consisted of 27 µl of the Taq Polymerase Kapa Biosystem Enzyme (Roche, USA), 1 µl of forward and reverse primers with a concentration of 10 M, 7–10 µl DNA samples, nuclease-free water until the total reaction mixture reached 50 µl. Amplification was carried out in a thermal cycler machine (Bio-Rad, USA) with the PCR condition: initial denaturation at 94 °C for 5 min, followed by 35 cycles of denaturation at 94 °C for 30 s, annealing at 50 °C for 30 s, extension at 72 °C for a minute, and final extension at 72 °C for 5 min. The PCR product was then electrophoresed on GelRed® (Biotium, Inc., Fremont, U.S.)-stained 2% Agarose gel with an electrophoresis machine (Bio-Rad, USA) then visualized under UV light using the Gel Doc™ XR machine (Bio-Rad, USA) and Quantity One software version 4.5 (Bio-Rad, USA).

The PCR product was then sent to the 1st Base Laboratory, Malaysia for direct sequencing (Sanger Sequencing). The sequencing results obtained were subsequently analyzed using Bioedit for Windows software version 7.2 (Tom Hall) with NCBI RefSeqGene: NG\_008731.1 (*Homo sapiens vitamin D receptor (VDR)*, RefSeqGene on chromosome 12) as reference gene.

### 2.4. Statistical analysis

Statistical analyses were performed using the IBM SPSS Statistics for Windows, Version 25.0 (IBM Corp. Armonk, NY, USA). The chi-square test was used to determine differences in nominal data between groups to compare parametric data between two groups and three groups. t-test and One-way ANOVA were used, respectively. Meanwhile, the Mann-Whitney test and Kruskal–Wallis test were used to compare non-parametric data between the groups, respectively. qPCR data were evaluated in real-time with the software Bio-Rad CFX™ Manager (version 3.1, Bio-Rad Lab. Inc., Hercules, CA, USA). Visualization of sequencing chromatogram and alignment to reference sequence (*Homo sapiens vitamin D receptor (VDR)*, RefSeqGene on chromosome 12 NCBI RefSeqGene: NG\_008731.1:g.64915G > C) was performed by using Bioedit 7.2.5 (Alzohairy, 2011; H91999). SNPstats web-based software (<https://www.snpsstats.net>) was used to analyze the Hardy-Weinberg equilibrium and association with active TB and positivity of IGRA, which was performed using a logistic regression model corrected for age, sex, and smoking status (Solé et al., 2006). A generalized linear model was used to examine the impact of factors to contribute to the incidence of PTB and IGRA positivity. Besides, it was also used to estimate the relationship between VDR polymorphism (SNP rs11574113) and VDR gene expression. In all studies,  $p$ -value  $\leq 0.05$  was statistically significant.

## 3. Result

### 3.1. Participants characterization

The characterization of the study participants is described in Table 1. In this study, positive IGRA was found in 61.64% (45/73) of household contacts. The age category across the groups did not differ ( $p = 0.281$ ). Male was significantly dominant in TB group (53%) than in HC group, 9th positive (31.1%) and negative (17.9%) IGRA ( $p = 0.001$ ). However, there was no substantial difference in gender between positive and negative IGRA HC ( $p = 0.21$ ). In addition, the acid fast-bacilli of index cases was not significantly different in positive IGRA HC compared to negative IGRA HC. There were significant differences in BMI between all three groups ( $p < 0.0001$ ), as well as between positive and negative IGRA HC ( $p = 0.0018$ ). Underweight participants were more common in the TB group (48.1%), and obese participants were more in the negative IGRA HC group (59.3%). Spouses were found more in positive IGRA HC than other relationships (62.2%).

**Table 1**  
Samples characterization.

	PTB N = 83	(+) IGRA HC N = 45	(-) IGRA HC N = 28	p-Value <sup>†</sup>	p- Value <sup>‡</sup>
Gender				<b>0.001*</b>	0.21
• Male	44 (53%)	14 (31.1%)	5 (17.9%)		
• Female	39 (47%)	31 (68.9%)	23 (82.1%)		
Age (years)				0.281	0.325
• 15–39	36 (43.4%)	22 (48.9%)	17 (60.7%)		
• ≥40	47 (56.6%)	23 (51.1%)	11 (39.3%)		
BMI classification (kg/m <sup>2</sup> )				<b>&lt;0.0001*</b>	<b>0.018*</b>
• Underweight ( $<18.5$ )	39 (48.1%)	6 (13.3%)	2 (7.4%) 6		
• Normal weight (18.5–22.9)	35 (43.2%)	19 (42.2%)	10 (35.7%)		
• Overweight (23–24.9)	3 (3.7%)	10 (22.2%)	16 (59.3%)		
• Obese ( $\geq 25$ )	4 (4.9%)	10 (22.2%)	16 (59.3%)		
Smoker	43 (51.8%)	11 (24.4%)	4 (14.3%)	<b>&lt;0.0001*</b>	0.296
Smear microscopic (PTB only)	37 (44.6%)				
• AFB 1+	35 (42.2%)				
• AFB 2+	11 (13.3%)				
• AFB 3+					
Smear microscopic of the index case (HC only)	25 (55.6%)	13 (28.9%)	13 (46.4%)		0.46
• AFB 1+	13 (28.9%)	12 (42.9%)	7 (24.4%)		
• AFB 2+	7 (15.6%)	3 (10.7%)	3 (10.7%)		
• AFB 3+					
Relationship to index cases (HC only)	11 (24.4%)	9 (32.1%)	9 (32.1%)		<b>0.025*</b>
• Parent/son/ daughter	28 (62.2%)	9 (32.1%)	9 (32.1%)		
• Spouse	6 (13.3%)	10 (44.4%)	10 (35.7%)		
• Others					

Remarks: <sup>†</sup> Chi-square test (among three groups); <sup>‡</sup> Chi-square test between (+) IGRA HC and (-) IGRA HC; \* significant p-value:  $p < 0.05$  (bold).  
PTB: new-active pulmonary tuberculosis patient; HC: household contact; (+) IGRA: positive IGRA; (-) IGRA: negative IGRA; BMI: body mass index.

**3.2. VDR gene expression in pulmonary tuberculosis patients and household contacts**

Based on the qPCR results (Table 2), VDR gene expression in negative IGRA HC was around five times higher than in PTB ( $p = 0.014$ ), and VDR gene expression in positive IGRA HC was approximately four times higher than in PTB ( $p = 0.011$ ). These results indicated that VDR gene expression in PTB was 0.2 (1:5) and 0.25 (1:4) downregulated compared to negative IGRA and positive IGRA HC, respectively. VDR gene expression in negative IGRA HC was higher than those in positive IGRA HC, but not significant ( $p = 0.945$ ).

**3.3. VDR SNP rs11574113 analysis**

The sequencing results of the VDR SNP rs11574113 are shown in Supplementary files: Fig. 1. The results obtained conform to the Hardy-Weinberg equilibrium. The G/G genotype dominated the samples in this study, both in the PTB group (66/83 [79.5%]) and household contacts (59/73 [80.8%]). Meanwhile, the proportion of genotype C/C was the

**Table 2**  
Comparison of relative quantitative VDR gene expression between active pulmonary TB, positive IGRA, and negative IGRA household contacts.

Biological Group	Expression (95% CI)	p- Value	p-Value ANOVA <sup>†</sup>
(-) IGRA HC	5.067 (1.149–22.339) <sup>a</sup>	<b>0.014*</b>	<b>0.013*</b>
(+) IGRA HC	3.994 (1.443–11.051) <sup>a</sup>	<b>0.011*</b>	
PTB	1	–	
(-) IGRA HC / (+) IGRA HC	1.269 (0.219–7.339) <sup>b</sup>	0.945	

Remarks: <sup>†</sup> the significance between three groups. \* significant p-value:  $p < 0.05$  (bold).

PTB: new-active pulmonary tuberculosis patient; HC: household contact; (+) IGRA: positive IGRA; (-) IGRA: negative IGRA.

<sup>a</sup> PTB as reference.

<sup>b</sup> (+) IGRA HC as the reference.

least in this study and was detected in only two samples of participants. In addition, it was found in the PTB group only (2/83 [127]), thus the value of OR of this genotype was unable to determine. The distribution of the genotype of the VDR SNP rs11574113 and its relationship to the incidence of active pulmonary tuberculosis and IGRA positivity are shown in Table 3 and Table 4.

The VDR gene expression of each genotype can be seen in Fig. 1. The VDR gene expression of the G/G genotype group was significantly different from that of G/C ( $p = 0.782$ ). However, when compared with the C/C genotype group, the expression of the VDR gene was significantly higher in the G/G and G/C genotype groups ( $p = 0.022$  and  $p = 0.03$ , respectively). By using generalized linear model analysis, G/G genotype and G/C genotype were associated with higher VDR gene expression compared to C/C genotype (OR<sup>a</sup> = 9.754 95% CI 1.482–64.195,  $p = 0.018$  and OR<sup>b</sup> = 9.723 95% CI 1.31–66.751,  $p = 0.021$ , respectively). The differences in the expression of the VDR gene by clustering the genotype VDR SNP rs11574113 in each sample group are detailed in Fig. 2. In the PTB group, the VDR gene expression was lower in the C/C genotype group ( $p = 0.037$ ), while in the IGRA negative HC group the VDR gene expression was significantly higher in the G/C group ( $p = 0.042$ ).

**3.4. Association of VDR gene expression with active pulmonary tuberculosis and IGRA positivity**

The VDR genes expression was found to be associated with active pulmonary tuberculosis after adjustments to other variables (OR<sup>a</sup> =

**Table 3**  
Genotype distribution of VDR SNP rs11574113 and its association to active pulmonary tuberculosis.

Model	Genotype	PTB N = 83	Household contact N = 73	OR 95% CI	p- Value
Codominant	G/G	66 (79.5%)	59 (80.8%)	1	0.2
	G/C	15 (18.1%)	14 (19.2%)	0.76 (0.29–2.02)	
	C/C	2 (2.4%)	0 (0%)	NA (0.00–NA)	
Dominant	G/G	66 (79.5%)	59 (80.8%)	1	0.81
	G/C-C/C	17 (20.5%)	14 (19.2%)	0.89 (0.34–2.31)	
Recessive	G-G-G/C	81 (97.6%)	73 (100%)	1	0.09
	C/C	2 (2.4%)	0 (0%)	NA (0.00–NA)	

Remarks: PTB: new-active pulmonary tuberculosis patient. OR: odds ratio.

**Table 4**  
Genotype distribution of VDR SNP rs11574113 and its association to active pulmonary tuberculosis.

Model	Genotype	(+) IGRA HC N = 45	(-) IGRA HC N = 28	OR 95% CI	p-Value
-	G/G	35 (77.8%)	24 (85.7%)	1.56 (0.39-6.19)	0.53
	G/C	10 (22.2%)	4 (14.3%)		

Remarks: HC: household contact; (+) IGRA: positive IGRA; (-) IGRA: negative IGRA; OR: odds ratio.

0.736 95% CI 0.542–0.998,  $p = 0.049$ ), as shown in Table 5. However, the association between VDR gene expression and IGRA positivity was not found significant in this study ( $OR^a = 0.783$  95% CI 0.539–1.137,  $p = 0.198$ ), as shown in Table 6.

**4. Discussion**

Early case detection and prevention of new infections are crucial in TB eradication efforts. Household contacts of tuberculosis patients are one group that has a high risk of suffering from tuberculosis (Fox et al., 2013). Household contact tracing of tuberculosis patients proved to be a better tool in detecting new tuberculosis cases than passive-case finding alone (Fox et al., 2018). Previous studies have revealed that some of these household contacts who had been exposed to *Mtb* for a long time did not experience infection, and the tuberculosis infection test results were still negative (Fox et al., 2013; Reichler et al., 2018). Several factors including genetic and immunological factors were investigated concerning an individual's resistance to tuberculosis infection (Allen et al., 2010; BoseDasgupta and Pieters, 2014; Wu et al., 2019; Harishankar et al., 2018).


In this study, VDR gene expression of PTB patients was significantly lower when compared to positive and negative IGRA HC, as shown in Fig. 1. This finding was similar to Panda's study, which also demonstrated the lower VDR expression in the active TB patient group than in household contacts with a negative tuberculin skin test. This result was

presumably due to *Mtb*'s efforts to decrease VDR gene expression for its benefit, to prevent the host's immune system and finally remain in the host (Mangin et al., 2014). Supporting this hypothesis, a study by Padhi et al. demonstrated that one of the *Mtb* lipoproteins, LprE, may inhibit CYP27B1 and VDR molecules to support intracellular *Mtb* survival (Padhi et al., 2019). When bacterial ligands downregulate VDR, the receptor is unable to express the enzymes essential to keep calcitriol in the normal range, and high calcitriol levels may occur (Miao and Goltzman, 2021). Increased calcitriol lowers VDR competency, suppresses macrophage activity, and may block the nuclear factor kappa-pat (Mangin et al., 2014).

In line with the results of this study, a previous microarray-based study demonstrated that VDR expression in *Mtb*-stimulated macrophages (U937) was downregulated (Xu et al., 2003). Contrary, a subsequent study that examined VDR mRNA expression in macrophages of TB patients stimulated with *Mtb*, showed an increase in VDR mRNA expression (Selvaraj et al., 2009; Fiske et al., 2019). This contrasting result may be due to the differences in the macrophage model used. Xu et al. used the U937 macrophage model (cell line) (Xu et al., 2003), while Selvaraj et al. and Fiske et al. used macrophages from TB patients, which demonstrated "trained" immunity from previous *Mtb* exposure, whereas macrophages from these patients remained "naïve" and responded quickly after re-stimulation with *Mtb* (Selvaraj et al., 2009; Fiske et al., 2019).

In addition to the efforts of *Mtb* itself, the low expression of the VDR gene experienced by TB patients themselves may lead to disruption of their immunity to be susceptible to tuberculosis. A decrease in the mRNA level of the VDR gene in TB cases indicates a disturbance in the activation process of inflammatory cytokines and innate immune cells, thereby reducing immunity to TB disease (Maruthai et al., 2020). Thus, the low expression of the VDR gene in TB patients may be a risk factor for TB infection as a result of *Mtb* infection, as some experts believe (Mangin et al., 2014).

The expression of the VDR gene for positive IGRA HC was found to be lower than that of negative IGRA HC but did not reach statistical significance (Table 3). To the best of our knowledge, this was the first study conducted to investigate the difference of the VDR expression gene



**Fig. 1.** VDR gene expression in each genotype of VDR SNP rs11574113. Sign\* indicates a significant difference ( $p < 0.05$ ) of VDR gene expression between two groups calculated by the Mann-Whitney test.




Fig. 2. VDR gene expression in each genotype of VDR SNP rs11574113 (clustered by the group). Sign\* indicates a significant difference ( $p < 0.05$ ) of VDR gene expression between two groups and three groups calculated by Mann-Whitney and Kruskal-Wallis test.

**Table 5**  
Association of variables and active pulmonary tuberculosis in multivariate analysis.

Variable	OR <sup>a</sup> (95% CI)	p-Value <sup>b</sup>
Age	1.022 (0.993–1.051)	0.142
Gender		
Male	0.97 (0.342–2.749)	0.954
Female	1	
BMI	<b>0.745 (0.664–0.837)</b>	<b>&lt;0.0001*</b>
Smoker		
No	<b>0.323 (0.112–0.933)</b>	<b>0.037*</b>
Yes	1	
Relative VDR gene expression (Log)	<b>0.736 (0.542–0.998)</b>	<b>0.049*</b>
VDR SNP rs11574113		
G/G	1.155 (0.437–3.051)	0.772
G/C/C/C	1	

Remarks: BMI: body mass index; VDR: Vitamin D Receptor; OR: odds ratio.

<sup>a</sup> Significant p-value :  $p \leq 0.05$  (bold).

<sup>b</sup> Analysis was adjusted to age, sex, BMI, smoking history, relative VDR gene expression (Log) and VDR SNP rs11574113 by using household contacts as a reference category.

between positive and negative IGRA people. IGRA-positive results in household contacts indicated that they might have *Mtb* infections, which could be the reason for lower VDR gene expression than negative IGRA household contacts, which may result in low levels of cathelicidin that are not sufficient to eliminate TB completely but may still be able to reduce excessive multiplication. Thus, its expression may be relatively higher when compared to active TB patients but lower when compared to IGRA-positive household contacts.

In this study, positive IGRA was found in 61.64% (45/73) of household contacts. This prevalence was almost similar to the study in Uganda (68.42%), also included in the list of 30 highly endemic nations with TB/HIV (Jones-López et al., 2013). In comparison to previous research in South Korea and Mongolia, however, the prevalence of

**Table 6**  
Association of variables and IGRA positivity of household contacts in multivariate analysis.

Variable	OR <sup>a</sup> (95% CI)	p-Value <sup>b</sup>
Age	1.001 (0.955–1.049)	0.97
Gender		
Male	1.336 (0.272–6.571)	0.772
Female	1	
BMI	<b>0.831 (0.733–0.942)</b>	<b>0.004*</b>
Smoker		
No	0.466 (0.084–3.106)	0.466
Yes	1	
Smear microscopic of the index case		
AFB 1+	0.538 (0.087–3.348)	0.507
AFB 2+	0.399 (0.061–2.593)	0.336
AFB 3+	1	
Relation to the index case		
Parent/son/daughter	2.157 (0.451–10.322)	0.336
Spouse	<b>9.201 (1.757–48.192)</b>	<b>0.009*</b>
Others	1	
Relative VDR gene expression (Log)	0.783 (0.539–1.137)	0.198
VDR SNP rs11574113		
G/G	0.564 (0.122–2.62)	<b>0.465</b>
G/C/C/C	1	

<sup>a</sup> Remarks: BMI: Body mass index; AFB: acid-fast bacilli; VDR: Vitamin D Receptor; OR: odds ratio.

<sup>b</sup> Significant p-value :  $p < 0.05$  (bold).

<sup>c</sup> Analysis was adjusted to age, sex, BMI, smoking history, AFB of the index case, relation to the index case, relative VDR gene expression (Log) and VDR SNP rs11574113 by using negative IGRA household contacts as a reference category.

positive IGRA in our study was relatively high (47.7% and 48.2%, respectively) (Eom et al., 2018; Gurjav et al., 2019). The difference in the prevalence of positive IGRA among household contacts may be due to several things, including differences in the character of the recruited contact sample, such as how close the contacts are to the index case, the immunity status of the contact, and other factors such as the infectious

rate of the index case (Eom et al., 2018). In this study, most of the household contacts recruited were spouses (50.69% (37/73)) of positive AFB-index cases, and 75.67% (28/37) of them have positive IGRA results. Among the 45 positive IGRA HC, 28 (62.2%) were spouses. This study represented an immense rate of TB infection in the spouse of index cases. This was consistent with the investigation by Crampin et al., which revealed that the couples are in close contact having significant TB risks (Crampin et al., 2011).

A positive IGRA suggests the synthesis in the region of the Difference1 segment (RD1) in response to two *Mtb*-specific antigens, namely early secretory target-6 (ESAT-6), and cuticle filtrate protein 10 (CFP-10) (Zellweger et al., 2020). Currently, it is used to detect TB infection and is included in the diagnostic pathway for latent TB infection (Chapman and Lauzardo, 2014; Kementerian Kesehatan Republik Indonesia, 2020). IGRA is not recommended as a diagnostic tool to differentiate active and latent TB (Zellweger et al., 2020). Therefore, in this study, the IGRA was carried out on household contacts who did not have any clinical symptoms of TB and without a previous history of TB, so the IGRA here aims to determine whether the household contacts have been latently infected with TB or not. The immunological response in latent TB hosts can control the infection but is not sufficient to eradicate the mycobacterium (Gideon et al., 2011). The prevalent assumption is that the bacilli in latent TB stop growing and enter an immobile phase because of the "strong" immune response created by the host, eventually becoming non-replicable while maintaining the capacity to sleep further under advantageous conditions in granulomas (Gideon et al., 2011; Ehlers and Schaible, 2013; Rao et al., 2019).

Several studies suggest that *VDR* expression is also influenced by the *VDR* polymorphism (Ogunkolade et al., 2002; Decker and Parker, 1995; Jurutka et al., 2000; Panda et al., 2019), thus possibly contributing to the development of TB disease (Xu and Shen, 2019; Cao et al., 2016; Panda et al., 2019; Lee and Song, 2015). Current research show varying results in various populations and several meta-analyses commonly reported *VDR* polymorphism have attempted to conclude the association of *VDR* polymorphisms with the incidence of TB (Xu and Shen, 2019; Cao et al., 2016; Huang et al., 2015; Gao et al., 2010; Chen et al., 2013). In this study, we tried to analyze the polymorphisms in the 3' UTR *VDR*, SNP rs1157413 G > C, which has not been investigated much, especially in tuberculosis patients. Since it was located in intron 8, rs1157413 do not alter the sequence of *VDR* protein. The rs1157413 G/C variant has been reported to be associated with a reduced risk of colorectal cancer only in those with high plasma vitamin D concentrations. (Budhathoki et al., 2016). Meanwhile, in this study, the rs1157413 G/C variant had a significantly higher *VDR* gene expression than G/G only in the IGRA HC negative group (Fig. 2). Overall, this study observed significantly higher *VDR* gene expression in the G/G and G/C when compared to C/C genotype groups (Fig. 1). Besides, our study attempted to estimate the impact of SNP rs1157413 G/C variant on *VDR* gene expression by controlling for several variables and found that G/G genotype and G/C genotype were associated with higher *VDR* gene expression compared to C/C genotype. However, this needs to be investigated further considering that the C/C genotype in this study was only found in two samples and was limited to the PTB group. Furthermore, it is necessary to conduct in a larger size to confirm this finding, whether this SNP may have a direct effect on *VDR* expression or transcripts stability, or are in disequilibrium with other functional polymorphisms that regulate *VDR* activity, such as single(A) repeat in exon 9 (Whitfield et al., 2001) or affect transcript stability, such as the *VDR* *TaqI*, also located on exon 9 (Hussain et al., 2019). In another study of TB, it was found that the A allele of rs1157413 had a protective effect against TB (Hu et al., 2016), but in our study, we did not observe the presence of the A allele in any of the participants.

Several previous studies have studied the correlation of *VDR* gene expression with vitamin D levels in the blood. A recent randomized controlled trial of monozygotic twins studied increased *VDR* gene expression and 25-hydroxyvitamin D levels after 60 days of vitamin D

(cholecalciferol) supplementation. In addition, the study observed that increasing levels of 25-hydroxyvitamin to levels greater than 50 ng/mL contributed to increased expression of the *VDR* gene (Medc et al., 2020). A US cohort involving patients with sickle cell anemia found that decreased vitamin D levels were associated with decreased expression of the *VDR* gene (Han et al., 2018). Contrary to these findings, a study of patients with oral neoplasms in India revealed an increase in *VDR* expression but not significantly compared to controls, while 25-hydroxyvitamin D levels expressed in vitamin D scores were found to be significantly lower in the case group of oral neoplasms than controls (Anand et al., 2017). Although this study did not measure the correlation of *VDR* gene expression with 25-hydroxyvitamin D levels, it can be seen that there was an opposite effect between *VDR* gene expression and 25-hydroxyvitamin D levels. Meanwhile, there were also studies showing that there was no correlation between *VDR* gene expression with 25-hydroxyvitamin D levels but with CYP24A1/CYP27B1 expression in epilepsy patients (Mazdeh et al., 2018). This study indicates that there may be a different correlation between 25-hydroxyvitamin D levels and *VDR* gene expression in various diseases and populations, or indeed there may not be a simple correlation between 25-hydroxyvitamin D levels and *VDR* gene expression as with 1,25 dihydroxyvitamin D levels described previously (Ogunkolade et al., 2002).

As for tuberculosis, downregulation of *VDR* gene expression has been associated with changes in 1,25-dihydroxyvitamin D levels in a number of studies. Selvaraj et al. revealed that in TB patients, *VDR* gene expression was downregulated which was thought to be caused by high levels of 1,25-dihydroxyvitamin D (Selvaraj et al., 2009). On the other hand, other studies have shown that levels of 1,25-dihydroxyvitamin D increase *VDR* mRNA production, stabilize *VDR* mRNA or protect *VDR* degradation thereby increasing overall *VDR* counts (Kongsbak et al., 2013; Liu et al., 2006). 1,25-dihydroxyvitamin D plays a role in converting the *VDR* to a functionally active protein that can bind to RXR and to specific gene sequences and coregulators required for modulation of gene expression (Pike and Meyer, 2011; Pike and Meyer, 2012; Haussler and Whitfield, 2013). Therefore, the availability of 1,25-dihydroxyvitamin D is a prerequisite for *VDR* activity. During immune reactions, the most likely source of 1,25-dihydroxyvitamin D is primarily the endogenous production of its precursor molecule, 25-dihydroxyvitamin D. Several studies on immune cells have revealed that 25-hydroxyvitamin D can be utilized and subsequently converted to 1,25-dihydroxyvitamin D via the action of the enzyme CYP27B1 (Jeffery et al., 2012). The increased synthesis of 1,25-dihydroxyvitamin D under these circumstances may lead to the use of available 25-hydroxyvitamin D (Selvaraj et al., 2009). Because 1,25-dihydroxyvitamin D has a lower half-life and is used for various metabolic processes, active pulmonary TB patients may suffer from a lack of substrate for the synthesis of the active metabolite, including 25-hydroxyvitamin D (Selvaraj et al., 2009).

We tried to assess the association of *VDR* gene expression with active pulmonary tuberculosis and the outcome of positive IGRA. After adjustments to other variables, the multivariate analysis indicated an association between *VDR* gene expression and active TB susceptibility (Table 5). However, there was no association between the *VDR* gene expression and the IGRA positivity result in household contact even after adjusted with other variables (Table 6). The factors that showed an association with the positivity of IGRA were BMI and the relationship to the index case (Table 6). BMI also appears to be significantly associated with active TB susceptibility (Table 5). This result may suggest two possibilities. First, the greater the BMI, the lower the chance of contracting TB, which is consistent with prior findings (Kim et al., 2018; Kumaratne et al., 2017). A high BMI would seem to protect the immune system. Adipose tissue is known to secrete leptin and TNF, an inflammatory mediator that helps defend the host from TB infection (Dorhoi and Kaufmann, 2014). Second, TB infection itself may cause weight loss, thus decreasing nutritional status (Cegielski and McMurray, 2004).

There were several limitations in this study. We realize that the sample used in this study is not large, but we hope that the picture obtained in this study can be developed with a larger number of samples in the future to obtain more adequate conclusions. In our study, we tried to involve variables such as smoker status, the relationship of household contacts with index cases, and the intensity of AFB which has been associated with the spread of *Mtb* germs at home (Crampin et al., 2011; Guwatudde et al., 2003; Sepkowitz, 1996). However, we did not examine other factors such as blood vitamin D levels, dietary factors or vitamin D supplementation that might influence *VDR* gene expression. A more comprehensive study including other components involved in the vitamin D metabolic pathway would certainly provide more adequate results in assessing the role of *VDR* in TB susceptibility. Likewise, analyses involving interactions between genes or even genes and the environment will certainly provide a better understanding of genetic influences on individual susceptibility to TB.

To date, there is still a lack of studies investigating the expression of the *VDR* gene in people who are latently infected with TB. The literature has explained how the role of *VDR* in eliminating *Mtb* through the vitamin D-dependent pathway (Chung et al., 2020). Several previous studies have also reported lower vitamin D (25-hydroxyvitamin D) concentrations in LTBI individuals than in active TB patients (Hong et al., 2019; Esteve Palau et al., 2015). The present study found that the expression of the *VDR* gene in IGRA-positive household contacts was higher than in active TB, and it may reflect the *VDR* protective efforts against TB in household contacts at the time of the examination. Whether with adequate *VDR*, cathelicidin as a mycobactericidal is continuously produced in sufficient quantities, or *VDR* may be involved in maintaining *Mtb* in the non-replicative phase is still not clearly understood. The specific role of *VDR* involved in the latency phase needs further investigation.

## 5. Conclusion

To conclude, *VDR* gene expression may be associated with active pulmonary tuberculosis (PTB) but not with IGRA positivity in this study. The *VDR* gene expression was downregulated in PTB rather than positive and negative IGRA HC. The high expression of the *VDR* gene may be influenced by *VDR* SNP rs11574113. This result may suggest a protective role of *VDR* in the progression of PTB. This study also found that among spouses of IGRA-based PTB patients, in comparison with other family members spouses are at greater risk of PTB infection. Spouses of PTB patients need adequate protection so that existing *Mtb* infections will not progress to the active phase.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.genrep.2022.101581>.

## CRedit authorship contribution statement

**Najdah Hidayah:** Methodology, Software, Investigation, Writing - original draft preparation  
**Irawaty Djaharuddin:** Conceptualization, Writing - Reviewing and Editing, Supervision  
**Ahyar Ahmad:** Conceptualization, Validation, Writing - Reviewing and Editing, Supervision  
**Agussalim Bukhari:** Methodology, Supervision, Validation  
**Ilhamjaya Patellongi:** Methodology, Software, Validation  
**Nur Ahmad Tabri:** Investigation, Validation, Writing - Reviewing and Editing  
**Sana Agus:** Investigation, Validation, Supervision  
**Subair Subair:** Investigation, Writing - Reviewing and Editing  
**Irda Handayani:** Investigation, Writing - Reviewing and Editing  
**Andi Tenriola:** Investigation, Methodology, Software  
**Handayani Halik:** Methodology, Software, Validation  
**Muhammad Nasrum Massi:** Conceptualization, Validation, Writing - Reviewing and Editing, Supervision.

## Declaration of competing interest

None.

## Acknowledgement

The authors thank Integrated Master-Doctor Program (PMDSU) by the Indonesian Ministry of Research and Technology for funding this study (Grant No. 4514/UN4.21/PL.00.00/2018). A great appreciation was also expressed to Community Lung Health Center (BBKPM) Makassar for their support and assistance in the sample compilation. The authors also thank all of the study participants.

## References

- World Health Organization, 2020. Global Tuberculosis Report 2020. World Health Organization, Geneva.
- Dye, C., Glaziou, P., Floyd, K., Ravignone, M., 2013. Prospects for tuberculosis elimination. *Annu. Rev. Public Health* 34, 271–286. <https://doi.org/10.1146/annurev-publhealth-031912-114431>.
- Sterling, T.R., Njie, G., Zenner, D., et al., 2020. Guidelines for the treatment of latent tuberculosis infection: recommendations from the National Tuberculosis Controllers Association and CDC, 2020. *Am. J. Transplant.* 20, 1196–1206. <https://doi.org/10.1111/ajt.15841>.
- Lee, S.H., 2016. Tuberculosis infection and latent tuberculosis. *Tuberc. Respir. Dis. (Seoul)* 79, 201–206. <https://doi.org/10.4046/trd.2016.79.4.201>.
- Fox, G.J., Barry, S.E., Britton, W.J., Marks, G.B., 2013. Contact investigation for tuberculosis: a systematic review and meta-analysis. *Eur. Respir. J.* 41, 140–156. <https://doi.org/10.1183/09031936.00070812>.
- Reichler, M.R., Khan, A., Sterling, T.R., et al., 2018. Risk and timing of tuberculosis among close contacts of persons with infectious tuberculosis. *J. Infect. Dis.* 218, 1000–10008. <https://doi.org/10.1093/infdis/jiy265>.
- Qidwai, T., Jamal, F., Khan, M.Y., 2012. DNA sequence variation and regulation of genes involved in pathogenesis of pulmonary tuberculosis. *Scand. J. Immunol.* 75, 568–587. <https://doi.org/10.1111/j.1365-3083.2012.02696.x>.
- van Tong, H., Velavan, T.P., Thye, T., Meyer, C.G., 2017. Human genetic factors in tuberculosis: an update. *Trop. Med. Int. Health* 22, 1063–1071. <https://doi.org/10.1111/tmi.12923>.
- Xu, X., Shen, M., 2019. Associations between vitamin D receptor genetic variants and tuberculosis: a meta-analysis. *Innate Immun.* 25, 305–313. <https://doi.org/10.1177/1753425919842643>.
- Cao, Y., Wang, X., Cao, Z., Cheng, X., 2016. Vitamin D receptor gene FokI polymorphisms and tuberculosis susceptibility: a meta-analysis. *Arch. Med. Sci.* 12, 1118–1134. <https://doi.org/10.5114/aoms.2016.60092>.
- Huang, L., Liu, C., Liao, G., et al., 2015. Vitamin D receptor gene foki polymorphism contributes to increasing the risk of tuberculosis: an update meta-analysis. *Med. (United States)* 94, e2256. <https://doi.org/10.1097/MD.0000000000002256>.
- Henrique, P., Gois, F., Ferreira, D., et al., 2017. Vitamin D and infectious diseases: simple bystander or contributing factor? *Nutrients* 9, 651. <https://doi.org/10.3390/nu9070651>.
- Ashenafi, S., Mazurek, J., Rehn, A., et al., 2018. Vitamin D3 status and the association with human cathelicidin expression in patients with different clinical forms of active tuberculosis. *Nutrients* 10, 721. <https://doi.org/10.3390/nu10060721>.
- Tesema, B., Moges, F., Habte, D., et al., 2017. Vitamin D deficiency among smear positive pulmonary tuberculosis patients and their tuberculosis negative household contacts in Northwest Ethiopia: a case-control study. *Ann. Clin. Microbiol. Antimicrob.* 16, 36. <https://doi.org/10.1186/s12941-017-0211-3>.
- Huang, S.-J., Wang, X.-H., Liu, Z.-D., et al., 2017. Vitamin D deficiency and the risk of tuberculosis: a meta-analysis. *Drug Des. Devel. Ther.* 11, 91–102. <https://doi.org/10.2217/14796678.5.1.15>.
- Junaid, K., Rehman, A., Jolliffe, D.A., et al., 2016. Vitamin D deficiency associates with susceptibility to tuberculosis in Pakistan, but polymorphisms in *VDR*, *DBP* and *CYP2R1* do not. *BMC Pulm. Med.* 16, 73. <https://doi.org/10.1186/s12890-016-0240-2>.
- Koo, H.K., Lee, J.S., Jeong, Y.J., et al., 2012. Vitamin D deficiency and changes in serum vitamin D levels with treatment among tuberculosis patients in South Korea. *Respirology* 17, 808–813. <https://doi.org/10.1111/j.1440-1843.2012.02172.x>.
- Yuvaraj, B., Sridhar, M.G., Kumar, S.V., Kadhiravan, T., 2016. Association of serum vitamin D levels with bacterial load in pulmonary tuberculosis patients. *Tuberc. Respir. Dis. (Seoul)* 79, 153–157. <https://doi.org/10.4046/trd.2016.79.3.153>.
- Sarin, P., Duffy, J., Mughal, Z., et al., 2016. Vitamin D and tuberculosis: review and association in three rural provinces of Afghanistan. *Int. J. Tuberc. Lung Dis.* 20, 383–388. <https://doi.org/10.5588/ijtld.15.0303>.
- Ganmaa, D., Munkhzul, B., Fawzi, W., et al., 2017. High-dose vitamin D3 during tuberculosis treatment in Mongolia: a randomized controlled trial. *Am. J. Respir. Crit. Care Med.* 196, 628–637. <https://doi.org/10.1164/rccm.201705-0936OC>.
- Tukvadze, N., Sanikidze, E., Kipiani, M., et al., 2015. High-dose vitamin D3 in adults with pulmonary tuberculosis: a double-blind randomized controlled trial. *Am. J. Clin. Nutr.* 102, 1059–1069. <https://doi.org/10.3945/ajcn.115.113886>.
- Bekele, A., Gebreselassie, N., Ashenafi, S., et al., 2018. Daily adjunctive therapy with vitamin D3 and phenylbutyrate supports clinical recovery from pulmonary tuberculosis: a randomized controlled trial in Ethiopia. *J. Intern. Med.* 284, 292–306. <https://doi.org/10.1111/joim.12767>.
- Ganmaa, D., Uyanga, B., Zhou, X., et al., 2020. Vitamin D supplements for prevention of tuberculosis infection and disease. *N. Engl. J. Med.* 383, 359–368. <https://doi.org/10.1056/nejmoa1915176>.

- Sutaria, N., Liu, C.T., Chen, T.C., 2014. Vitamin D status, receptor gene polymorphisms, and supplementation on tuberculosis: a systematic review of case-control studies and randomized controlled trials. *J. Clin. Transl. Endocrinol.* 1, 151–160. <https://doi.org/10.1016/j.jcte.2014.08.001>.
- Coussens, A.K., Wilkinson, R.J., Martineau, A.R., 2015. Phenylbutyrate is bacteriostatic against *Mycobacterium tuberculosis* and regulates the macrophage response to infection, synergistically with 25-hydroxy-vitamin D. *PLoS Pathog.* 11, e1005007. <https://doi.org/10.1371/journal.ppat.1005007>.
- Belyaeva, L.V., Churilov, L.P., Robertova, L., Ms, M., 2017. Vitamin D, cathelicidin, prolactin, autoantibodies, and cytokines in different forms of pulmonary tuberculosis versus sarcoidosis. *Isr. Med. Assoc. J.* 19, 499–505.
- Solé, X., Guinó, E., Valls, J., et al., 2006. SNPStats: A web tool for the analysis of association studies. *Bioinformatics* 22, 1928–1929. <https://doi.org/10.1093/bioinformatics/btl268>.
- Sonawane, A., Santos, J.C., Mishra, B.B., et al., 2011. Cathelicidin is involved in the intracellular killing of mycobacteria in macrophages. *Cell. Microbiol.* 13, 1601–1617. <https://doi.org/10.1111/j.1462-5822.2011.01644.x>.
- Chung, C., Silwal, P., Kim, I., et al., 2020. Vitamin D-cathelicidin axis: at the crossroads between protective immunity and pathological inflammation during infection. *Immune Netw.* 20, e12. <https://doi.org/10.4110/in.2020.20.e12>.
- Choi, K.Y., Chow, L.N.Y., Mookherjee, N., 2012. Cationic host defence peptides: multifaceted role in immune modulation and inflammation. *J. Innate Immun.* 4, 361–370. <https://doi.org/10.1159/000336630>.
- Paik, S., Kim, J.K., Chung, C., Jo, E.K., 2019. Autophagy: a new strategy for host-directed therapy of tuberculosis. *Virulence* 10, 448–459. <https://doi.org/10.1080/21505594.2018.1536598>.
- Bizzaro, G., Antico, A., Fortunato, A., Bizzaro, N., 2017. Vitamin D and autoimmune diseases: is vitamin D receptor (VDR) polymorphism the culprit? *Isr. Med. Assoc. J.* 19, 438–443.
- Liu, J., Jiang, T., Wei, L., et al., 2013. The discovery and identification of a candidate proteomic biomarker of active tuberculosis. *BMC Infect. Dis.* 13, 506. <https://doi.org/10.1186/1471-2334-13-506>.
- Medapati, R.V., Suvvari, S., Godi, S., Gangiseti, P., 2017. NRAMP1 and VDR gene polymorphisms in susceptibility to pulmonary tuberculosis among Andhra Pradesh population in India: a case-control study. *BMC Pulm. Med.* 17, 89. <https://doi.org/10.1186/s12890-017-0431-5>.
- Salimi, S., Farajian-mashadi, F., Alavi-Naini, R., et al., 2015. Association between vitamin D receptor polymorphisms and haplotypes with pulmonary tuberculosis. *Biomed. Rep.* 3, 189–194. <https://doi.org/10.3892/br.2014.402>.
- Mahmoud, A.A., Ali, A.H.K., 2014. Vitamin D receptor gene polymorphism and 25 hydroxy vitamin D levels in Egyptian patients with pulmonary tuberculosis. *Egypt J. Chest Dis. Tuberc.* 63, 651–655. <https://doi.org/10.1016/j.ejcdt.2014.02.015>.
- Silva-Ramirez, B., Saenz-Saenz, C.A., Bracho-Vela, L.A., et al., 2019. Association between vitamin D receptor gene polymorphisms and pulmonary tuberculosis in a Mexican population. *Indian J. Tuberc.* 66, 70–75. <https://doi.org/10.1016/j.ijt.2018.04.005>.
- Trajman, A., Steffen, R.E., Menzies, D., 2013. Interferon-gamma release assays versus tuberculin skin testing for the diagnosis of latent tuberculosis infection: an overview of the evidence. *Pulm. Med.* 601737. <https://doi.org/10.1155/2013/601737>.
- Eom, J.S., Kim, I., Kim, W., et al., 2018. Household tuberculosis contact investigation in a tuberculosis-prevalent country. *Medicine (Baltimore)* 97, e9681. <https://doi.org/10.1097/MD.00000000000009681>.
- Andraos, C., 2011. CpG Islands of the Vitamin D Receptor Gene: Differential Methylation and Tuberculosis Predisposition. University of Johannesburg, South Africa.
- Hu, Q., Chen, Z., Liang, G., et al., 2016. Vitamin D receptor gene associations with pulmonary tuberculosis in a Tibetan Chinese population. *BMC Infect. Dis.* 1–7. <https://doi.org/10.1186/s12879-016-1699-4>.
- Halliday, A., Whitworth, H., Kottoor, S.H., et al., 2017. Stratification of latent mycobacterium tuberculosis infection by cellular immune profiling. *J. Infect. Dis.* 215, 1480–1487. <https://doi.org/10.1093/infdis/jix107>.
- Qiagen, 2016. QuantIFERON® - TB Gold Plus (QFT® - Plus) ELISA Package Insert 2. [www.quantiferon.com/imm/content/PLI%90AQFT/PLUS/2PK-Elisa/UK.pdf](http://www.quantiferon.com/imm/content/PLI%90AQFT/PLUS/2PK-Elisa/UK.pdf). (Accessed 20 September 2018).
- Salehi-tabar, R., Memari, B., Wong, H., et al., 2018. The tumor suppressor FBW7 and the vitamin D receptor are mutual cofactors. *Mol. Cancer Res.* 17, 709–719. <https://doi.org/10.1158/1541-7786.MCR-18-0991>.
- Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔCT</sup> method. *Methods* 25, 402–408. <https://doi.org/10.1006/meth.2001.1262>.
- Alzohairy, A.M., 2011. BioEdit: an important software for molecular biology. *GERF Bull. Biosci.* 2, 60–61.
- Hall, T., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* 95–98.
- Fox, G.J., Nhung, N.V., Sy, D.N., et al., 2018. Household-contact investigation for detection of tuberculosis in Vietnam. *N. Engl. J. Med.* 378, 221–229. <https://doi.org/10.1056/nejmoa1700209>.
- Allen, M., Bailey, C., Cahatol, I., et al., 2015. Mechanisms of control of mycobacterium tuberculosis by NK cells: role of glutathione. *Front. Immunol.* 6, 508. <https://doi.org/10.3389/fimmu.2015.00508>.
- BoseDasgupta, S., Pieters, J., 2014. Striking the right balance determines TB or not TB. *Front. Immunol.* 5, 455. <https://doi.org/10.3389/fimmu.2014.00455>.
- Wu, S., Wang, M.G., Wang, Y., He, J.Q., 2019. Polymorphisms of cytokine genes and tuberculosis in two independent studies. *Sci. Rep.* 9, 2507. <https://doi.org/10.1038/s41598-019-39249-4>.
- Harishankar, M., Selvaraj, P., Bethunaickan, R., 2018. Influence of genetic polymorphism towards pulmonary tuberculosis susceptibility. *Front. Med.* 5, 213. <https://doi.org/10.3389/fmed.2018.00213>.
- Mangin, M., Sinha, R., Fincher, K., 2014. Inflammation and vitamin D: the infection connection. *Inflamm. Res.* 63, 803–819. <https://doi.org/10.1007/s00011-014-0755-z>.
- Padhi, A., Pattanaik, K., Biswas, M., et al., 2019. Mycobacterium tuberculosis LprE suppresses TLR2-dependent cathelicidin and autophagy expression to enhance bacterial survival in macrophages. *J. Immunol.* 203, 2665–2678. <https://doi.org/10.4049/jimmunol.1801301>.
- Miao, D., Goltzman, D., 2021. Probing the scope and mechanisms of calcitriol actions using genetically modified mouse models. *JBM Plus* 5, e10434. <https://doi.org/10.1002/jbm4.10434>.
- Xu, Y., Xie, J., Li, Y., et al., 2003. Using a cDNA microarray to study cellular gene expression altered by *Mycobacterium tuberculosis*. *Chin. Med. J.* 116, 1070–1073.
- Selvaraj, P., Prabhu Anand, S., Harishankar, M., Alagarasu, K., 2009. Plasma 1,25 dihydroxy vitamin D3 level and expression of vitamin D receptor and cathelicidin in pulmonary tuberculosis. *J. Clin. Immunol.* 29, 470–478. <https://doi.org/10.1007/s10875-009-9277-9>.
- Fiske, C.T., Blackman, A., Maruri, F., et al., 2019. Increased vitamin D receptor expression from macrophages after stimulation with *M. tuberculosis* among persons who have recovered from extrapulmonary tuberculosis. *BMC Infect. Dis.* 19, 266. <https://doi.org/10.1186/s12879-019-3958-7>.
- Maruthai, K., Sankar, S., Subramanian, M., 2020. Methylation status of VDR gene and its association with vitamin D status and VDR gene expression in pediatric tuberculosis disease D status and VDR gene expression in pediatric tuberculosis. *Immunol. Investig.* 00, 1–16. <https://doi.org/10.1080/08820139.2020.1810702>.
- Jones-López, E.C., Namugga, O., Mumbowa, F., et al., 2013. Cough aerosols of mycobacterium tuberculosis predict new infection: a household contact study. *Am. J. Respir. Crit. Care Med.* 187, 1007–1015. <https://doi.org/10.1164/rccm.201208-1422OC>.
- Gurjav, U., Ankhbat, M., Ganbaatar, G., et al., 2019. Vitamin D deficiency is associated with tuberculosis infection among household contacts in Ulaanbaatar, Mongolia. *Int. J. Tuberc. Lung Dis.* 23, 919–923. <https://doi.org/10.5588/ijtld.19.0047>.
- Crampin, A.C., Kasimba, S., Mwaungulu, N.J., et al., 2011. Married to *M. tuberculosis*: risk of infection and disease in spouses of smear positive tuberculosis patients. *Trop. Med. Int. Health* 16, 811–818. <https://doi.org/10.1111/j.1365-3156.2011.02763.x>.
- Zellweger, J.P., Sotgiu, G., Corradi, M., Durando, P., 2020. The diagnosis of latent tuberculosis infection (LTBI): currently available tests, future developments, and perspectives to eliminate tuberculosis (tb). *Med. Lav.* 111, 170–183. <https://doi.org/10.23749/mdl.v11i13.9983>.
- Chapman, H.J., Lauzardo, M., 2014. Advances in diagnosis and treatment of latent tuberculosis infection. *J. Am. Board Fam. Med.* 27, 704–712. <https://doi.org/10.3122/jabfm.2014.05.140062>.
- Kementerian Kesehatan Republik Indonesia, 2020. Petunjuk Teknis Penanganan Infeksi Laten Tuberculosis (ILTb). Jakarta.
- Gideon, H.P., Flynn, J.L., Flynn, J.L., 2011. Latent tuberculosis: what the host “ sees ”? *Immunol. Res.* 50, 202–212. <https://doi.org/10.1007/s12026-011-8229-7>.
- Ehlers, S., Schaible, U.E., 2013. The granuloma in tuberculosis: dynamics of a host-pathogen collusion. *Front. Immunol.* 3, 411. <https://doi.org/10.3389/fimmu.2012.00411>.
- Rao, M., Ippolito, G., Mfinanga, S., et al., 2019. Latent TB infection (LTBI) – mycobacterium tuberculosis pathogenesis and the dynamics of the granuloma battleground. *Int. J. Infect. Dis.* 80, S58–S61. <https://doi.org/10.1016/j.ijid.2019.02.035>.
- Ogunkolade, B., Boucher, B., Prah, J., et al., 2002. Vitamin D receptor gene polymorphisms influence insulin secretion in Bangladeshi Asians. *Diabetes* 51, 2291–2300. <https://doi.org/10.2337/diabetes.47.4.688>.
- Decker, C.L., Parker, R., 1995. Diversity of cytoplasmic functions for the 3' untranslated region of eukaryotic transcripts. *Curr. Opin. Cell Biol.* 7, 386–392. [https://doi.org/10.1016/0955-0674\(95\)80094-8](https://doi.org/10.1016/0955-0674(95)80094-8).
- Jurutka, P.W., Remus, L.S., Whitfield, G.K., et al., 2000. The polymorphic N terminus in human vitamin D receptor isoforms influences transcriptional activity by modulating interaction with transcription factor IIB. *Mol. Endocrinol.* 14, 401–420. <https://doi.org/10.1210/mend.14.3.0435>.
- Panda, S., Tiwari, A., Luthra, K., et al., 2019. Association of FokI VDR polymorphism with vitamin D and its associated molecules in pulmonary tuberculosis patients and their household contacts. *Sci. Rep.* 9, 15251. <https://doi.org/10.1038/s41598-019-51803-8>.
- Lee, Y.H., Song, G.G., 2015. Vitamin D receptor gene FokI, TaqI, BsmI, and ApaI polymorphisms and susceptibility to pulmonary tuberculosis: a meta-analysis. *Genet. Mol. Res.* 14, 9118–9129. <https://doi.org/10.4238/2015.August.7.21>.
- Gao, L., Tao, Y., Zhang, L., Jin, Q., 2010. Vitamin D receptor genetic polymorphisms and tuberculosis: updated systematic review and meta-analysis. *Int. J. Tuberc. Lung Dis.* 14, 15–23. <https://doi.org/10.1002/mas.21461>.
- Chen, C., Liu, Q., Zhu, L., et al., 2013. Vitamin D receptor gene polymorphisms on the risk of tuberculosis: a meta-analysis of 29 case-control studies. *PLoS One* 8, e83843. <https://doi.org/10.1371/journal.pone.0164648>.
- Budhathoki, S., Yamaji, T., Iwasaki, M., et al., 2016. Vitamin D receptor gene polymorphism and the risk of colorectal cancer: a nested case-control study. *PLoS One* 11, 1–13. <https://doi.org/10.1371/journal.pone.0164648>.
- Whitfield, G.K., Remus, L.S., Jurutka, P.W., et al., 2001. Functionally relevant polymorphisms in the human nuclear vitamin D receptor gene. *Mol. Cell. Endocrinol.* 177, 145–159.
- Hussain, T., Naushad, S.M., Ahmed, A., et al., 2019. Association of vitamin D receptor TaqI and ApaI genetic polymorphisms with nephrolithiasis and end stage renal

- disease: a meta-analysis. *BMC Med. Genet.* 20, 1–9. <https://doi.org/10.1186/s12881-019-0932-6>.
- Medeiros, J.F.P., de Oliveira Borges, M.V., Soares, A.A., et al., 2020. The impact of vitamin D supplementation on VDR gene expression and body composition in monozygotic twins: randomized controlled trial. *Sci. Rep.* 10, 1–10. <https://doi.org/10.1038/s41598-020-69128-2>.
- Han, J., Zhang, X., Saraf, S.L., et al., 2018. Risk factors for vitamin D deficiency in sickle cell disease. *Br. J. Haematol.* 181, 828–835. <https://doi.org/10.1111/bjh.15270>.
- Anand, A., Singh, S., Sonkar, A.A., et al., 2017. Expression of vitamin D receptor and vitamin D status in patients with oral neoplasms and effect of vitamin D supplementation on quality of life in advanced cancer treatment. *Wspolczesna Onkol.* 21, 145–151. <https://doi.org/10.5114/wo.2017.68623>.
- Mazdeh, M., Ghafouri-Fard, S., Hatami, M., et al., 2018. Expression analysis of vitamin D signaling pathway genes in epileptic patients. *J. Mol. Neurosci.* 64, 551–558. <https://doi.org/10.1007/s12031-018-1059-5>.
- Kongsbak, M., Levring, T.B., Geisler, C., Von, Essen M.R., 2013. The vitamin D receptor and T cell function. *Front. Immunol.* 4, 1–10. <https://doi.org/10.3389/fimmu.2013.00148>.
- Liu, P.T., Stenger, S., Li, H., et al., 2006. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 311, 1770–1773. <https://doi.org/10.1126/science.1123933>.
- Pike, J.W., Meyer, M.B., 2011. The vitamin D receptor: new paradigms for the regulation of gene expression by 1,25-dihydroxyvitamin D<sub>3</sub>. *Endocrinol. Metab. Clin. N. Am.* 39, 255–269. <https://doi.org/10.1016/j.ecd.2010.02.007>.
- Pike, J.W., Meyer, M.B., 2012. The vitamin D receptor: new paradigms for the regulation of gene expression by 1, 25-dihydroxyvitamin D<sub>3</sub>. *Rheum. Dis. Clin.* 38, 13–27.
- Hausler, M.R., Whitfield, G.K., 2013. Molecular mechanisms of vitamin D action. *Calcif. Tissue Int.* 92, 77–98. <https://doi.org/10.1007/s00223-012-9619-0>.
- Jeffery, L.E., Wood, A.M., Qureshi, O.S., et al., 2012. Availability of 25-hydroxyvitamin D<sub>3</sub> to APCs controls the balance between regulatory and inflammatory T cell responses. *J. Immunol.* 189, 5155–5164.
- Kim, S.J., Ye, S., Ha, E., Chun, E.M., 2018. Association of body mass index with incident tuberculosis in Korea. *PLoS One* 13, e0195104. <https://doi.org/10.1371/journal.pone.0195104>.
- Kumaratne, M., Early, G., Cisneros, J., 2017. Vitamin D deficiency and association with body mass index and lipid levels in Hispanic American adolescents. *Glob. Pediatr. Health* 4, 1–6. <https://doi.org/10.1177/2333794x17744141>.
- Dorhoi, A., Kaufmann, S.H.E., 2014. Tumor necrosis factor alpha in mycobacterial infection. *Semin. Immunol.* 26, 203–209. <https://doi.org/10.1016/j.smim.2014.04.003>.
- Cegielski, J.P., McMurray, D.N., 2004. The relationship between malnutrition and tuberculosis: evidence from studies in humans and experimental animals. *Int. J. Tuberc. Lung Dis.* 8, 286–298.
- Guwatudde, D., Nakakeeto, M., Maganda, A., Chiunda, A., 2003. Tuberculosis in household contacts of infectious cases in Kampala, Uganda. *Am. J. Epidemiol.* 158, 887–898. <https://doi.org/10.1093/aje/kwg227>.
- Sepkowitz, K.A., 1996. How contagious is tuberculosis? *Clin. Infect. Dis.* 23, 954–962. <https://doi.org/10.1093/clinfid/23.5.954>.
- Hong, Y., Kim, Y., Lee, J.J., et al., 2019. Levels of vitamin D-associated cytokines distinguish between active and latent tuberculosis following a tuberculosis outbreak. *BMC Infect. Dis.* 19, 151. <https://doi.org/10.1186/s12879-019-3798-5>.
- Esteve Palau, E., Sánchez Martínez, F., Knobel Freud, H., et al., 2015. Tuberculosis: plasma levels of vitamin D and its relation with infection and disease. *Med. Clin. (Engl. Ed.)* 144, 111–114. <https://doi.org/10.1016/j.medcle.2015.05.003>.

# 19.\_Agussalim\_Bukhari-Najdah2.pdf

## ORIGINALITY REPORT

22%

SIMILARITY INDEX

15%

INTERNET SOURCES

17%

PUBLICATIONS

5%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://m.cafe.daum.net">m.cafe.daum.net</a> Internet Source	1%
2	<a href="http://www.plantpath.cornell.edu">www.plantpath.cornell.edu</a> Internet Source	1%
3	<a href="http://www.nsc.nagoya-cu.ac.jp">www.nsc.nagoya-cu.ac.jp</a> Internet Source	1%
4	Submitted to Universitas Hasanuddin Student Paper	<1%
5	Omer Erdag, Mahfuz Turan, Rifki Ucler, Mehmet Berkoz, Mehmet Fatih Garca, Nazım Bozan, Ahmet Faruk Kiroglu, Hakan Cankaya. "Is Nasal Polyposis Related to Levels of Serum Vitamin D and Vitamin D Receptor Gene Expression?", Medical Science Monitor, 2016 Publication	<1%
6	Vitamin D, 2010. Publication	<1%
7	<a href="http://mdpi-res.com">mdpi-res.com</a> Internet Source	<1%

8

Kathirvel Maruthai, Saranya Sankar, Mahadevan Subramanian. "Methylation Status of VDR Gene and its Association with Vitamin D Status and VDR Gene Expression in Pediatric Tuberculosis Disease", Immunological Investigations, 2020

Publication

<1 %

9

[oro.open.ac.uk](http://oro.open.ac.uk)

Internet Source

<1 %

10

"Handbook of vitamin D in human health", Wageningen Academic Publishers, 2013

Publication

<1 %

11

[assets.researchsquare.com](https://assets.researchsquare.com)

Internet Source

<1 %

12

Minuk, G.Y.. "Rebound hepatitis following withdrawal of immunosuppressive therapy in patients with chronic Hepatitis B viral infections", Hepatology Research, 200409

Publication

<1 %

13

Shabroza Bashir, Ajaz A. Shah, Jahangir I. Dar, Irshad A. Misgar et al. "Association of VDR gene BsmI polymorphism with temporomandibular joint disorders: A case control study in Kashmiri population", Gene Reports, 2022

Publication

<1 %

14	Internet Source	<1 %
15	"Integrated Physiology/Obesity", Diabetes, 2013 Publication	<1 %
16	Submitted to A'Sharqiyah University, Oman Student Paper	<1 %
17	<a href="https://theses.gla.ac.uk">theses.gla.ac.uk</a> Internet Source	<1 %
18	<a href="http://www.koreascience.or.kr">www.koreascience.or.kr</a> Internet Source	<1 %
19	Jun Kyu Song. "Novel Compound Heterozygous Mutations in the Vitamin D Receptor Gene in a Korean Girl with Hereditary Vitamin D Resistant Rickets", Journal of Korean Medical Science, 2011 Publication	<1 %
20	<a href="http://mafiadoc.com">mafiadoc.com</a> Internet Source	<1 %
21	"Understanding the Host Immune Response Against Mycobacterium tuberculosis Infection", Springer Science and Business Media LLC, 2018 Publication	<1 %
22	<a href="http://www.wjgnet.com">www.wjgnet.com</a> Internet Source	<1 %

23	Submitted to Korea University Student Paper	<1 %
24	Tizaoui, Kalthoum, Wajih Kaabachi, Agnès Hamzaoui, and Kamel Hamzaoui. "Association between vitamin D receptor polymorphisms and multiple sclerosis: systematic review and meta-analysis of case-control studies", Cellular and Molecular Immunology, 2014. Publication	<1 %
25	Submitted to Queen Mary and Westfield College Student Paper	<1 %
26	Sudhasini Panda, Ambrish Tiwari, Kalpana Luthra, S. K. Sharma, Archana Singh. "Association of Fok1 VDR polymorphism with Vitamin D and its associated molecules in pulmonary tuberculosis patients and their household contacts", Scientific Reports, 2019 Publication	<1 %
27	napier-repository.worktribe.com Internet Source	<1 %
28	obgyn.pericles-prod.literatumonline.com Internet Source	<1 %
29	bio-protocol.org Internet Source	<1 %
30	citeseerx.ist.psu.edu Internet Source	<1 %

---

31	<a href="http://epdf.tips">epdf.tips</a> Internet Source	<1 %
32	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	<1 %
33	Mochammad Hatta. "Detection and identification of mycobacteria in sputum from suspected tuberculosis patients", BMC Research Notes, 2010 Publication	<1 %
34	Submitted to Universiti Putra Malaysia Student Paper	<1 %
35	<a href="http://d-nb.info">d-nb.info</a> Internet Source	<1 %
36	<a href="http://springerplus.springeropen.com">springerplus.springeropen.com</a> Internet Source	<1 %
37	Submitted to Lebanese American University Student Paper	<1 %
38	A UITTERLINDEN. "Genetic Vitamin D Receptor Polymorphisms and Risk of Disease", Vitamin D, 2005 Publication	<1 %
39	Doaa Gadallah, Ahmed Sedky, Hend Mohamed Esmaeel. "Vitamin D among Patients of Pulmonary and Extrapulmonary Tuberculosis: A Prospective Case-control	<1 %

## Study", Current Respiratory Medicine Reviews, 2020

Publication

- 
- |    |  |      |
|----|--|------|
| 40 | Upendra Yadav, Pradeep Kumar, Vandana Rai. "FokI polymorphism of the vitamin D receptor (VDR) gene and susceptibility to tuberculosis: Evidence through a meta-analysis", Infection, Genetics and Evolution, 2021<br>Publication | <1 % |
| 41 | Submitted to iGroup<br>Student Paper   | <1 % |
| 42 | <a href="http://www.ijsrp.org">www.ijsrp.org</a><br>Internet Source  | <1 % |
| 43 | <a href="http://www.sciencepub.net">www.sciencepub.net</a><br>Internet Source  | <1 % |
| 44 | "Final Technical Progress Report", 'Office of Scientific and Technical Information (OSTI)'<br>Internet Source  | <1 % |
| 45 | <a href="http://docksci.com">docksci.com</a><br>Internet Source  | <1 % |
| 46 | <a href="http://dokumen.pub">dokumen.pub</a><br>Internet Source  | <1 % |
| 47 | <a href="http://gala.gre.ac.uk">gala.gre.ac.uk</a><br>Internet Source  | <1 % |
| 48 | <a href="http://www.scitepress.org">www.scitepress.org</a><br>Internet Source  | <1 % |
-

49 Bao, B.. "Catfish hepcidin gene is expressed in a wide range of tissues and exhibits tissue-specific upregulation after bacterial infection", *Developmental and Comparative Immunology*, 2005  
Publication

<1 %

50 Kali Prasad Pattanaik, Srabasti Sengupta, Bimal Prasad Jit, Riddhi Kotak, Avinash Sonawane. "Host-mycobacteria conflict: Immune responses of the host vs. the mycobacteria TLR2 and TLR4 ligands and concomitant host-directed therapy", *Microbiological Research*, 2022  
Publication

<1 %

51 [backend.orbit.dtu.dk](https://backend.orbit.dtu.dk)  
Internet Source

<1 %

52 [journals.lww.com](https://journals.lww.com)  
Internet Source

<1 %

53 [mail.scialert.net](https://mail.scialert.net)  
Internet Source

<1 %

54 [rupress.org](https://rupress.org)  
Internet Source

<1 %

55 [www.zora.uzh.ch](https://www.zora.uzh.ch)  
Internet Source

<1 %

56 Jeane Franco Pires Medeiros, Michelle Vasconcelos de Oliveira Borges, Aline Alves

<1 %

Soares, Jessica Cavalcante dos Santos et al.  
"The impact of vitamin D supplementation on  
VDR gene expression and body composition  
in monozygotic twins: randomized controlled  
trial", Scientific Reports, 2020

Publication

---

57

Margaret K. Guerriero, Mary W. Redman,  
Kelsey K. Baker, Renato G. Martins et al.  
"Racial disparity in oncologic and quality-of-  
life outcomes in patients with locally  
advanced head and neck squamous cell  
carcinomas enrolled in a randomized phase 2  
trial", Cancer, 2018

Publication

---

<1 %

58

P. Selvaraj. "Plasma 1,25 Dihydroxy Vitamin  
D3 Level and Expression of Vitamin D  
Receptor and Cathelicidin in Pulmonary  
Tuberculosis", Journal of Clinical Immunology,  
02/14/2009

Publication

---

<1 %

59

[academic.oup.com](http://academic.oup.com)

Internet Source

---

<1 %

60

[cyberleninka.org](http://cyberleninka.org)

Internet Source

---

<1 %

61

[ddd.uab.cat](http://ddd.uab.cat)

Internet Source

---

<1 %

62

[etheses.whiterose.ac.uk](http://etheses.whiterose.ac.uk)

Internet Source

<1 %

---

63 [repositorio.ufmg.br](https://repositorio.ufmg.br)  
Internet Source

<1 %

---

64 [tesisenred.net](https://tesisenred.net)  
Internet Source

<1 %

---

65 [www.karger.com](https://www.karger.com)  
Internet Source

<1 %

---

66 [www.oncotarget.com](https://www.oncotarget.com)  
Internet Source

<1 %

---

67 "Extraskkeletal Effects of Vitamin D", Springer  
Science and Business Media LLC, 2018  
Publication

<1 %

---

68 Assel G. Zhumina, Konstantin Li, Anna A.  
Konovalova, Yelena A. Li et al. "Plasma 25-  
Hydroxyvitamin D Levels and VDR Gene  
Expression in Peripheral Blood Mononuclear  
Cells of Leukemia Patients and Healthy  
Subjects in Central Kazakhstan", Nutrients,  
2020  
Publication

<1 %

---

69 Ester Lilian Acen, Irene Andia Biraro,  
Mudarshiru Bbuye, David Patrick Kateete,  
Moses L. Joloba, William Worodria.  
"Hypovitaminosis D among newly diagnosed

<1 %

pulmonary TB patients and their household contacts in Uganda", Scientific Reports, 2022

Publication

---

70

Jian Zhao, Cen Wen, Ming Li. "Association Analysis of Interleukin-17 Gene Polymorphisms with the Risk Susceptibility to Tuberculosis", Lung, 2016

Publication

---

<1 %

71

Ju Yeon Ban. "Promoter Polymorphism (rs3755724, -55C/T) of Tissue Inhibitor of Metalloproteinase 4 (TIMP4) as a Risk Factor for Kawasaki Disease with Coronary Artery Lesions in a Korean Population", Pediatric Cardiology, 04/2009

Publication

---

<1 %

72

Li-Ching Lee, Ying-Chun Cho, Pei-Jung Lin, Ting-Chi Yeh, Chun-Yen Chang, Ting-Kuang Yeh. "Influence of Genetic Variants of the N-Methyl-D-Aspartate Receptor on Emotion and Social Behavior in Adolescents", Neural Plasticity, 2016

Publication

---

<1 %

73

Rasoul Abdollahzadeh, Mohammad Hossein Shushizadeh, Mina Barazandehrokh, Sepideh Choopani et al. "Association of Vitamin D receptor gene polymorphisms and clinical/severe outcomes of COVID-19

<1 %

patients", Infection, Genetics and Evolution,  
2021

Publication

---

74

Victoria Naranjo, Ursula HÄ¶fle, JoaquÄ¶n Vicente, Ma Paz MartÄ¶n et al. " Genes differentially expressed in oropharyngeal tonsils and mandibular lymph nodes of tuberculous and nontuberculous European wild boars naturally exposed to ", FEMS Immunology & Medical Microbiology, 2006

Publication

---

<1 %

75

Wen-Feng Li. "Genetics of osteoporosis: accelerating pace in gene identification and validation", Human Genetics, 12/12/2009

Publication

---

<1 %

76

Zhu, Andy Z.X., Qian Zhou, Lisa Sanderson Cox, Sean P. David, Jasjit S. Ahluwalia, Neal L. Benowitz, and Rachel F. Tyndale. "Association of CHRNA5-A3-B4 SNP rs2036527 with smoking cessation therapy response in African American smokers", Clinical Pharmacology & Therapeutics, 2014.

Publication

---

<1 %

77

[etd.aau.edu.et](http://etd.aau.edu.et)

Internet Source

---

<1 %

78

[fjfsdata01prod.blob.core.windows.net](http://fjfsdata01prod.blob.core.windows.net)

Internet Source

---

<1 %

79	<a href="http://medic.upm.edu.my">medic.upm.edu.my</a> Internet Source	<1 %
80	<a href="http://pr.hec.gov.pk">pr.hec.gov.pk</a> Internet Source	<1 %
81	<a href="http://publichealthinafrica.org">publichealthinafrica.org</a> Internet Source	<1 %
82	<a href="http://soar.suny.edu">soar.suny.edu</a> Internet Source	<1 %
83	<a href="http://unbscholar.lib.unb.ca">unbscholar.lib.unb.ca</a> Internet Source	<1 %
84	<a href="http://westminsterresearch.westminster.ac.uk">westminsterresearch.westminster.ac.uk</a> Internet Source	<1 %
85	<a href="http://www.db-thueringen.de">www.db-thueringen.de</a> Internet Source	<1 %
86	Abeer Ramadan, Sara F. Sallam, Mai S. Elsheikh, Sally R. Ishak et al. "VDR gene expression in asthmatic children patients in relation to vitamin D status and supplementation", Gene Reports, 2019 Publication	<1 %
87	Fernando Lecanda, Pamela M. Warlow, Linda R. Halstead, Thomas H. Steinberg, Roberto Civitelli. "Impaired intramembranous bone formation in connexin43 null mice", Bone, 1998 Publication	<1 %

88

Frances C. Bach, Kirsten Rutten, Kristyanne Hendriks, Frank M. Riemers et al. " The Paracrine Feedback Loop Between Vitamin D (1,25(OH) D ) and PTHrP in Prehypertrophic Chondrocytes ", Journal of Cellular Physiology, 2014

Publication

&lt;1 %

89

Francisca Dayse Martins de Sousa, Iury de Paula Souza, Ednelza da Silva Graça Amoras, Sandra Souza Lima et al. "Low levels of TNFA gene expression seem to favor the development of pulmonary tuberculosis in a population from the Brazilian Amazon", Immunobiology, 2023

Publication

&lt;1 %

90

Kieran G Meade. "Antigen stimulation of peripheral blood mononuclear cells from Mycobacterium bovis infected cattle yields evidence for a novel gene expression program", BMC Genomics, 2008

Publication

&lt;1 %

91

María Jiménez-Sousa, José Jiménez, Amanda Fernández-Rodríguez, Oscar Brochado-Kith et al. "VDR rs2228570 Polymorphism Is Related to Non-Progression to AIDS in Antiretroviral Therapy Naïve HIV-Infected Patients", Journal of Clinical Medicine, 2019

Publication

&lt;1 %

92

Nibiao Zheng, Christopher C. Whalen, Andreas Handel. "Modeling the Potential Impact of Host Population Survival on the Evolution of M. tuberculosis Latency", PLoS ONE, 2014

Publication

<1 %

93

Wafa Nouari, Lamia Ysmail-Dahlouk, Mourad Aribi. "Vitamin D3 enhances bactericidal activity of macrophage against Pseudomonas aeruginosa", International Immunopharmacology, 2016

Publication

<1 %

94

Wei Seong Toh, Zheng Yang, Boon Chin Heng, Tong Cao. "Chapter 23 Differentiation of Human Embryonic Stem Cells Toward the Chondrogenic Lineage", Springer Science and Business Media LLC, 2007

Publication

<1 %

95

[daneshyari.com](http://daneshyari.com)

Internet Source

<1 %

96

[digiresearch.vut.ac.za](http://digiresearch.vut.ac.za)

Internet Source

<1 %

97

[elischolar.library.yale.edu](http://elischolar.library.yale.edu)

Internet Source

<1 %

98

[eprints.nirt.res.in](http://eprints.nirt.res.in)

Internet Source

<1 %

99

[gaffi.org](http://gaffi.org)

Internet Source

<1 %

100 [jcm.asm.org](http://jcm.asm.org)  
Internet Source

<1 %

101 [journal.unhas.ac.id](http://journal.unhas.ac.id)  
Internet Source

<1 %

102 [journal.usm.my](http://journal.usm.my)  
Internet Source

<1 %

103 [liu.diva-portal.org](http://liu.diva-portal.org)  
Internet Source

<1 %

104 [megaaktiva.umkendari.ac.id](http://megaaktiva.umkendari.ac.id)  
Internet Source

<1 %

105 [ojs.ptbioch.edu.pl](http://ojs.ptbioch.edu.pl)  
Internet Source

<1 %

106 [orapp.aut.ac.nz](http://orapp.aut.ac.nz)  
Internet Source

<1 %

107 [peerj.com](http://peerj.com)  
Internet Source

<1 %

108 [publikationen.uni-tuebingen.de](http://publikationen.uni-tuebingen.de)  
Internet Source

<1 %

109 [repository-tnmgrmu.ac.in](http://repository-tnmgrmu.ac.in)  
Internet Source

<1 %

110 [static.elsevier.es](http://static.elsevier.es)  
Internet Source

<1 %

111	<a href="https://translational-medicine.biomedcentral.com">translational-medicine.biomedcentral.com</a> Internet Source	<1 %
112	<a href="https://www.biorxiv.org">www.biorxiv.org</a> Internet Source	<1 %
113	<a href="https://www.futuremedicine.com">www.futuremedicine.com</a> Internet Source	<1 %
114	<a href="https://www.pure.ed.ac.uk">www.pure.ed.ac.uk</a> Internet Source	<1 %
115	<a href="https://www.scirp.org">www.scirp.org</a> Internet Source	<1 %
116	<a href="https://www.tara.tcd.ie">www.tara.tcd.ie</a> Internet Source	<1 %
117	<a href="https://www.thieme-connect.com">www.thieme-connect.com</a> Internet Source	<1 %
118	Antimicrobial Peptides, 2016. Publication	<1 %
119	Barbara J Boucher. "Why do so many trials of vitamin D supplementation fail?", Endocrine Connections, 2020 Publication	<1 %
120	Guangyuan Chen, Cong Hu, Yuxuan Song, Mengxi Xiu, Wanfeng Liang, Ningjing Ou, Xiaoqiang Liu, Peng Huang. "Relationship Between the Apal (rs7975232), Bsml (rs1544410), FokI (rs2228570), and TaqI	<1 %

(rs731236) Variants in the Vitamin D Receptor Gene and Urolithiasis Susceptibility: An Updated Meta-Analysis and Trial Sequential Analysis", *Frontiers in Genetics*, 2020

Publication

---

121

Sergio Díaz-Fernández, Raquel Villar-Hernández, Zoran Stojanovic, Marco Fernández et al. "Study of CD27, CD38, HLA-DR and Ki-67 immune profiles for the characterization of active tuberculosis, latent infection and end of treatment", *Frontiers in Microbiology*, 2022

Publication

---

<1 %

122

"ASBMR 22nd annual meeting", *Journal of Bone and Mineral Research*, 09/2000

Publication

---

<1 %

123

Emad Yuzbashian, Golaleh Asghari, Mehdi Hedayati, Maryam Zarkesh, Parvin Mirmiran, Alireza Khalaj. "Determinants of vitamin D receptor gene expression in visceral and subcutaneous adipose tissue in non-obese, obese, and morbidly obese subjects", *The Journal of Steroid Biochemistry and Molecular Biology*, 2018

Publication

---

<1 %

124

Kazuhiko Yoneda, Hiroki Chikumi, Takeshi Murata, Naomasa Gotoh et al. "Measurement of multidrug efflux pumps by

<1 %

quantitative real-time polymerase chain reaction ", FEMS Microbiology Letters, 2005

Publication

---

125 Keiji Takasu, Jemimah N. Ndabarua, Hieu Thi Pham, Shun-ichiro Takano. "Pupal-adult parasitism of the coconut hispine beetle by the koinobiont pupal parasitoid *Tetrastichus brontispae*", Biological Control, 2018 <1 %

Publication

---

126 Lin He. "Administration of Pioglitazone in Low-Density Lipoprotein Receptor-Deficient Mice Inhibits Lesion Progression and Matrix Metalloproteinase Expression in Advanced Atherosclerotic Plaques", Journal of Cardiovascular Pharmacology, 11/2006 <1 %

Publication

---

127 M. Tiwari, M.K. Verma, P.K. Singh, D. Bharti. "Role of Fok I VDR polymorphism in TB risk assessment; A Study in Central India population", Meta Gene, 2021 <1 %

Publication

---

128 Maulana, Adi, Akira Imai, Theo Van Leeuwen, Koichiro Watanabe, Kotaro Yonezu, Takanori Nakano, Adrian Boyce, Laurence Page, and Anders Schersten. "Origin and geodynamic setting of Late Cenozoic granitoids in Sulawesi, Indonesia", Journal of Asian Earth Sciences, 2016. <1 %

Publication

---

129

Retsepile E. Maphasa, Mervin Meyer, Admire Dube. "The Macrophage Response to Mycobacterium tuberculosis and Opportunities for Autophagy Inducing Nanomedicines for Tuberculosis Therapy", Frontiers in Cellular and Infection Microbiology, 2021

Publication

<1 %

---

130

Submitted to Salisbury University

Student Paper

<1 %

---

131

Teodor T. Postolache, Faisal Akram, Ellen E. Lee, Christopher A. Lowry et al. "Increased brain vitamin D receptor expression and decreased expression of cathelicidin antimicrobial peptide in individuals who died by suicide", Journal of Psychiatric Research, 2020

Publication

<1 %

---

132

W. J. Fassbender, B. Goertz, K. Weismüller, B. Steinhauer, H. Stracke, D. Auch, T. Linn, R. G. Bretzel. "VDR Gene Polymorphisms are Overrepresented in German Patients with Type 1 Diabetes Compared to Healthy Controls without Effect on Biochemical Parameters of Bone Metabolism", Hormone and Metabolic Research, 2002

Publication

<1 %

---

---

Exclude quotes      On

Exclude matches      < 5 words

Exclude bibliography      On