

POLYMORPHISM OF VASCULAR ENDOTHELIAL GROWTH FACTOR REGIO PROMOTER C(-634)G AS A RISK FACTOR OF BALINESE TYPE-2 DIABETIC RETINOPATHY

by Dr Habibah Muhiddin

FILE	EGIO_PROMOTER_C_- 634_G_AS_A_RISK_FACTOR_OF_BALINESE_TYPE-2_D.PDF (322.93K)		
TIME SUBMITTED	13-FEB-2021 08:34AM (UTC+0700)	WORD COUNT	2654
SUBMISSION ID	1508407527	CHARACTER COUNT	12955

6
**POLYMORPHISM OF VASCULAR ENDOTHELIAL
GROWTH FACTOR REGIO PROMOTER C(-634)G AS
A RISK FACTOR OF BALINESE
TYPE-2 DIABETIC RETINOPATHY**

¹Mas-Putrawati, A. A., ²Bakta, M., ²Suastika, K., ³Habiba-Muhiddin, H. S., ¹Niti-Susila, N. K.,

¹Departement of Ophthalmology, Faculty of Medicine, Udayana University/
Sanglah General Hospital, Bali-Indonesia.

²Department of Internal Medicine, Faculty of Medicine, Udayana University, Bali-Indonesia

³Departement of Ophthalmology, Faculty of Medicine, Hasanuddin University,
Makasar-Indonesia

12
Background: Diabetic retinopathy (DR) is one of the complications on diabetic mellitus (DM) patient as a micro vascular retina disorder which caused by a long term of hyperglycemia. This is one of the blindness causes in the world. This research aims to find out Polymorphism of VEGFC(-634)G gene as a risk factor of DR on the Balinese with DM type-2 (DMT2). **Method:** This study is applying two designs, analytical cross sectional and case control. The case is DMT2 patient with DR(+), DMT2 without DR as control. The sequencing technique was performed to evaluate polymorphism and plasma VEGF levels were determined by ELISA. **Results:** Cross sectional study (hospital based) came out with quite high number of DR, 57 people of 97 study samples. This study shows the existence of genetic variation on Gen VEGF C(-634)G, with most often genotype of CG (51.5%). Polymorphism C(-634)G as the risk factor of DR (OR=1.815 CI-95%= 1.077-3.057, $p=0.025$), and high level of VEGF were also significant (QR=3.75, CI-95% 1.34-10.20, $p=0.008$). VEGF level with genotype of CG, CC and GG, not found any difference ($p=0.245$). Logistic regression shows that the most influential variable as the risk factor of DR is VEGF level ($p=0.007$), polymorphism of VEGF C(-634)G ($p=0.022$) and systolic blood pressure ($p=0.023$). **Conclusions:** Polymorphism of VEGF C(-634)G gene and high level of VEGF as the risk factor of DR, and can be used as a reference in handling early stage of DR patient to prevent blindness.

Keywords: diabetic mellitus; retinopathy; polymorphism; VEGF levels.

11

INTRODUCTION

Diabetic retinopathy (DR) is one of complications of diabetes mellitus (DM) patients. This was marked by retina macro vascular disorder caused by long term of hyperglycemic and as a cause of blindness worldwide.

DR risk factors are multi factors one of them is genetic factor. VEGF C(-634)G polymorphism gene was studied due to VEGF as an important factor cause of retina neo-vascularisation. In addition, research regardless of VEGF gene in some countries were varies and inconsistent. Researchs by Awata, *et al* (2002) in Japan, Suganthalakshmi, *et al* (2006) in India, and Yang, *et al* (2010) in China reported that VEGF C-634G gene polymorphism a risk factor of DR.¹⁻³

Address for correspondence:

A. A. Mas Putrawati

Departement of Ophthalmology, Faculty of Medicine,
Udayana University/Sanglah General Hospital,
Bali-Indonesia.

E-Mail: masputra06@gmail.com

Research by Freathy, *et al* (2006) in UK, Nakamura, *et al* (2009) in Japan, and Chun, *et al* (2010) in Korea results in insignificant risk factor of DR.⁴⁻⁶

This study aims to support DR pathogenesis theory through the role of genetic factor of VEGF C(-634)G polymorphism gene on promoter region to induce DR. Specifically this study observed VEGF C(-634)G Balinese variances gene and VEGF levels for evaluating risk factor of DR.

MATERIALS AND METHODS

This study applied two designs, i.e. cross sectional and case control. Case and control were recruited from cross-sectional research. Case was DMT2 with DR complication patient and control was DMT2 without DR complication. This research was carried out at Internal Disease Division of Endocrine and Eyes Clinic Sanglah General Hospital, Bali-Indonesia. Sequencing technique was applied to evaluate polymorphism and ELISA for observing VEGF levels.

1

Open access: www.balimedicaljournal.org and www.ojs.unud.ac.id

DMT2 of Balinese patients age of 40-60 years, 5-15 years suffering from the disease and willing to follow the research by signing informed consent were included in this study. Patient with glaucoma, eye pressure ≥ 21 mmHg, infection, intra ocular trauma, uveitis, vitreous bleeding, intra ocular operation history, intravitreal injection of (anti VEGF, steroid), laser photocoagulation < 6 month, stroke, and myocardial infarction were excluded. DR diagnosis was performed on the basis of slit-lamp bio-microscopic observation applying 78 or 90 dioptri condensing lens and retinal photofundus.

RESULTS

Hospital based cross sectional in this study reveals that 57 (58.7%) of 97 patients were DR. Average age of patients was 52.78 ± 7.59 years, 59.8% of patients were male, DM duration was 6.62 ± 2.33 years, 66% of family history with DM, systolic blood pressure of 131.55 ± 14.09 mmHg, diastolic blood pressure of 89 ± 8.10 mmHg, BMI 24.67 ± 3.7 kg/m², and HbA1C of 8.5 ± 2.55 %. This study obtained the present of genetic variants on VEGF C(-634)G gene with the most frequent genotype was CG (51.5%).

In the case control study, age, sexes, DM duration, BMI and HbA1C parameters were not different between case and control, except for blood pressure is different significantly (Table 1).

Table 1
Subject Characteristic between Case and Control Groups

Characteristic	Case (n=36)	Control (n=36)	p
Age (years)	52.33 \pm 7.94	52.89 \pm 8.78	0.640
Sexes			
Male	22	23	0.500
Female	14	13	
DM (years)	6.32 \pm 3.86	5.71 \pm 5.11	0.580
BMI (kg/m ²)	24.46 \pm 3.54	24.52 \pm 4.03	0.640
Blood pressure (mmHg)			
Systolic	134.17 \pm 12.16	125.42 \pm 11.98	0.004
Diastolic	86.39 \pm 7.13	81.81 \pm 7.09	0.005
HbA1c (%)	8.61 \pm 2.51	8.13 \pm 2.52	0.970

Significant at $p < 0.05$

C(-634)G polymorphism as a risk factor of DR was 1.815 times as can be seen in Table 2 within OR=1.815, CI 95% = 1.077-3.057, $p = 0.025$). In this study, it was obtained that there was no different between distribution of allele frequencies CC, CG, and GG for case and control groups ($p=0.365$). However, allele C was higher in DR(+) compare to DR(-), in contrast allele G was higher in DR(-) compare to DR(+) as can be seen from Table 3. High VEGF levels is 3.75 times as a risk factor of DR (OR=3.75, CI 95% = 1.34-10.2, $p=0.008$) (Table 3). In this study, it was also gained that average VEGF levels with CG, CC and GG

were not different significantly ($p = 0.245$), however, VEGF level of CG genotype was higher compare to other genotypes.

Table 2
VEGF C-634G Polymorphism Gene for Case and Control Groups as Risk Factor of DR

VEGF Polymorphism	Case N=36	Control N=36	p
CG	26	15	0.023
CC	1	5	
GG	9	16	

Significance at $p < 0.05$

Table 3
Allele Distribution of VEGF C(-634)G Polymorphism Gene

Variable	Allele G	Allele C	p
DR(+)	44 (61.10%)	28 (38.90%)	0.365
DR(-)	47 (65.30%)	25 (34.70%)	

Significance at $p < 0.05$

6

Logistic regression analysis indicates that the most influential variables as risk factor of DR were VEGF levels ($p = 0.008$), followed by VEGF C(-634)G polymorphism gene ($p = 0.036$) and systolic blood pressure ($p = 0.038$) as indicates by Table 4 and Figure 1.

Table 4
High VEGF Levels in Case and Control Groups as Risk Factor of DR

VEGF levels	Case	Control	p
High	20	9	0.008
Low	16	27	

Significance at $p < 0.05$




Figure 1
Comparison of VEGF Serum Levels to Polymorphism of VEGF C(-634)G gene, genotype of CC, CG and GG
■ case ■ control

DISCUSSION

Results of VEGF C(-634)G polymorphism gene research worldwide were inconsistent. This is probably due to ethnic and environment difference.^{1,2} DR hospital based research in others countries on the basis of DMT2 resulted in similar results to this research, i.e. > 50%. It has already known that 1/3 of DMT2 population will gradually suffer from DR. Therefore, number of people with DR will be increase in line with increasing of DMT2 patients.^{7,8}

This study found that 66 % of the samples had history of DM. It has already known well that patients with DM family history will have a risk to gain DR. Duration of suffering to DM has also a risk to gain DR. It was reported that 2/3 of DM patients above 15 years will gain DR.^{7,8}

Blood pressure was also affect the present of DR. DM patients frequently have high blood pressure (hypertension). WESDR reported that 17% of DMT2 patients will have hypertension and this will increase to be 25% during 10 years. Undercontrol hypertension decreases retinopathy progresivity risk until 34%.⁶

Research by Chun, *et al* (2010) and Nakamura, *et al* (2008) obtained that insignificant results between the present of DR and VEGF C(-634)G gene.^{5,6} However, research by Awata, *et al* (2002), Suganthalaksmi, *et al* (2006) and Yang, *et al* (2010) gained a significant results between the present of DR and VEGF C(-634)G gene. They reporten that VEGF C(-634) gene is a risk factor of DR.^{1,3}

This study resulted that VEGF C(-634)G gene CG genotype is a risk factor of DR (OR=1.815. CI 95% = 1.077-3.057, $p = 0.023$). Regardless of varies results, it was probably due to ethnic and environment different.

Correlation of allele frequencies as a risk factor of DR was also obtained varies between some researches. Allele frequencies correlated to risk marker for development of DR and increase activity on promoter region. Bleda, *et al* (2011) reported that G allele was significantly higher in DR(+) group ($p=0.004$).⁹ On the other hand, Nakamura, *et al* (2008) found insignificant risk factor in either C and G allele to develop DR ($p=0.244$).⁵ Research by Awata, *et al* (2002) found that C allele is higher in case group (0.48) compare to C allele in control group (0.35).¹

This study obtained that allele frequencies is insignificant different between two group. However, we gained that C allele is higher in DR(+) group compare to DR(-) group. On the other hand, G allele is higher in DR(-) group compare to DR(+) group.

Research by Awata, *et al* (2002) and Yang, *et al* (2010) reported that VEGF level is higher in DR(+) group compare to DR(-) group.^{1,2} This study was in line with study by Awata and Yang *et al*.

This study obtained that high VEGF levels were observed higher in case compare to control group (OR = 3.75, CI 95% = 1.34-10.2, $p=0.008$). DM patients with DR complication have an increased of VEGF levels, therefore, it can be used as a predictive factor and guidance to manage DM patients with DR complication.

VEGF levels, generally can be affected by many factors, such as tissue hypoxia, hyperglycemic, stress oxidative, environment, and other gene polymorphism besides VEGF C(-634)G gene polymorphism. Plasma VEGF levels are not similar to VEGF levels in vitreous and aqueous tissues. VEGF expression in endothelial vascular leads to increase of permeability of blood vein that induce increased of angiogenesis activity. Awata, *et al* (2002) found that VEGF level increase on VEGF C(-634)G gene CC genotype. This finding makes the research consistent with strong genetic correlation on C allele. VEGF C(-634)G gene related to lipo-polysaccharide (LPS) that induce VEGF production from peripheral blood mononuclear cells (PBMCs) in which in gene level that can increase transcription factor.¹

Research by Yang, *et al* (2010) found that VEGF C(-634)G gene CC genotype has VEGF levels significantly higher compare to other genotypes. Therefore, they concluded that for people of Han China, CC genotype as genetic risk factor and main factor to induce high VEGF levels.²

Research by Bleda, *et al* (2011) found that serum VEGF levels with CC, CG and GG genotypes were not insignificantly different as risk factors of DR. Research by Nakamura, *et al* ((2008) also found that there was insignificant different between VEGF with VEGF C(-634)G gene polymorphism in either on vitreous and plasma.⁹

In this study, it was found that serum VEGF levels were insignificant different between CG, CC, and GG genotypes. However, average VEGF levels on DR group with CG genotype has a higher trend compare to CC and GG genotypes.

Polymorphism of C(-634)G gene was a binding site of myeloid zinc finger protein, MZF1 a transcription factor. MZF1 has a function of to bind DNA promoter target and regulate transcription. Their mechanisms have not been understood properly. It could probably due to luciferase reporter system.¹

Further developed research is evaluating mRNA levels on VEGF gene. This was probably strongly correlated to DR development and progresivity. Finding of genetic predisposition on DR can be applied to manage therapy based on individual genetic history and preferably carried out at initial stage DM diagnoses. Therefore, it will be more effective to inhibit DR progresivity.⁵

Logistic regression analysis resulted in that for all predicted independent variables correlated to DR, such as blood pressure (systolic and diastolic),

VEGF levels, and VEGF C(-634)G polymorphism gene, in which the most significant was VEGF levels ($p= 0.007$), followed by VEGF C(-634)G polymorphism gene ($p=0.022$) and finally systolic blood pressure ($p=0.023$).

CONCLUSION

Polymorphism of VEGF C(-634)G gene and high VEGF levels are risk factors of DR on Balinese DMT2 patients.

REFERENCES

1. Awata, T., Inoue, K., Kurihara, S., Ohkubo, T., Watanabe, M., Inukai, K., Inoue, I., Katayama, S. 2002. A common polymorphism in the 5'-untranslated region of the VEGF gene is associated with diabetic retinopathy in type 2 diabetes. *Diabetes*, 51: 1635-1639.
2. Suganthalaksmi, B., Anand, R., Kim, R., Mahalaksmi, R., Kartihikprakash, S., Namperumalsamy, P., *et al.* 2006. Association of VEGF and eNOS gene polymorphisms in type 2 diabetic retinopathy. *Molecular Vision*, 12: 336-341.
3. Yang, Y., Andresen, B.A., Yang, K., Zhang, Y., Li, X. 2010. Association of vascular endothelial growth factor -634C/G polymorphism and diabetic retinopathy in type 2 diabetic Han Chinese. *Experimental Biology and Medicine*, 235:1204-1211.
4. Freathy, R.M., Weedon, M.N., Shields, B., Hitman, G.A., Walker, M., McCarthy, M.I., *et al.* 2006. Functional variation in VEGF is not associated with type 2 diabetes in a united kingdom caucasian population. *Journal of the Pancreas*, 7: 295-302.
5. Nakamura S., Iwasaki N., Funatsu H., Kitano S., Iwamoto. 2008. Impact of variants in the VEGF gene on progression of proliferative diabetic retinopathy. *Graefe's Archive for Clinical and Experimental Ophthalmology*, 247(1):11-15.
6. Chun, M.Y., Hwang, H.S., Cho, H.Y. 2010. Association of vascular endothelial growth factor polymorphism with nonproliferative and proliferative diabetic retinopathy. *The Journal of Clinical Endocrinology and Metabolism*, 95(7): 3547-3551.
7. ADA (American Diabetes Association). 2012. Diagnosis dan Classification of Diabetes Mellitus. *Diabetes care*, 35:564-71.
8. AAO (American Academy of Ophthalmology and Staff). 2012. *Retina and Vitreous Basic and Clinical Science Course*. Section-12. San Francisco: AAO. p. 109-132.
9. Bleda, S., Haro, J.D., Varela, C., Esparza, L., Ferruelo, A., Acin, F. 2011. VEGF are involved in the late vascular complication in type II diabetic patients. *Diabetes & Vascular disease research* 9(1): 68-74



This work is licensed under
a Creative Commons Attribution

POLYMORPHISM OF VASCULAR ENDOTHELIAL GROWTH FACTOR REGIO PROMOTER C(-634)G AS A RISK FACTOR OF BALINESE TYPE-2 DIABETIC RETINOPATHY

ORIGINALITY REPORT

% **8**

SIMILARITY INDEX

% **4**

INTERNET SOURCES

% **7**

PUBLICATIONS

% **2**

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to King's College

Student Paper

% **1**

2

Satagopan Uthra, Rajiv Raman, Bickol N. Mukesh, Padmaja Kumari R et al. " Intron 4 VNTR of Endothelial Nitric Oxide Synthase (eNOS) Gene and Diabetic Retinopathy in Patients in Southern India ", Ophthalmic Genetics, 2009

Publication

% **1**

3

T. Awata, K. Inoue, S. Kurihara, T. Ohkubo, M. Watanabe, K. Inukai, I. Inoue, S. Katayama. "A Common Polymorphism in the 5'-Untranslated Region of the VEGF Gene Is Associated With Diabetic Retinopathy in Type 2 Diabetes", Diabetes, 2002

Publication

% **1**

4

Priyanka Biswas, Amit Kr. Chakraborty, Anumoy Mukherjee, Baidyanath Pal, Debasish Maji, Madhusnata De. "Elevated Vascular Endothelial

% **1**

Growth Factor Level in Association with Mean Platelet Volume Are Emerging Risk Factors for Vascular Complications in T2DM Patients", Health, 2019

Publication

-
- 5 "Abstracts of the 2012 Meeting of the International Neuro-Ophthalmology Society, Singapore", Neuro-Ophthalmology, 2012
Publication % 1
-
- 6 www.genes2cognition.org
Internet Source % 1
-
- 7 ascopubs.org
Internet Source <% 1
-
- 8 onlinelibrary.wiley.com
Internet Source <% 1
-
- 9 Yuan, Yazhen, Zhuan Wen, Yongqing Guan, Yanhua Sun, Jie Yang, Xiaohui Fan, Xinzhi Yang, and Rui Liu. "The relationships between type 2 diabetic retinopathy and VEGF-634G/C and VEGF-460C/T polymorphisms in Han Chinese subjects", Journal of Diabetes and its Complications, 2014.
Publication <% 1
-
- 10 www.ias.ac.in
Internet Source <% 1
-

Thapa, R, G Paudyal, N Maharjan, and PS

11

Bernstein. "Demographics and awareness of diabetic retinopathy among diabetic patients attending the vitreo-retinal service at a tertiary eye care center in Nepal", Nepalese Journal of Ophthalmology, 2012.

Publication

<% 1

12

Netasha Khan, Andrew D. Paterson, Delnaz Roshandel, Ali Raza, Muhammad Ajmal, Nadia K. Waheed, Maleeha Azam, Raheel Qamar. "Association of IGF1 and VEGFA polymorphisms with diabetic retinopathy in Pakistani population", Acta Diabetologica, 2019

Publication

<% 1

13

Mostafa Fegghi, Abdolrahim Nikzamir, Alireza Esteghamati, Touraj Mahmoudi, Mir Saeed Yekaninejad. "Relationship of vascular endothelial growth factor (VEGF) +405 G/C polymorphism and proliferative retinopathy in patients with type 2 diabetes", Translational Research, 2011

Publication

<% 1

14

K.-C. Hsueh, Y.-J. Lin, J.-S. Chang, L. Wan, Y.-H. Tsai, C.-H. Tsai, C.-P. Chen, F.-J. Tsai. "Association of Vascular Endothelial Growth Factor C-634 G Polymorphism in Taiwanese Children With Kawasaki Disease", Pediatric Cardiology, 2007

Publication

<% 1

15

El-Shazly, Sherien F, Mohamed H El-Bradey, and Mohamed K Tameesh. "Vascular endothelial growth factor gene polymorphism prevalence in patients with diabetic macular oedema and its correlation with anti-vascular endothelial growth factor treatment outcomes : Vascular endothelial growth factor gene", *Clinical and Experimental Ophthalmology*, 2013.

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5 WORDS