

Vitreous and Serum Concentrations of Vascular Endothelial Growth Factor and Platelet-Derived Growth Factor in Proliferative Diabetic Retinopathy

by Habibah Muhiddin

Submission date: 22-Mar-2021 10:46PM (UTC-0700)

Submission ID: 1540082557

File name: ndothelial_Growth_Factor_and_Platelet-Derived_Growth_Factor2.pdf (566.93K)

Word count: 4461

Character count: 23804

Vitreous and Serum Concentrations of Vascular Endothelial Growth Factor and Platelet-Derived Growth Factor in Proliferative Diabetic Retinopathy

3
This article was published in the following Dove Press journal:
Clinical Ophthalmology

Habibah Setyawati Muhiddin
Muhammad Irfan Kamaruddin 
Andi Muhammad Ichsan
Budu 
39
Department of Ophthalmology, Faculty of
Medicine, Hasanuddin University,
Makassar, South Sulawesi, Indonesia

57
Purpose: This study aimed to investigate the concentrations of vascular endothelial growth factor (VEGF) and platelet-derived growth factor (PDGF) in vitreous and serum samples, analyze the ratio, and compare among proliferative diabetic retinopathy (PDR) subgroups.

Patients and Methods: This study included 17 eyes of patients with PDR, identified as the PDR group which was divided into three subgroups (vitreous hemorrhage [VH], VH with fibrotic tissues, and tractional retinal detachment), and five control eyes (nucleus and intraocular lens drop). Vitreous and serum samples were obtained on the same day. The VEGF-A and PDGF-AB concentrations were calculated by enzyme-linked immunosorbent assay.

Results: The VEGF-A and PDGF-AB concentrations in vitreous samples were significantly higher in the PDR group (630.72 ± 342.81 pg/mL) compared with those in the control group (153.58 ± 145.85 pg/mL); however, they were not detected in serum samples. The vitreous/serum ratio of the VEGF-A concentration in the PDR group (2.1 ± 1.8) was significantly higher compared with that in the control group (0.31 ± 0.33). The VEGF-A concentrations in vitreous samples were highest in the VH group and lowest in the VH with fibrotic tissue subgroup (mean difference 536.16 pg/mL). The vitreous VEGF-A/PDGF-AB concentration ratios were also significantly different among the PDR subgroups.

Conclusion: High concentrations of VEGF and PDGF in vitreous samples of PDR eyes indicate its local related activity in PDR pathology. There is a possibility of PDGF involvement in the pathogenesis of PDR. The VEGF/PDGF concentration ratios possibly play a significant role in the formation of fibrotic tissue in PDR.

Keywords: vascular endothelial growth factor, platelet-derived growth factor, proliferative diabetic retinopathy, fibrotic tissue

Introduction


The neovascularization process in the retina depends on the growth factor production induced by hypoxic condition.¹ Vascular endothelial growth factor (VEGF) is a dominant proangiogenic factor in proliferative diabetic retinopathy (PDR), which triggers migration and proliferation of endothelial cells and blood flow increase, together with microvascular permeability.²⁻⁴ It was proven to have significant role on the retina of patients with PDR, especially on retinal pigment epithelial (RPE) cells, glial cells, vitreous fibroblast, ganglion cells, and Muller cells.^{1,5,6} The amount and duration of VEGF needed to pass through the blood-brain barrier are possibly less than the concentrations needed to induce neovascularization. Hence,

Correspondence: Habibah Setyawati
Muhiddin
Email muhiddinhabibah@yahoo.co.id

26
submit your manuscript | www.dovepress.com
Dovepress    
<http://dx.doi.org/10.2147/OPTH.S240812>

Clinical Ophthalmology 2020:14 1547-1552

1547

 © 2020 Muhiddin et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at <https://www.dovepress.com/terms.php> and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (<http://creativecommons.org/licenses/by-nc/3.0/>). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 42 and 5 of our Terms (<https://www.dovepress.com/terms.php>).

monoclonal antibodies against VEGF were introduced and used widely in the form of anti-VEGF agents, such as pegaptanib, ranibizumab, and bevacizumab.^{5,7,8}

Another growth factor found to contribute to the pathology of PDR is platelet-derived growth factor (PDGF).^{9–11} Both VEGF and PDGF have very similar characteristics.^{2,3} PDGF could directly trigger cellular proliferation and movement. It is an autocrine growth stimulator in the RPE cells, which participates in the retinal wound repair and epiretinal membrane formation.¹² There are four different PDGF chains: PDGF-A, PDGF-B, PDGF-C, and PDGF-D. They later form five isoforms that bind to two different PDGF receptors (PDGFR α , PDGFR β), with different affinities.¹³ The three isoforms of PDGF (AA, AB, and BB) were proven not only to be related to the neovascularization processes in PDR but also to play important role in the formation of fibrotic tissues in the retina of patients with PDR.¹⁴ This study aimed to investigate the concentrations of VEGF and PDGF in vitreous and serum samples, analyze the ratio, and compare it among clinical PDR stage.

Patients and Methods

Study Design and Subjects

This is an analytical observational study with a cross-sectional design. We enrolled consecutively attending adults (>18 years old) who had a clinically confirmed diagnosis of PDR, nucleus drop, and intraocular lens (IOL) drop. Inclusion criterion for PDR group was patients with PDR who were candidates for vitreoretinal surgery for long-standing (>3 months) vitreous or preretinal hemorrhage or tractional retinal detachment (TRD). The PDR group was divided into three subgroups: subgroup I, vitreous hemorrhage (VH); subgroup II, VH with fibrotic tissues; subgroup III, TRD. The control group consisted of patients without diabetes who underwent pars plana vitrectomy due to a whole nucleus drop or IOL drop. Exclusion criteria for the PDR group were previous ocular surgery within the last 2 years, existing or a history of ocular inflammation, rubeosis iridis or neovascular glaucoma, and rhegmatogenous retinal detachment.

The study was conducted in August 2018–March 2019 and was approved by the Institutional Review Board of the Medical Faculty of Hasanuddin University of Makassar (ethics committee reference number 1106/H4.8.4.5.31/PP36-KOMETIK/2018). Written informed consent for surgery, blood sampling, and vitreous sampling was obtained from

all participants in accordance with the Declaration of Helsinki.

Sample Collection and Measurement of VEGF-A and PDGF-AB Concentrations

Blood samples were collected from all participants on the day of surgery and were left for 2 h at room temperature until clotted. Serum was separated by centrifugation at 4°C for 15 min at 1000 × g, divided into aliquots, and then stored at –80°C until assayed.

At the onset of vitrectomy, 0.5–0.8 mL of undiluted vitreous samples was obtained by aspiration into a 1 mL sterile syringe attached to the vitreous cutter (Alcon Constellation) with the stopcock of the infusion closed. The samples were transferred into sterilized Corning microcentrifuge tubes (1.5 mL), placed immediately on ice, and centrifuged for 20 min at 4°C at 1000 × g. Supernatants without sediment were divided into aliquots and immediately frozen at –80°C until assayed.

The VEGF-A and PDGF-AB concentrations were measured in the vitreous and serum samples by enzyme-linked immunosorbent assay (ELISA) using the *Elabscience Human VEGF-A and PDGF-AB ELISA Kit*. Each assay was performed in duplicate according to the manufacturer's instructions, using 100 μ L aliquots of vitreous or serum samples, then diluted accordingly to comply with the detection range of the relevant assay, and eventually to contain the same amount of protein. The data presented are per milligram of protein. The optical density was determined at 450 and 570 nm using an absorption spectrophotometer.

Statistical Analyses

All values are expressed as mean \pm standard deviation, and all data are presented in the tables and figures. Statistical analyses were conducted using Statistical Package for Social Science version 21.0. The differences between groups were tested using Mann–Whitney *U*-test and Kruskal–Wallis test. A value of $p < 0.05$ was considered statistically significant.

Results

Patient Characteristics

A total of 22 eyes were enrolled, 17 eyes of 17 patients with PDR and 5 eyes of 5 patients with nucleus and IOL drop as controls. Based on the PDR subgroups, subgroup I (VH) was diagnosed in six patients; whereas, subgroup II (VH with fibrotic tissues) was diagnosed in five patients and subgroup III (TRD) in six patients. Nineteen eyes had

46 best-corrected visual acuity (BCVA) between 0–3/60 and three eyes had BCVA better than 3/60 (Table 1).

VEGF-A and PDGF-AB Concentrations in Vitreous and Serum Samples

The mean vitreous VEGF-A concentrations in the PDR group (659.01 ± 332.92 pg/mL) were significantly higher compared with that in the control group (190.94 ± 138.05 pg/mL and 57.01 ± 20.83 pg/mL) ($p < 0.05$) (Table 2). The vitreous PDGF-AB concentrations in the PDR group (124.52 ± 53.87 pg/mL) were also significantly higher compared with that in the control group (57.01 ± 20.83 pg/mL) ($p < 0.05$) (Table 2).

The serum concentrations of both VEGF-A and PDGF-AB in the PDR group (551.38 ± 461.84 pg/mL and 98.69 ± 38.56 pg/mL) were not significantly different compared with those in the control group (577.85 ± 336.29 pg/mL and 140.13 ± 80.93 pg/mL) ($p = 0.493$ and $p = 0.319$) (Table 2). Interestingly, the concentrations of PDGF-AB in the control group were higher compared with that in the PDR group.

The VEGF-A/PDGF-AB concentration ratios in both vitreous and serum samples were not significantly higher in the diabetic patients compared with those in the control group (Table 3). The vitreous/serum ratio of the VEGF-A

Table 1 Patients' Characteristics

Variable	Value	Groups	
		PDR	Control
Gender			
Male	12	8	4
Female	10	9	1
Eyes			
Right	8	6	2
Left	14	11	3
BCVA			
0–3/60	19	15	4
>3/60	3	2	1
Hypertension			
Yes	9	9	0
No	13	8	5
Age (years)	52.64 ± 10.1	52.29 ± 10.74	53.80 ± 8.47
IOP (mmHg)	13.86 ± 2.1	13.82 ± 2.30	14.00 ± 1.60
Blood Glucose (mg/dl)	161 ± 74.34	179.00 ± 75.50	99.80 ± 12.00

Notes: The data are expressed as number and mean \pm SD (according to data variable).

Abbreviations: PDR, proliferative diabetic retinopathy; BCVA, best-corrected visual acuity; IOP, intraocular pressures.

Table 2 The VEGF-A and PDGF-AB Concentrations

Variable	Group	Group		P
		PDR (pg/mL)	Control (pg/mL)	
VEGF-A	Vitreous	630.72 ± 342.81	153.58 ± 145.85	0.001*
	Serum	551.38 ± 461.84	577.85 ± 336.29	0.493
PDGF-AB	Vitreous	119.43 ± 56.21	63.67 ± 23.40	0.031*
	Serum	98.69 ± 38.56	140.13 ± 80.93	0.319

Notes: The data are expressed as mean \pm SD. *Significant difference between PDR and control group using Mann–Whitney test ($P < 0.05$).

Abbreviations: PDR, proliferative diabetic retinopathy; VEGF-A, vascular endothelial growth factor-A; PDGF-AB, platelet-derived growth factor-AB.

concentrations was significantly higher in the PDR group compared with that in the control group ($p < 0.05$), but not significant in the PDGF-AB concentrations ($p = 0.058$) (Table 3).

VEGF-A and PDGF-AB Concentrations in Vitreous and Serum Samples of Each PDR Group

There was no significant difference in the serum concentrations of both VEGF-A ($p = 0.586$) and PDGF-AB ($p = 0.265$) among all PDR subgroups in our study. The vitreous VEGF-A concentrations were significantly different among all PDR subgroups ($p < 0.05$). *Post hoc* Bonferroni shows that there were significantly lower concentrations of vitreous VEGF-A in subgroup II compared with subgroup I (mean difference 536.16 pg/mL; $*p < 0.05$) (Figure 1).

There are interesting patterns of the VEGF-A and PDGF-AB concentrations in vitreous samples: in subgroup II, the vitreous VEGF-A concentrations were lowest, whereas the vitreous PDGF-AB concentrations were highest. There was no significant difference in the vitreous/serum ratio of both the VEGF-A and PDGF-AB concentrations among the PDR subgroups ($p = 0.754$ (VEGF = A); $p = 0.482$ (PDGF = AB)) (Table 4). The VEGF-A/PDGF-AB concentration ratios in

Table 3 The VEGF-A and PDGF-AB Concentration Ratios on Vitreous and Serum

Ratio Variable	Variable	PDR	Control	P
Vitreous/Serum	VEGF-A	2.1 ± 1.8	0.31 ± 0.33	0.005*
	PDGF-AB	1.32 ± 0.68	0.66 ± 0.52	0.058
VEGF-A/PDGF-AB	Vitreous	6.21 ± 4.03	2.66 ± 2.15	0.071
	Serum	5.82 ± 5.3	6.39 ± 5.96	0.940

Notes: The data are expressed as mean \pm SD. *Significant difference between PDR and control group using Mann–Whitney test ($P < 0.05$).

Abbreviations: PDR, proliferative diabetic retinopathy; VEGF-A, vascular endothelial growth factor-A; PDGF-AB, platelet-derived growth factor-AB.




Figure 1 The Vitreous VEGF-A and PDGF-AB concentrations in each PDR group. **Notes:** Group classifications, I: PDR with VH; II: PDR with VH and fibrotic tissues; III: PDR with TR. *Significant difference using Post hoc Bonferroni test ($P < 0.05$). **Abbreviations:** PDR, proliferative diabetic retinopathy; VH, vitreous hemorrhage; TRD, tractional retinal detachment; VEGF-A, vascular endothelial growth factor-A; PDGF-AB, platelet-derived growth factor-AB.

vitreous samples were significantly different among each PDR subgroup ($p < 0.01$), but not in serum samples ($p = 0.591$) (Table 4). The vitreous VEGF-A/PDGF-AB concentration ratios were lowest in subgroup II (2.54) (Table 4).

Discussion

In this study, we found that the vitreous VEGF-A concentrations in patients with PDR were significantly increased compared with those in controls, which is supported by the

Table 4 The VEGF-A and PDGF-AB Concentration Ratios in Vitreous and Serum in Each PDR Group

Ratio Variable	PDR Groups	Ratio	P
VEGF-A/PDGF-AB Vitreous	I	8.04±2.62	*0.008
	II	2.54±1.26	
	III	7.45±4.94	
VEGF-A/PDGF-AB Serum	I	4.85±2.54	0.591
	II	3.83±2.91	
	III	8.43±7.95	
VEGF-A Vitreous/Serum	I	2.43±1.97	0.754
	II	1.53±1.07	
	III	2.25±2.25	
PDGF-AB Vitreous/Serum	I	1.06±0.50	0.482
	II	1.72±1.02	
	III	1.25±0.39	

Notes: The data are expressed as mean±SD. Group classifications, I: PDR with VH; II: PDR with VH and fibrotic tissues; III: PDR with TRD. *Significant difference using Kruskal-Wallis test among all PDR groups ($P < 0.05$)

Abbreviations: PDR, proliferative diabetic retinopathy; VH, vitreous hemorrhage; TRD, tractional retinal detachment; VEGF-A, vascular endothelial growth factor-A; PDGF-AB, platelet-derived growth factor-AB.

evidence that the vitreous/serum ratio of the VEGF-A concentrations was also significantly higher in the PDR group compared with that in the control group. Similar results have been reported by Praidou et al.¹⁰ Baharivand et al.¹⁵ and Wang et al.¹⁶ Chernykh et al.¹⁷ even reported that vitreous VEGF concentrations of patients with PDR were 17 times higher compared with those of controls. This result is supported by well-known theories about VEGF's role in PDR pathology.

The concentrations of vitreous PDGF-AB in our study were also increased in PDR eyes compared with controls and even though its vitreous/serum concentration ratios were not significantly high, our findings with respect to ratios were complex. Freyberger et al.⁹ reported that the PDGF-AB concentrations were significantly increased in patients with PDR, with a higher level in rubeosis iridis eyes. Praidou et al.¹⁰ also reported that PDGF concentrations of all isoforms were increased significantly in patients with PDR. Our result not only suggests that PDGF has a role in the pathology of PDR; it also provides evidence supporting how anti-PDGF could possibly work on the neovascularization processes in PDR. Previously, Phase IIb study by Jaffe et al.¹⁸ already reported how the combination of anti-VEGF and anti-PDGF was superior to anti-VEGF alone in the treatment of neovascular AMD.

In this study, we found that there was no significant difference in the serum VEGF-A concentrations between PDR and controls. Praidou et al.¹⁰ also confirmed such findings in their study. This finding was contrary to the theory that diabetic patients have high serum VEGF levels which will lead to diabetes complications, not only retinopathy but also nephropathy, neuropathy, atherosclerosis and other vascular disturbances.¹⁹ Baharivand et al.¹⁵ found in their study that the serum VEGF levels in patients with PDR were increased compared with those in controls. The difference in our result may be due to the fact that our control group was not free from other systemic conditions, which could cause increase in VEGF concentration. The reason was supported by the fact that in the PDR group, the vitreous VEGF-A concentrations were also not significantly different from the serum VEGF-A concentrations, meaning that our serum VEGF-A concentrations were possibly high. In this study, we also found that the vitreous VEGF-A concentrations of the VH subgroup were significantly high compared with the VH with fibrotic tissue subgroup. The vitreous VEGF-A concentrations were highest in the VH subgroup and lowest in the VH with fibrotic tissue subgroup. Contrarily, the vitreous PDGF-AB concentrations were

highest in the VH with fibrotic tissue subgroup and lowest in the VH subgroup. To the best of our knowledge, there are no studies yet that have similar results to ours. A study by Pennock et al²⁰ found that VEGF-A was a competitor of PDGF toward PDGFRs. In 2014, Pennock et al²¹ even reported that VEGF-A was worked via PDGFR α to maintain the sustainability of cells that express both VEGF and PDGF receptors. A recent study in 2016 by Pennock et al²² reported that VEGF-A not only worked on endothelial cells to maintain and stimulate angiogenesis but also worked on PDGFR α to maintain the sustainability of cells enduring hypoxia. These studies' results could explain our results, why when hypoxia happen (active PDR, VH group), the PDGF-AB concentrations decrease, possibly because high concentrations of VEGF-A at that stage act as a competitor of PDGF toward PDGFR α , indirectly decreasing the concentrations of PDGF in vitreous samples.

Another question is why the VEGF-A concentrations are low in the VH with fibrotic tissue subgroup. A study by Kuiper et al²³ could explain the situation; they found that after the increase in the VEGF-A concentration in the VH stage, the level of connective tissue growth factor (CTGF) also increases and inactivates VEGF-A, so its concentrations decrease gradually. Van Geest RF et al²⁴ even reported that the CTGF/VEGF ratio was a strong predictive factor in the formation of fibrotic tissue in PDR. These results further support that anti-VEGF therapy should not be given when there are already fibrovascular membranes present because it could possibly aggravate angiofibrotic switch that could accelerate TRD.

In this study, we also reported interesting results regarding the vitreous VEGF-A/PDGF-AB ratio in each PDR subgroup. To the best of our knowledge, this is the first study to report the vitreous VEGF-A/PDGF-AB concentration ratios in PDR group. The ratio of the VH with fibrotic tissue subgroup was lower than that of the VH subgroup, and surprisingly, the ratio was not only influenced by lower VEGF-A concentration but also by higher PDGF-AB concentration. Based on these findings, we assume that there is an indication that the vitreous VEGF-A/PDGF-AB ratio attempted to reach an equilibrium state in the VH with fibrotic tissue subgroup and that the ratio was possibly related to the process of fibrotic tissue formation.

The empirical results reported herein should be considered in light of some limitations. It is quite difficult to find truly healthy subjects to be included in the control group in our study; thus, they are subject to biases and confounding factors that may influence our results. However, we attempted

to exclude any control eyes with conditions that could influence growth factor concentrations, including fragmented nucleus drop. Besides, for the patients with PDR in our study, advanced examinations, such as fundus fluorescein angiography and optical coherence tomography, were not performed to clearly state the ischemic condition of the retina. We also have a relatively small sample size compared with the other previous related studies, but we used proper statistical analyses to maintain good results.

Conclusion

High concentrations of VEGF and PDGF in vitreous samples of PDR eyes, but not in serum samples, indicate its local related activity in PDR pathology. The vitreous VEGF-A concentrations in the VH with fibrotic tissue subgroup was low in contrast to the PDGF-AB concentrations. The vitreous VEGF-A/PDGF-AB ratio is possibly associated with the formation of fibrotic tissue in PDR. Further investigations and experimental studies using animal models may be required in the future.

Acknowledgments

We would like to express our great appreciation to Dr Burhanuddin Bahar, MS for assistance with statistical analyses, and Dr Rizalinda Sjahril, MSc, PhD for her valuable and constructive suggestions during the planning and development of this research work. The authors also would like to thank Enago for the English language review.

Disclosure

The authors report no conflicts of interest in this work.

References

- Perrin R, Konopatskaya O, Qiu Y, et al. Diabetic retinopathy is associated with a switch in splicing from anti- to pro-angiogenic isoforms of vascular endothelial growth factor. *Diabetologia*. 2005;48:2422–2427. doi:10.1007/s00125-005-1951-8
- Ferrara N. Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev*. 2004;25:581–611.
- Young T, Miller J. Intraocular neovascularization. In: Ryan SJ, Schachat AP, editors. *Retina*. 4th ed. Mosby: St. Louis; 2006:2403–2411.
- Frank R. Etiologic mechanisms in diabetic retinopathy. In: Ryan SJ, Schachat AP, editors. *Retina*. 4th ed. Mosby: St. Louis; 2006:1241–1270.
- Shams N, Ianchuleaev T. Role of vascular endothelial growth factor in ocular angiogenesis. *Ophthalmol Clin N Am*. 2006;9:335–344.
- Boulton M, Foreman D, Williams G. VEGF localization in diabetic retinopathy. *Br J Ophthalmol*. 1998;82:561–568. doi:10.1136/bjo.82.5.561

7. Jorge R, Costa RA, Calucci D, et al. Intravitreal bevacizumab for persistent new vessels in diabetic retinopathy (IBEPE Study). *Retina*. 2006;26:1006–1013. doi:10.1097/01.iae.0000246884.76018.63
8. Andreoli CM, Miller JW. Anti-vascular endothelial growth factor therapy for ocular neovascular disease. *Curr Opin Ophthalmol*. 2007;18(6):502–508. doi:10.1097/ICU.0b013e3282f0ca54
9. Freyberger H, Brocker M, Yakut H, et al. Increased levels of platelet-derived growth factor in the vitreous fluid of patients with proliferative diabetic retinopathy. *Exp Clin Endocrinol Diabetes*. 2000;108(02):106–109. doi:10.1055/s-2000-5803
10. Praidou A, Klangas I, Papakonstantinou E, et al. Vitreous and serum levels of platelet-derived growth factor and their correlation in patients with proliferative diabetic retinopathy. *Curr Eye Res*. 2009;34(2):152–161. doi:10.1080/02713680802585920
11. Praidou A, Papakonstantinou E, Andraudi S, et al. Vitreous and serum levels of vascular endothelial growth factor and platelet-derived growth factor and their correlation in patients with non-proliferative diabetic retinopathy and clinically significant macula oedema. *Acta Ophthalmol*. 2011;89(3):248–254. doi:10.1111/j.1755-3768.2009.01661.x
12. Cassidy L, Barry P, Shaw C, et al. Platelet derived growth factor and fibroblast growth factor basic levels in the vitreous of patients with vitreoretinal disorders. *Br J Ophthalmol*. 1998;82(2):181–185. doi:10.1136/bjo.82.2.181
13. Li X, Eriksson U. Novel PDGF family members: PDGF-C and PDGF-D. *Cytokine Growth Factor Rev*. 2003;14(2):91–98. doi:10.1016/S1359-6101(02)00090-4
14. Klaassen I, de Vries EW, Vogels Ilse MC, et al. Identification of proteins associated with clinical and pathological features of proliferative diabetic retinopathy in vitreous and fibrovascular membranes. *PLoS One*. 2017;12(11):e0187304. doi:10.1371/journal.pone.0187304
15. Baharivand N, Zarghami N, Panahi F, et al. Relationship between vitreous and serum vascular endothelial growth factor levels, control of diabetes and microalbuminuria in proliferative diabetic retinopathy. *Clin Ophthalmol*. 2012;6:185–191. doi:10.2147/OPHTH.S27423
16. Wang J, Chen S, Jiang F, et al. Vitreous and plasma VEGF levels as predictive factors in the progression of proliferative diabetic retinopathy after vitrectomy. *PLoS One*. 2014;9(10):e110531. doi:10.1371/journal.pone.0110531
17. Chernykh VV, Varvarinsky EV, Smimov EV, Chernykh DV, Trunov AN. Proliferative and inflammatory factors in the vitreous of patients with proliferative diabetic retinopathy. *Indian J Ophthalmol*. 2015;63(1):33–36. doi:10.4103/0301-4738.151464
18. Jaffe GJ, Ciulla TA, Ciardella AP, et al. Dual antagonism of PDGF and VEGF in neovascular age-related macular degeneration a phase IIb, multicenter, randomized controlled trial. *Ophthalmology*. 2017;124:224–234. doi:10.1016/j.ophtha.2016.10.010
19. Guo L, Jiang F, Tang Y-T, et al. The association of serum vascular endothelial growth factor and ferritin in diabetic microvascular disease. *Diabetes Technol Ther*. 2014;16(4):224–234. doi:10.1089/dia.2013.0181
20. Pennock S, Kazlauskas A. Vascular endothelial growth factor A competitively inhibits platelet-derived growth factor (PDGF)-dependent activation of PDGF receptor and subsequent signaling events and cellular responses. *Mol Cell Bio*. 2012;32:1955–1966.
21. Pennock S, Haddock LJ, Mukai S, Kazlauskas A. Vascular endothelial growth factor acts primarily via platelet-derived growth factor receptor α to promote proliferative vitreoretinopathy. *Am J Pathol*. 2014;184(11):3052–3068. doi:10.1016/j.ajpath.2014.07.026
22. Pennock S, Kim LA, Kazlauskas A. Vascular endothelial cell growth factor A acts via platelet-derived growth factor receptor α to promote viability of cells enduring hypoxia. *Mol Cell Biol*. 2016;36(18):2314–2327. doi:10.1128/MCB.01019-15
23. Kuiper EJ, Van Nieuwenhoven FA, de Smet MD, et al. The angio-fibrotic switch of VEGF and ctgf in proliferative diabetic retinopathy. *PLoS One*. 2008;3(7):e2675. doi:10.1371/journal.pone.0002675
24. Van Geest RJ, Lesnik-Oberstein SY, Tan HS, et al. A shift in the balance of vascular endothelial growth factor and connective tissue growth factor by bevacizumab causes the angiofibrotic switch in proliferative diabetic retinopathy. *Br J Ophthalmol*. 2012;96(4):587–590. doi:10.1136/bjophthalmol-2011-301005

Clinical Ophthalmology

Dovepress

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed

Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>

Vitreous and Serum Concentrations of Vascular Endothelial Growth Factor and Platelet-Derived Growth Factor in Proliferative Diabetic Retinopathy

ORIGINALITY REPORT

30%

SIMILARITY INDEX

21%

INTERNET SOURCES

29%

PUBLICATIONS

11%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Austin Peay State University 3%
Student Paper
- 2** research.birmingham.ac.uk 3%
Internet Source
- 3** www.dovepress.com 2%
Internet Source
- 4** Ali Keles, Kenan Sonmez, Yasemin Ozdamar Erol, Sema Nur Ayyıldız, Elmas Ogus. "Vitreous levels of vascular endothelial growth factor, stromal cell–derived factor-1 α , and angiopoietin-like protein 2 in patients with active proliferative diabetic retinopathy", Graefe's Archive for Clinical and Experimental Ophthalmology, 2020 1%
Publication
- 5** www.researchsquare.com 1%
Internet Source
- 6** www.pubmedcentral.nih.gov 1%
Internet Source

7 Submitted to Padjadjaran University 1%

Student Paper

8 Wei Chen, Qianyi Lu, Li Lu, Huaijin Guan. 1%

"Increased levels of alphaB-crystallin in vitreous fluid of patients with proliferative diabetic retinopathy and correlation with vascular endothelial growth factor", Clinical & Experimental Ophthalmology, 2017

Publication

9 Xiangke Yin, Xianchai Lin, Xiangrong Ren, Bo 1%

Yu, Lixian Liu, Zhimin Ye, Qishan Chen, Chunsik Lee, Weisi Lu, Dechao Yu, Xuri Li. "Novel multi-targeted inhibitors suppress ocular neovascularization by regulating unique gene sets", Pharmacological Research, 2019

Publication

10 www.oncotarget.com 1%

Internet Source

11 Flyn Mckinnirey, Benjamin Herbert, Graham 1%

Vesey, Sharon McCracken. "Mesenchymal Stem Cell Mediated Immune Modulation Is Donor Sex Specific", Research Square, 2021

Publication

12 Submitted to Universitas Hasanuddin 1%

Student Paper

13

Internet Source

1%

14

Jian Jang You. "ELEVATION OF ANGIOGENIC FACTOR CYSTEINE-RICH 61 LEVELS IN VITREOUS OF PATIENTS WITH PROLIFERATIVE DIABETIC RETINOPATHY :", *Retina*, 01/2012

Publication

1%

15

journals.plos.org

Internet Source

1%

16

www.lib.kobe-u.ac.jp

Internet Source

1%

17

s3-eu-west-1.amazonaws.com

Internet Source

<1%

18

Duh, E.J.. "Vitreous levels of pigment epithelium-derived factor and vascular endothelial growth factor: implications for ocular angiogenesis", *American Journal of Ophthalmology*, 200404

Publication

<1%

19

L Cassidy, P Barry, C Shaw, M J Duffy, S Kennedy. "Platelet derived growth factor and fibroblast growth factor basic levels in the vitreous of patients with vitreoretinal disorders", *British Journal of Ophthalmology*, 1998

Publication

<1%

20

Yu, Jing. "Expression of IGFBP-6 in proliferative vitreoretinopathy rat models and its effects on retinal pigment epithelial-J cells", *Molecular Medicine Reports*, 2013.

Publication

<1%

21

mafiadoc.com

Internet Source

<1%

22

pure.uva.nl

Internet Source

<1%

23

www.spandidos-publications.com

Internet Source

<1%

24

Submitted to Free University of Bolzano

Student Paper

<1%

25

Mojca Urbančič, Veronika Kloboves Prevodnik, Daniel Petrovič, Mojca Globočnik Petrovič. "A Flow Cytometric Analysis of Vitreous Inflammatory Cells in Patients with Proliferative Diabetic Retinopathy", *BioMed Research International*, 2013

Publication

<1%

26

epub.ub.uni-muenchen.de

Internet Source

<1%

27

Amir Ramadan Gomaa, Eman Tayae Elsayed, Reham Fadi Moftah. "MicroRNA-200b Expression in the Vitreous Humor of Patients with Proliferative Diabetic Retinopathy",

<1%

28

DEMET NAK, Nuriye Ozlem Kucuk, Emre Can Celebioglu, Mehmet Sadik BILGIC, Serhat HAYME, Metin Kemal KIR. "The Role of F18-FLT PET / CT in Assessing Early Response to TARE and TACE in Patients with Primary and Metastatic Liver Tumors", Research Square, 2021

Publication

<1%

29

Submitted to Mansoura University

Student Paper

<1%

30

Pennock, Steven, Leo A. Kim, and Andrius Kazlauskas. "VEGF-A acts via PDGFR α to promote viability of cells enduring hypoxia", Molecular and Cellular Biology, 2016.

Publication

<1%

31

"Research progress on the role of connective tissue growth factor in fibrosis of diabetic retinopathy", International Journal of Ophthalmology, 2018

Publication

<1%

32

Pedro Brito, Jorge Costa, Nuno Gomes, Sandra Costa, Jorge Correia-Pinto, Rufino Silva. "Serum pro-inflammatory factors as predictors of persistent diabetic macular oedema with limited anatomic response to anti- : association with

<1%

intravitreal injection treatment profiles in real-world setting ", Acta Ophthalmologica, 2019

Publication

33

Qader Motarjemizadeh, Naser Samadi Aidenloo, Siamak Sepehri. "A comparative study of different concentrations of topical bevacizumab on the recurrence rate of excised primary pterygium: a short-term follow-up study", International Ophthalmology, 2015

Publication

<1%

34

www.hindawi.com

Internet Source

<1%

35

www.karger.com

Internet Source

<1%

36

D.Y. Chen, G.F. Su. "Tumor necrosis factor-like weak inducer of apoptosis association with proliferative diabetic retinopathy and promotes proliferation and collagen synthesis in retinal ARPE-19 cells", Genetics and Molecular Research, 2016

Publication

<1%

37

Qian Zhang, Wanting Xing, Jie Zhang, Junwen Hu, Lunan Qi, Bangde Xiang. "Circulating Tumor Cells Undergoing the Epithelial–Mesenchymal Transition: Influence on Prognosis in Cytokeratin 19-Positive Hepatocellular Carcinoma", OncoTargets and

<1%

38

Submitted to University of Wales Institute,
Cardiff

Student Paper

<1%

39

Zulvikar Syambani Ulhaq, Gita Vita Soraya,
Budu, Lely Retno Wulandari. "The role of IL-6-
174 G/C polymorphism and intraocular IL-6
levels in the pathogenesis of ocular diseases: a
systematic review and meta-analysis", Scientific
Reports, 2020

Publication

<1%

40

Farid Panahi, Nader Baharivand, Zarghami,
Dokht Ghafari M. Yazdan, Ali Mahdavi Fard,
Mohajeri Abas. "Relationship between vitreous
and serum vascular endothelial growth factor
levels, control of diabetes and microalbuminuria
in proliferative diabetic retinopathy", Clinical
Ophthalmology, 2012

Publication

<1%

41

www.onlinejacc.org

Internet Source

<1%

42

Dawit Kumilachew Yimenu, Eyerusalem Shello
Adelo, Ebrahim Abdela Siraj, Tadael Assaye
Kassie et al. "

<1%

Health Professionals Oral Health Knowledge

and Practice: Unleashing the Hidden Challenges

", Journal of Multidisciplinary Healthcare, 2020

Publication

43

Grisanti, Swaantje, Qi Zhu, Olcay Tatar, Julia Lueke, Matthias Lueke, Aysegul Tura, and Salvatore Grisanti. "DIFFERENTIAL EXPRESSION OF VASCULAR ENDOTHELIAL GROWTH FACTOR-A ISOFORMS IN NEOVASCULAR AGE-RELATED MACULAR DEGENERATION :", Retina, 2014.

Publication

<1%

44

Rafael Simo, Esther Carrasco, Marta Garcia-Ramirez, Cristina Hernandez. "Angiogenic and Antiangiogenic Factors in Proliferative Diabetic Retinopathy", Current Diabetes Reviews, 2006

Publication

<1%

45

Rui Cheng, Jian-xing Ma. "Angiogenesis in diabetes and obesity", Reviews in Endocrine and Metabolic Disorders, 2015

Publication

<1%

46

"Retinal Detachment Surgery and Proliferative Vitreoretinopathy", Springer Nature America, Inc, 2018

Publication

<1%

47

Bing Li, Meng-Da Li, Jun-Jie Ye, Zhe Chen, Zi-

<1%

Jian Guo, Yu Di. "Vascular endothelial growth factor concentration in vitreous humor of patients with severe proliferative diabetic retinopathy after intravitreal injection of conbercept as an adjunctive therapy for vitrectomy", Chinese Medical Journal, 2020

Publication

48

M Yokoi. "Elevations of AGE and vascular endothelial growth factor with decreased total antioxidant status in the vitreous fluid of diabetic patients with retinopathy", British Journal of Ophthalmology, 2005

Publication

49

Yong Jie Qin, Sun On Chan, Hong Liang Lin, Yu Qiao Zhang et al. "Increased Expression of Growth Hormone–Releasing Hormone in Fibrinous Inflammation of Proliferative Diabetic Retinopathy", American Journal of Ophthalmology, 2020

Publication

50

link.springer.com

Internet Source

51

repub.eur.nl

Internet Source

52

Evangelos S. Gragoudas. "Pegaptanib for Neovascular Age-Related Macular Degeneration", New England Journal of

<1%

<1%

<1%

<1%

<1%

53

Rogério A. Costa, Rodrigo Jorge, Daniela Calucci, José A. Cardillo, Luiz A. S. Melo, Ingrid U. Scott. "Intravitreal Bevacizumab for Choroidal Neovascularization Caused by AMD (IBeNA Study): Results of a Phase 1 Dose-Escalation Study", *Investigative Ophthalmology & Visual Science*, 2006

Publication

<1%

54

Shalin S Shah, Lori Vidal Denham, Jasmine R Elison, Partha S Bhattacharjee, Christian Clement, Tashfin Huq, James M Hill. "Drug delivery to the posterior segment of the eye for pharmacologic therapy", *Expert Review of Ophthalmology*, 2014

Publication

<1%

55

Shinoda, Kei, Susumu Ishida, Shinichi Kawashima, Toshiko Wakabayashi, Motoko Uchita, Tadayuki Matsuzaki, Michiyo Takayama, Ken Shinmura, and Masakazu Yamada. "Clinical factors related to the aqueous levels of vascular endothelial growth factor and hepatocyte growth factor in proliferative diabetic retinopathy", *Current Eye Research*, 2000.

Publication

<1%

56

ikee.lib.auth.gr

Internet Source

<1%

57

Kinya Tsubota, Yoshihiko Usui, Yoshihiro Wakabayashi, Jun Suzuki, Shunichiro Ueda, Hiroshi Goto. "

<1%

Effectiveness of prophylactic intravitreal bevacizumab injection to proliferative diabetic retinopathy patients with elevated preoperative intraocular VEGF in preventing complications after vitrectomy

", Clinical Ophthalmology, 2019

Publication

58

Laura Liu, Wei-Chi Wu, Ling Yeung, Nan-Kai Wang et al. "Ghost Cell Glaucoma After Intravitreal Bevacizumab for Postoperative Vitreous Hemorrhage Following Vitrectomy for Proliferative Diabetic Retinopathy", Ophthalmic Surgery, Lasers, and Imaging, 2010

<1%

Publication

59

Rong Li, Arvydas Maminishkis, Fei E. Wang, Sheldon S. Miller. "PDGF-C and -D Induced Proliferation/Migration of Human RPE Is Abolished by Inflammatory Cytokines", Investigative Ophthalmology & Visual Science, 2007

<1%

Publication

60

Jiaxing Wang, Song Chen, Feng Jiang, Caiyun You, Chunjie Mao, Jinguo Yu, Jindong Han,

<1%

Zhuhong Zhang, Hua Yan. "Vitreous and Plasma VEGF Levels as Predictive Factors in the Progression of Proliferative Diabetic Retinopathy after Vitrectomy", PLoS ONE, 2014

Publication

61

Joana Mesquita, João Paulo Castro-de-Sousa, Sara Vaz-Pereira, Arminda Neves, Luís A. Passarinha, Cândida T. Tomaz. "Evaluation of the growth factors VEGF-a and VEGF-B in the vitreous and serum of patients with macular and retinal vascular diseases", Growth Factors, 2018

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off