

# On Weak Solution of Cauchy Problem for Nonlinear Parabolic Equations

*by*

---

FILE	ARIS_2019_J._PHYS.__CONF._SER._1341_062038.PDF (597.72K)		
TIME SUBMITTED	18-NOV-2019 12:41PM (UTC+0700)	WORD COUNT	2998
SUBMISSION ID	1216037425	CHARACTER COUNT	12040

PAPER • OPEN ACCESS

## On Weak Solution of <sup>2</sup>Cauchy Problem for Nonlinear Parabolic Equations

To cite this article: Naimah Aris *et al* 2019 *J. Phys.: Conf. Ser.* **1341** 062038

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices  
to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of  
every title for free.

## On Weak Solution of Cauchy Problem for Nonlinear Parabolic Equations

Naimah Aris<sup>1</sup>, Muh. Rifki Nisardi<sup>1</sup>, Kasbawati<sup>1</sup>, Jeffry Kusuma<sup>1</sup>, Budi Nurwahyu<sup>1</sup>, Mawardi<sup>1</sup>

<sup>1</sup>Department of Mathematics, Hasanuddin University, Makassar, Indonesia

E-mail: newima@gmail.com

**Abstract.** In this paper, we investigate weak solutions of Cauchy problem for nonlinear parabolic equations. By using Galerkin approximation, the existence and uniqueness of weak solutions are proved.

### 1. Introduction

There are several problems in partial differential equations, of which classical solution is difficult to solve. Different approaches are needed to ensure the existence of a solution of the equations. One way to determine the solution of a differential equation is to expand the space of solution which contains the smooth solution and the non-smooth solution. The solution contained in this space is called a weak solution [17].

The existence of the weak solution of nonlinear parabolic equations has been investigated in many papers. Qiang *et al.*, [2] have shown the existence and some properties of the weak solution of nonlinear parabolic equations without formulating it into abstract Cauchy problem.

In partial differential equations theory, some mathematical problems endowed by appropriate conditions and spaces can be transformed into an abstract Cauchy problem. Many papers deal with the Cauchy problem for nonlinear parabolic equations, [2-8]. Choe and Lee [4] explain the existence and uniqueness of Cauchy problem for degenerate parabolic equations using Harnack estimate. Shang [6] study the existence and non-existence of local and global solutions of the Cauchy problem by a priori estimate and compactness methods. The properties of the solutions of the Cauchy problem for non-divergent form with density, appear in Rainbek's paper in 2015.

In this paper, we discuss the weak solution of the abstract Cauchy problem of nonlinear parabolic equations

$$\begin{aligned} \frac{\partial u}{\partial t} + Au &= f(u) + \delta \\ u(0) &= u_0 \end{aligned} \quad (1)$$

where  $A$  is a continuous linear operator that does not depend on time,  $A = -\Delta$  and  $f$  is a continuous function. We use Galerkin approximation [1] to show the existence of the weak solution. For the proof, we use the following notation

$$\Psi = \{u \in \mathcal{D}(\Omega), \operatorname{div} u = 0\}$$

$$V = \text{Closure from } \Psi \text{ in } H_0^1(\Omega)$$

$$H = \text{Closure from } \Psi \text{ in } L^2(\Omega)$$

Let  $V$  and  $H$  be Hilbert spaces.  $V \subset H$ ,  $V$  dense in  $H$  with injective dan continuous function. Scalar product and norm of  $V$  and  $H$  are denoted by  $((\cdot, \cdot))$ ,  $\|\cdot\|$ ,  $(\cdot, \cdot)$ ,  $|\cdot|$ .  $H$  is recognizable using its dual,  $H'$  which can be identified using by a dense subspace  $V$ , hence

$$V \subset H \subset V'$$

Scalar product in of  $f \in H$  and  $u \in V$  given by

$$\langle f, u \rangle = (f, u), \forall f \in H, \forall u \in V. \quad (2)$$

For every  $u$  in  $V$

$$v \in V \rightarrow ((u, v)) \in \mathbb{R}$$

be a linear and continuous map in  $V$ . There exist an element in  $V'$  denote by  $Au$  such that

$$\langle Au, v \rangle = ((u, v)), \forall v \in V. \quad (3)$$

where  $A: V \rightarrow V'$  be a linear and continuous operator.  $A \in \mathcal{L}(V, V')$  define on  $V$ , so that  $A$  is a canonic isomorfism of  $V$  on  $V'$ .

Assuming that  $u$  is the solution to equation (1), with  $u \in C^2(\Omega)$ . If  $v$  is element of  $\Psi$ , we obtain

$$\left(\frac{du}{dt}, v\right) + ((u, v)) = (f, v) + (\delta, v).$$

From the continuity of equation (3), for each  $v \in V$  apply

$$\left(\frac{du}{dt}, v\right) = \frac{d}{dt} (u, v).$$

It is the basis to form a weak solution to a problem in the equation (1). For  $f \in L^2(0, T; V')$  and the given  $u_0 \in H$ , so will be determined  $u$  that fulfil

$$u \in L^2(0, T; V)$$

and

$$\frac{d}{dt} (u, v) + ((u, v)) = \langle f, v \rangle + \langle \delta, v \rangle, \forall v \in V.$$

Then the problem (1) can be formulated as follows

$$\begin{aligned} u' + Au &= f(u) + \delta \text{ untuk } t \in (0, T), \\ u(0) &= u_0. \end{aligned}$$

Equation (2) is an equation distributed on  $V'$ . Based on (2), Equation (3) can be formulated as

$$\begin{aligned} u' &= (f(u) + \delta) - Au \\ \frac{d}{dt} \langle u, v \rangle &= \langle (f(u) + \delta) - Au, v \rangle, \forall v \in V. \end{aligned} \quad (4)$$

Since  $A$  linear operator and continuous from  $V$  to  $V'$  and  $u \in L^2(0, T; V)$ , so that function  $Au$  is found on  $L^2(0, T; V')$ . Hence,  $(f(u) + \delta) - Au \in L^2(0, T; V')$  and from Equation (4) as well as Lemma 1 indicate that

$$u' \in L^2(0, T; V').$$

In this case, a function should determine  $u \in L^2(0, T; V) \cap C^0(0, T; H)$  in such a way that

$$u_t = \frac{du}{dt} \in L^2(0, T; V').$$

#### Lemma 1

Let  $\Omega$  be an open bounded set in  $\mathbb{R}^m$ , and  $g_j$  is a sequence function on  $L^p(\Omega)$  with

$$\|g_j\|_{L^p(\Omega)} \leq C \quad \text{for each } j \in \mathbb{Z}^+$$

If  $g \in L^p(\Omega)$  and  $g_j \rightarrow g$  almost in every point on  $g_j \rightarrow g$  di  $L^p(\Omega)$ .

**Theorem 1**

Let  $X \subset H \subset Y$  is Banach space, with  $X$  is reflexive space. Given that  $u_n$  is a limited uniform term on  $L^2(0, T; X)$  and  $\frac{du_n}{dt}$  limited uniform on  $L^p(0, T; Y)$ , for  $p > 1$ . Then, a strong convergent sub term exists on  $L^2(0, T; H)$ .

**Lemma 2**

Let  $X$  is a reflexive Banach space and  $x_n$  is a limited-term  $X$ . Then,  $x_n$  has a weak convergent sub term on  $X$ .

The following conditions will use throughout the paper

**Assumption 1**

$f$  is a function of  $L^2(0, T; V')$  fulfilling

$$-k - \alpha_1 |s|^p \leq f(s)s \leq k - \alpha_2 |s|^p, \quad p > 2 \tag{5}$$

$$f'(s) \leq l. \tag{6}$$

**2. Result**

Equation (1) fulfil an equation at  $L^q(0, T; H^{-s}(\Omega))$ , where  $q$  is conjugate pair at equation (5). Also (1) fulfil similarity at  $L^q(0, T; H^{-s}(\Omega))$ , hence for any  $v \in L^p(0, T; H^s(\Omega))$  we have

$$\begin{aligned} \left\langle \frac{du}{dt}, v \right\rangle + \langle Au, v \rangle &= \langle f(u), v \rangle + \langle \delta, v \rangle \\ \left\langle \frac{du}{dt}, v \right\rangle + ((u, v)) &= \langle f(u), v \rangle + \langle \delta, v \rangle \end{aligned}$$

for almost each  $t \in [0, T]$ .

Using Sobolev embedding theorem, obtained rules

$$s \geq m(p - 2)/2p$$

given  $p$  dependent on function given on eq (14) on (5). If chosen  $m = 1$  or  $2$ , then obtained  $s = 1$ , so that the equation obtained in familiar space  $L^q(0, T; H^{-1}(\Omega))$  or  $L^q(0, T; V^*(\Omega))$ .

**Theorem 2 (Weak Solution)**

Equation (1) and  $f$  is a function of  $C^1$  fulfilling by assumption A, have a unique weak solution  $u$  for any  $T > 0$  and given  $u_0 \in H$  with

$$u \in L^2(0, T; H_0^1(\Omega) \cap L^p(\Omega_T)) \text{ And } u \in C^0([0, T]; L^2(\Omega))$$

in addition,  $u_0 \mapsto u(t)$  continuous on  $L^2(\Omega)$ .

**Proof:**

First, assume that  $V$  separated. Form independent linear sequence of complete elements  $V; \omega_1, \dots, \omega_m, \dots$ . Observe for any value of  $m$  can be defined as a form of approach solution  $u_m$  for Equation (1) defined as follows

$$u_m(t) = \sum_{i=1}^m g_{im} \omega_i.$$

Function  $g_{im}, 1 \leq i \leq m$  is a function of scalar defined in  $[0, T]$ , and multiplication result of  $\omega_j$ , Equation (1) and  $u_m$  obtained

$$\begin{aligned} \left\langle \frac{du_m}{dt} + Au_m, \omega_j \right\rangle &= \langle f(u) + \delta, \omega_j \rangle \\ \left\langle \frac{du_m}{dt}, \omega_j \right\rangle + \langle Au_m, \omega_j \rangle &= \langle f(u), \omega_j \rangle + \langle \delta, \omega_j \rangle \\ \langle u_m', \omega_j \rangle + \langle (u_m, \omega_j) \rangle &= \langle f(u), \omega_j \rangle + \langle \delta, \omega_j \rangle \\ u_m(0) &= u_{0m} \quad \omega_j \in V. \end{aligned}$$

$u_{0m}$  is orthogonal projection on  $H$  from  $u_0$  in space spanned by  $\omega_1, \dots, \omega_n$ .

Multiply Equation (7) with  $g_{jm}(t)$  so that obtained

$$\langle u_m', \omega_j \rangle g_{jm}(t) + \langle (u_m, \omega_j) \rangle g_{jm}(t) = \langle f(u_m), \omega_j \rangle g_{jm}(t) + \langle \delta, \omega_j \rangle g_{jm}(t).$$

Then sum both term obtained

$$\sum \langle u_m', \omega_j \rangle g_{jm}(t) + \sum \langle (u_m, \omega_j) \rangle g_{jm}(t) = \sum \langle f(u_m), \omega_j \rangle g_{jm}(t) + \sum \langle \delta, \omega_j \rangle g_{jm}(t).$$

Since  $\sum \omega_j g_{jm}(t) = \sum g_{jm}(t) \omega_j = u_m(t)$ , we have

$$\langle u_m'(t), u_m \rangle + \langle (u_m, u_m) \rangle = \langle f(u_m), u_m \rangle + \langle \delta, u_m \rangle.$$

Since

$$2\langle u_m'(t), u_m(t) \rangle = \frac{d}{dt} |u_m(t)|^2$$

we obtain

$$\frac{1}{2} \frac{d}{dt} |u_m|^2 + \|u_m\|^2 = \int_{\Omega} f(u_m) u_m \, dx + \int_{\Omega} \delta u_m \, dx.$$

Using the Equation (5), we obtain

$$\frac{1}{2} \frac{d}{dt} |u_m|^2 + \|u_m\|^2 \leq \int_{\Omega} k - \alpha_2 |u_m|^p \, dx + \delta \int_{\Omega} u_m \, dx$$

$$\frac{1}{2} \frac{d}{dt} |u_m|^2 + \|u_m\|^2 \leq \int_{\Omega} k - \alpha_2 |u_m|^p \, dx + |\delta| \|u_m\|.$$

Using Young inequality  $ab \leq \frac{1}{2}(a^2 + b^2)$

$$\frac{1}{2} \frac{d}{dt} |u_m|^2 + \|u_m\|^2 \leq \int_{\Omega} k - \alpha_2 |u_m|^p \, dx + \frac{1}{2} (|\delta|^2 + \|u_m\|^2)$$

$$\frac{1}{2} \frac{d}{dt} |u_m|^2 + \|u_m\|^2 \leq \int_{\Omega} k - \alpha_2 |u_m|^p \, dx + \frac{1}{2} |\delta|^2 + \frac{1}{2} \|u_m\|^2$$

$$\frac{1}{2} \frac{d}{dt} |u_m|^2 + \frac{1}{2} \|u_m\|^2 \leq \int_{\Omega} k - \alpha_2 |u_m|^p \, dx + \frac{1}{2} |\delta|^2.$$

Integrate both term from 0 to  $T$

$$\begin{aligned} & \frac{1}{2} \int_0^T \frac{d}{dt} |u_m(t)|^2 dt + \frac{1}{2} \int_0^T \|u_m(t)\|^2 dt \\ & \leq \int_0^T \int_{\Omega} k dx dt - \int_0^T \int_{\Omega} \alpha_2 |u_m(t)|^p dx dt + \int_0^T \frac{1}{2} |\delta|^2 dt \\ \frac{1}{2} |u_m(T)|^2 - \frac{1}{2} |u_m(0)|^2 + \frac{1}{2} \int_0^T \|u_m(t)\|^2 dt + \int_0^T \int_{\Omega} \alpha_2 |u_m(x,t)|^p dx dt & \leq kT|\Omega| + \frac{1}{2} |\delta|^2 T \\ \frac{1}{2} |u_m(T)|^2 + \frac{1}{2} \int_0^T \|u_m(t)\|^2 dt + \int_0^T \int_{\Omega} \alpha_2 |u_m(x,t)|^p dx dt & \leq \frac{1}{2} |u_0|^2 + kT|\Omega| + \frac{1}{2} |\delta|^2 T. \end{aligned}$$

Given  $|\Omega|$  is the measurement of  $\Omega$ ,  $|\Omega| = \int_{\Omega} dx$ . Based on the above inequality, we obtain

$$\begin{aligned} \sup |u_m(T)|^2 & \leq 2\gamma \\ \int_0^T \|u_m(t)\|^2 & \leq 2\gamma \\ \int_0^T \int_{\Omega} |u_m(x,t)|^p dx dt & \leq \gamma/\alpha_2 \end{aligned}$$

where  $\gamma = \frac{1}{2} |u_0|^2 + kT|\Omega| + \frac{1}{2} |\delta|^2 T$ , hence, based on the boundary, we can write

$$\begin{aligned} u_m & \text{ bounded uniform on } L^{\infty}(0, T; H). \\ u_m & \text{ bounded uniform on } L^2(0, T; V). \\ u_m & \text{ bounded uniform on } L^p(\Omega_T). \end{aligned}$$

Sequence  $u_m$  that is bounded uniform on  $L^p(\Omega_T)$  can be used to obtain the limit on a nonlinear function of  $f(u_m)$  since

$$|f(s)| \leq \beta(|s|^{p-1} + 1).$$

From Equation (5), boundary on  $u_m$  di  $L^p(\Omega_T)$  can be used to obtain the boundary of  $f(u_m)$  on  $L^q(\Omega_T)$  with  $(p, q)$  is a conjugate pair, hence

$$\begin{aligned} \|f(u_m)\|_{L^q(\Omega_T)}^q & = \int_0^T \left( \int_{\Omega} |f(u_m)|^q dx \right) dt \\ \|f(u_m)\|_{L^q(\Omega_T)}^q & \leq \beta \int_0^T \left( \int_{\Omega} |u_m|^{p-1} + 1 \right)^q dx dt \\ \|f(u_m)\|_{L^q(\Omega_T)}^q & \leq C \int_0^T \left( \int_{\Omega} |u_m|^{q(p-1)} + 1 dx \right) dt. \end{aligned}$$

Since  $\frac{1}{p} + \frac{1}{q} = 1$ , such that  $q(p-1) = p$ , then

$f(u_m)$  bounded uniform on  $L^q(\Omega_T)$ .

Furthermore, bounded uniform is required to derive equation (1). It is worth noted that both spaces  $L^2(0, T; V^*)$  and  $L^q(0, T; L^q(\Omega))$  contain continue on  $L^q(0, T; H^{-s}(\Omega))$ , with  $q < 2$  karena  $p > 2$ . Then use (4)

$$\frac{du_m}{dt} = -Au_m + P_m f(u_m) + \delta$$

So that  $\frac{du_m}{dt}$  bounded uniform on  $L^q(0, T; H^{-s}(\Omega))$ .

Using Lemma 2 to derive a sub term with weak convergence, namely  $u_m$

$$\begin{aligned} u_m &\rightharpoonup u \text{ on } L^2(0, T; V) \\ u_m &\rightharpoonup u \text{ on } L^p(\Omega_T) \\ f(u_m) &\rightharpoonup \chi \text{ on } L^q(\Omega_T). \end{aligned}$$

Applying compactness property on Theorem 1, an additional sub term can be defined as

$$u_m \rightharpoonup u \text{ di } L^2(0, T; H). \quad (8)$$

It has been revealed previously that  $u_m$  bounded uniform on  $L^2(0, T; H_0^1(\Omega))$  and also  $\frac{du_m}{dt}$  bounded uniform on  $L^q(0, T; H^{-s}(\Omega))$ . Since  $H_0^1(\Omega) \subset\subset L^2(\Omega) \subset H^{-s}(\Omega)$  and  $H_0^1(\Omega)$  is Hilbert space, so  $H_0^1(\Omega)$  is reflexive. Therefore, we use compactness theorem to acquire (8)

In the next step, we prove the convergence of  $P_m f(u_m) \rightharpoonup \chi$  di  $L^q(\Omega_T)$ .

$$\int_{\Omega_T} (P_m f(u_m) - \chi) \phi dxdt = \int_{\Omega_T} (f(u_m) - \chi) \phi dxdt - \int_{\Omega_T} Q_m f(u_m) \phi dxdt$$

for any given  $\phi \in L^p(\Omega_T)$ . It has been revealed that the first part of the right term close to 0. For the second part of the right term, we define function  $\phi$  as

$$\phi = \sum_{j=1}^m \alpha_j(t) \phi_j$$

Let  $\alpha_j \in L^p(0, T)$ ,  $\phi_j \in C_c^\infty(\Omega)$  dense on  $L^p(\Omega_T)$ . So for function

$$\int_{\Omega_T} Q_m f(u_m) \sum_{j=1}^m \alpha_j(t) \phi_j dxdt = \int_{\Omega_T} f(u_m) \sum_{j=1}^m \alpha_j(t) Q_m \phi_j dxdt.$$

Because  $Q_m \phi_j \rightarrow 0$  di  $L^p(\Omega)$  for each  $j$ . Hence, we can prove that  $P_m f(u_m)$  is convergent.

The facts imply each of the equation found in space  $L^2(0, T; V) \cap L^p(\Omega_T)$  is convergent on those dual spaces. The dual space is  $L^2(0, T; V^*) \cap L^q(\Omega_T)$ . Equation

$$\frac{du}{dt} + Au = \chi. \quad (9)$$

Fulfil similarity in this room. To make more clear, providing the fact that

$$L^2(0, T; V^*) \cap L^q(\Omega_T) \subset L^q(0, T; H^{-s}).$$

Hence, it is absolute that Equation (9) fulfil similarity in that space.

Furthermore, we go on proving that  $\chi = f(u)$ . To do this, we use the weak convergence of  $u_m$  to  $u$  di  $L^2(\Omega_T)$  and also Lemma 2. Since  $u_m \rightharpoonup u$  di  $L^2(\Omega_T)$ , using the Lemma 1 to guarantee the existence of sub term  $u_{n_j}$  in such a way that  $u_{n_j}(x, t) \rightarrow u(x, t)$  for each  $(x, t) \in \Omega_T$ .

So that by using the continuity of  $f$ , that  $f(u_{n_j}(x, t)) \rightarrow f(u(x, t))$  for almost every  $(x, t) \in \Omega_T$  along the boundary on  $f(u_m)$  di  $L^q(\Omega_T)$  provided on Equation (7), we apply Lemma 1 to conclude that  $f(u_{n_j}(x, t)) \rightarrow f(u(x, t))$  di  $L^q(\Omega_T)$ . From singularity of *weak limits*, cause  $\chi = f(u)$ .

To prove that solution  $u(t)$  continue from  $[0, T]$  on  $L^2(\Omega)$ , we investigate  $u(0, T; V) \cap L^p(\Omega_T)$ , using Theorem 1 to conclude  $u \in C^0([0, T]; L^2(\Omega))$ .

To prove  $u(0) = u_0$ , take  $\phi \in C^1([0, T]; V) \cap L^p(\Omega)$ , with  $\phi(T) = 0$ . Notice that,  $\phi \in L^2(0, T; V) \cap L^p(\Omega_T)$ . So that, by *limiting equation*

$$\left\langle \frac{du}{dt}, v \right\rangle + a(u, v) = \langle f(u), v \rangle + \langle \delta, v \rangle.$$

Implementing partial integral to variable  $t$ ,

$$\int_0^T -\langle u, \phi' \rangle + a(u, \phi) ds = \int_0^T \langle f(u(s)), \phi \rangle + \langle \delta, \phi \rangle ds + (u(0), \phi(0))$$

and applying Galerkin approach methods, we obtain

$$\int_0^T -\langle u_m, \phi' \rangle + a(u_m, \phi) ds = \int_0^T \langle P_m f(u_m(s)), \phi \rangle + \langle \delta, \phi \rangle ds + (u_0, \phi(0)).$$

Since  $u_m(0) = P_m u_0 \rightarrow u_0$  we have  $u(0) = u_0$ .

In order to prove the singularity of solution, Let  $u_0$  and  $v_0$  be initial values on  $H$ . Then, form  $w(t) = u(t) - v(t)$ ,

so that

$$\frac{dw}{dt} + Aw = f(u) - f(v), \quad w(0) = u_0 - v_0.$$

By multiplying it with  $w$  and integrating it along  $\Omega$ , we obtain

$$\frac{1}{2} \frac{d}{dt} |w|^2 + \|w\|^2 = (f(u) - f(v), u - v).$$

It is worth noted that the boundary on  $f'$  is found in Equation (6). This prove that

$$\begin{aligned} (f(u) - f(v), u - v) &= \int_{\Omega} [f(u(x)) - f(v(x))] (u(x) - v(x)) dx \\ (f(u) - f(v), u - v) &= \left( \int_{v(x)}^{u(x)} f'(s) ds \right) (u(x) - v(x)) dx \\ &\leq l |u(x) - v(x)|^2 dx \\ &\leq l |u - v|^2 \end{aligned}$$

hence

$$\frac{1}{2} \frac{d}{dt} |w|^2 \leq l |u - v|^2.$$

we then integrate to acquire

$$|u(x) - v(x)| \leq |u_0 - v_0| e^{lt}.$$

We prove the singularity if  $u_0 = v_0$  an also depend on the continuity of the initial condition.

#### Acknowledgment

This work was financially supported by the LP2M (Lembaga Penelitian dan Pengabdian pada Masyarakat) of Hasanuddin University Makassar and The Higher Education Ministry of Indonesia.

### References

- [1] Zeidler E 1990 *Nonlinear Functional Analysis and its Applications II/B: Nonlinear Monotone Operators* Springer Science+Business Media, LLC.
- [2] Qiang L, Zheng Y, Wenshu Z 2007 Existence and some properties of weak solutions for a singular nonlinear parabolic equations. *Math. Meth. Appl. Sci.* **30** 1329–53
- [3] Robinson R C 2001 *Infinite-Dimensional Dynamical Systems: An Introduction to Dissipative Parabolic PDEs and the Theory of Global Attractor* Cambridge Texts in Applied Mathematics.
- [4] Choe H J, Lee J H 1998 Cauchy problem for nonlinear parabolic equations *Hokkaido Mathematical Journal* **27** 751-759.
- [5] Shang H F 2011 On the Cauchy problem for the singular parabolic equations with gradient term, *J. Math. Anal. Appl.* **378** 578–591.
- [6] Shang H F 2014 Cauchy problem for nonlinear parabolic equations with a gradient term, *J. Differential Equations* **257** 2801–25.
- [7] Nakao M, Aris N 2007 On global attractor for nonlinear parabolic equations of m-Laplacian type *J. Math. Anal. Appl.* **331** 793–809.
- [8] Jakhongir J R 2015 The Properties of the Solutions for Cauchy Problem of Nonlinear Parabolic Equations in Non-Divergent Form with Density *Journal of Siberian Federal University Mathematics & Physics* **8(2)** 192-200.

# On Weak Solution of Cauchy Problem for Nonlinear Parabolic Equations

---

## ORIGINALITY REPORT

---

%**9**

SIMILARITY INDEX

%**3**

INTERNET SOURCES

%**7**

PUBLICATIONS

%**5**

STUDENT PAPERS

---

## PRIMARY SOURCES

---

**1**

Guo, . "One-Dimensional Landau–Lifshitz Equations", Frontiers of Research with the Chinese Academy of Sciences, 2008.

Publication

%**1**

---

**2**

Ya. Belopolskaya. "Probabilistic approach to solution of nonlinear PDES arising in financial mathematics", Journal of Mathematical Sciences, 2010

Publication

%**1**

---

**3**

de Araujo, G.M.. "Unilateral problem for the Navier-Stokes operators with variable viscosity in noncylindrical domain", Nonlinear Analysis, 20060215

Publication

%**1**

---

**4**

Submitted to University of Stellenbosch, South Africa

Student Paper

%**1**

---

**5**

[egms-web.org](http://egms-web.org)

Internet Source

%**1**

---

6	"Numerical Analysis and Its Applications", Springer Nature, 2013 Publication	% 1
7	<a href="http://www.water.state.co.us">www.water.state.co.us</a> Internet Source	<% 1
8	Submitted to University College London Student Paper	<% 1
9	Haifeng Shang, Lihua Deng. "Local and global existence for quasilinear parabolic systems with a strongly nonlinear source", Nonlinear Analysis: Theory, Methods & Applications, 2012 Publication	<% 1
10	<a href="http://doc-test.utsp.utwente.nl">doc-test.utsp.utwente.nl</a> Internet Source	<% 1
11	Ning Ju. Nonlinearity, 05/2000 Publication	<% 1
12	<a href="http://www.j3.jstage.jst.go.jp">www.j3.jstage.jst.go.jp</a> Internet Source	<% 1
13	<a href="http://nanojournal.ifmo.ru">nanojournal.ifmo.ru</a> Internet Source	<% 1
14	<a href="http://ux1.mat.mfc.us.edu.pl">ux1.mat.mfc.us.edu.pl</a> Internet Source	<% 1
15	"Delay Differential Equations and Applications", Springer Nature, 2006 Publication	<% 1

16

Analysis and Optimization of Differential Systems, 2003.

Publication

<% 1

17

Asfaw, Teffera M.. "A new topological degree theory for pseudomonotone perturbations of the sum of two maximal monotone operators and applications", Journal of Mathematical Analysis and Applications, 2016.

Publication

<% 1

18

Submitted to Universitaet Trier

Student Paper

<% 1

19

Submitted to Indian Institute of Science Education and Research IISER, Trivandrum

Student Paper

<% 1

20

P. Rosenau, S. Kamin. "Non-linear diffusion in a finite mass medium", Communications on Pure and Applied Mathematics, 1982

Publication

<% 1

21

Submitted to Karunya University

Student Paper

<% 1

22

Klaus Gürlebeck, Klaus Habetha, Wolfgang Sprößig. "Application of Holomorphic Functions in Two and Higher Dimensions", Springer Nature, 2016

Publication

<% 1

---

EXCLUDE QUOTES ON

EXCLUDE  
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5  
WORDS