

43_-_Sep_2021_-
_Concordance_between_Triglyc
eride_Glucose_Index_and_Admi
ssion_Insulin_Resistance_Index_
in_Non-
diabetic_Subjects_of_Acute_Cor
onary_Syndrom.pdf

by

Submission date: 19-Jan-2023 01:17PM (UTC+0700)

Submission ID: 1995239344

File name: 43_-_Sep_2021_-
_Concordance_between_Triglyceride_Glucose_Index_and_Admission_Insulin_Resistance_Index_in_Non-
diabetic_Subjects_of_Acute_Coronary_Syndrom.pdf (406.94K)

Word count: 3286

Character count: 17420



Concordance between Triglyceride Glucose Index and Admission Insulin Resistance Index in Non-diabetic Subjects of Acute Coronary Syndrome

Andi Irham¹, Husaini Umar^{2*}, Fabiola Maureen Shinta Adam², Idar Mappangara³, Syakib Bakri⁴, Risna Halim⁵, Alfian Zainuddin⁶

¹Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia; ²Department of Internal Medicine, Division of Endocrine Metabolic and Diabetes, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia; ³Department of Internal Medicine, Division of Cardiology and Vascular Medicine, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia; ⁴Department of Internal Medicine, Division of Nephrology and Hypertension, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia; ⁵Department of Internal Medicine, Division of Tropical Medicine and Infectious Disease, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia; ⁶Department of Biostatistics, Faculty of Public Health, Hasanuddin University, Makassar, Indonesia

Abstract

Edited by: Ksenija Bogoeva-Kostovska
Citation: Irham A, Umar H, Adam FMS, Mappangara I, Bakri S, Halim R, Zainuddin A. Concordance between Triglyceride Glucose Index and Admission Insulin Resistance Index in Non-diabetic Subjects of Acute Coronary Syndrome. *Open Access Maced J Med Sci*. 2021 Oct 01; 9(B):1044-1047. <https://doi.org/10.3889/oamjms.2021.7075>
Keywords: Insulin resistance; Triglyceride glucose index; Admission insulin resistance index; Non-diabetic of acute coronary syndrome
*Correspondence: Husaini Umar, Division of Endocrine Metabolic and Diabetes, Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.
E-mail: husaini_umar65@yahoo.co.id
Received: 16-Aug-2021
Revised: 04-Sep-2021
Accepted: 21-Sep-2021
Copyright: © 2021 Andi Irham, Husaini Umar, Fabiola Maureen Shinta Adam, Idar Mappangara, Syakib Bakri, Risna Halim, Alfian Zainuddin
Funding: This research did not receive any financial support
Competing Interests: The authors declare that they have no competing interests.
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: Acute coronary syndrome (ACS) is the leading cause of death in the world and is closely related to insulin resistance (IR). The examination of the triglyceride glucose (TyG) index and admission IR index (AIRI) can be an alternative examination to assess IR.

AIM: The aim of the study was to analyze the concordance between the TyG index and AIRI in assessing IR in non-diabetic subjects of ACS.

METHODS: This study was a cross-sectional study using secondary data from ACS patient medical records in Wahidin Sudirohusodo Hospital Makassar from September 2020 until April 2021. An examination of RBG, TyG, and fasting insulin was performed. The statistical test results were considered significant if the $p < 0.05$.

RESULTS: This study included 67 subjects, of which 54 were male and 13 were female, there were 24 people aged <50 years, 43 people aged 50 years old, with a mean of 55.15 ± 12.71 years, a diagnosis of ST-elevation myocardial infarction (STEMI) 25 people, non-STEMI 34 people and unstable angina pectoris 8 people, $p < 0.05$. The receiver operating characteristic curve showed that the cut-off value of the TyG index in predicting AIRI was 3141, with a sensitivity of 88.98% and specificity of 59.09%. The OR value revealed that subjects with a TyG index of >9.06 indicated an 11-fold greater risk for IR than subjects with a TyG index ≤ 9.06 .

CONCLUSIONS: There was a significant concordance between the TyG index and the AIRI in non-diabetic acute coronary syndrome patient.

Introduction

Cardiovascular disease is the leading cause of death in the world and is often associated with various comorbidities such as obesity, abnormal lipid profile, and insulin resistance (IR) [1], [2]. The most common cardiovascular disease is acute coronary syndrome (ACS). ACS is divided into three groups, namely, ST-elevation myocardial infarction (STEMI), non-STEMI (NSTEMI), and unstable angina (UA) [3].

The high population with chronic disease associated with IR has been widely reported worldwide [4]. The mechanism of IR in the heart is complex and multifactorial. Impaired insulin-stimulated glucose absorption is still a common problem, changes in signaling kinases can be increased or decreased,

thus impacting cardiac structure and function. IR generally occurs mainly as a result of obesity, excess calories, lack of physical activity, genetics, and age. IR is associated with many serious medical conditions such as type 2 diabetes mellitus, hypertension, atherosclerosis, and metabolic syndrome (MetS) [5].

Systemic inflammation also contributes to IR, proinflammatory cytokines induce IR which can affect glucose and fatty acid utilization by inducing inflammation increases the heart's dependence on triglycerides from the liver and free fatty acids from adipose tissue for energy [6]. Obesity can increase circulating concentrations of inflammatory cytokines, such as interleukin-6 tumor necrosis factor- α and leptin [2].

Hyperinsulinemia Euglycemic Clamp is still the gold standard to diagnose IR, but this method is expensive, invasive, and difficult to perform in health

facilities [16]. There are various kinds of IR tests, such as quantitative insulin sensitivity check index, fasting glucose to insulin ratio), dan insulin-modified feasibility intravenous glucose tolerance test but each of these methods is not currently available in all health services [8]. Alternative examination that has relatively low cost, fast results and can easily predict RI is admission IR index (AIRI) and triglyceride glucose (TyG) index [5], [9]. Even though AIRI is expensive than TyG Index, but the subjects does not need to fast.

Luo et al. conducted a study of 1092 patients with STEMI and received percutaneous coronary intervention. This study divided patients into 4 groups according to their level of TyG Index, the patients were then conducted a study for 1 year. The results showed that the higher the TyG Index, the higher the incidence of STEMI [10]. The study of Aman et al. showed the TyG index in predicting HOMA-IR was 4.66, with the sensitivity of 86.2% and specificity of 44.1%. The OR value revealed that subjects with TyG index of ≥ 4.66 indicated a 5-time greater risk for IR than subjects with a TyG index < 4.66 [11].

This study aims to investigate the concordance between the TyG index and AIRI in assessing IR in nondiabetic subjects of ACS.

Methods

Research design

This study was cross-sectional study using secondary data from ACS patient medical records in Wahidin Sudirohusodo Hospital Makassar from September 2020 until April 2021.

Research subject

Research subjects were taken with consecutive sampling from all patients who entered the emergency room Wahidin Sudirohusodo Hospital Makassar from September 2020 until April 2021 and met the research criteria. Inclusion criteria were ACS subjects aged > 18 years old, no history of diabetic, and willing to take part in research. Exclusion criteria were when the subject aged ≤ 18 years old and refuse to take part in research.

Research data collection

Research subjects were non-diabetic subjects of ACS who met the research criteria then we collected their triglyceride levels, RBG levels, insulin levels (without fasting) and used formula to determined TyG index and AIRI subjects.

Data analysis

Data analysis was performed using SPSS. The statistical test used Anova test if the data is normally distributed or the Kruskal Wallis test if the data is not normally distributed. To find the correlation between AIRI and TyG use the Pearson Correlation test if the data is normally distributed or the Spearman Correlation test if the data is not normally distributed. The results were considered significant if the $p < 0.05$.

Research permit and ethical eligibility

This research protocol has been approved by the Health Research Ethics Committee, Medical Faculty, Hasanuddin University Makassar, according to the ethical approval recommendation letter number: 165/UN4.6.4.5.31/PP36/2020.

Results

The study was conducted at the emergency unit installation and the treatment installation at Wahidin Sudirohusodo Hospital and a network hospital from September 2020 to April 2021. This study involved 67 subjects who met the inclusion criteria. Statistical analysis carried out was descriptive statistical calculations and statistical tests of spearman correlation, and receiver operating characteristic (ROC) curve. The results of statistical tests were significant if $p < 0.05$.

Data analysis was conducted on 67 subjects of which 54 were male and 13 were female. The incidence of heart disease will increase at the age of 50 years, 39 in this study found 24 people aged < 50 years, 50 years 43 people, with a mean of 55.15 ± 12.71 years, a diagnosis of STEMI 25 people, NSTEMI 34 people and unstable angina pectoris (UAP) 8 people. The characteristics of the research subjects had fasting blood glucose (FBG) 108.04 ± 20.52 , TG 141.22 ± 52.67 , AIRI 28.87 ± 15.98 , and TyG index 8.86 ± 0.39 (Table 1).

Table 1: Sample characteristic (n=67)

Variables	n (mean \pm SD)	%/median (min-max)
Gender		
Male	54	80.6
Female	13	9.4
Age (years)		
< 50	24	35.8
≥ 50	43	63.2
Diagnosis		
ST elevation myocardial infarction	25	37.3
Non-ST elevation myocardial infarction	34	50.7
Unstable angina pectoris	8	11.9
Triglyceride	141.22 ± 52.67	132 (58–330)
RBG	134.46 ± 27.60	129 (81–220)
Fasting blood glucose	108.04 ± 20.52	(68–191)
Admission insulin resistance index	28.87 ± 15.98	25.39 (8.51–98.76)
Triglyceride glucose	8.86 ± 0.39	8.86 (7.80–9.84)

Figure 1 shows the results of the ROC curve analysis showed the area under curve value was 0.799




Figure 1: Receiver operating characteristic curve of triglyceride glucose index on homeostasis model assessment of insulin resistance

($p < 0.01$). This means that the TyG Index is quite sensitive in predicting the AIRI value with a cutoff value of 31.41. Figure 2. illustrates the positive correlation between the TyG index and AIRI ($p = 0.001$) with a correlation strength of 0.359 is in the category of moderate correlation and the direction is positive (the higher the value of AIRI, the higher the value of TyG or otherwise).




Figure 2: Positive correlation between triglyceride glucose index and admission insulin resistance index

Table 2, shows that there is a positive correlation between TyG and AIRI index with $p < 0.001$, sensitivity reached 88.80% and specificity 59.09%. Based on OR, subjects with TyG index >9.06 had an 11.556 times greater risk of experiencing RI than subjects with TyG index 8062.

Table 2: Triglyceride glucose index with admission insulin resistance index based on receiver operating characteristic curve

Triglyceride glucose	Admission insulin resistance index		n	p value	Sensitivity	Specify	OR
	≤ 31.41	> 31.41					
≤ 9.06	40	9	49	0.001	88.80%	59.09%	11.556
> 9.06	5	13	18				
n	45	22	67				

The number of subjects of this study was 67 which 54 were male and 13 were female Non-Diabetic subjects ACS. The incidence of heart disease will increase at the age of 50 years [12], in this study we found 24 people aged <50 years, ≥ 50 years 43 people, with a mean of 55.15 ± 12.71 years, a diagnosis of STEMI 25 people, NSTEMI 34 people and UAP 8 people. The characteristics of the research subjects had FBG (108.04 ± 20.52 , TG 141.22 ± 52.67 , AIRI 28.87 ± 15.98 , and TyG index 8.86 ± 0.39).

In this study, the relationship between AIRI and TyG was found where ($p = 0.001$) with a correlation strength of 0.359 was in the category of moderate correlation and the direction was positive. From the results of this study, the TyG index and AIRI can be used as markers to detect resistance. IR can increase the risk of hyperglycemia, hypertension, and dyslipidemia which increases the risk of inflammation, changes in coagulation factors, atherosclerosis, and coronary heart disease death [13], [14]. The TyG index and AIRI are easy tests to detect the presence of subjects who are at high risk of suffering from cardiovascular disease, high triglyceride levels and fasting glucose are components of MetS which is one of the most important markers of heart disease [15]. A study conducted by Lee *et al.* concluded that the TyG index was strongly associated with an increased risk of ACS in asymptomatic subjects with type 2 diabetes, especially if they had cardiovascular risk factors [16]. The other study research by Hong *et al.* which uses the TyG index as a tool to detect the presence of subjects with high cardiovascular risk [17]. AIRI examination itself conducted by Refaie and Elewa A showed the results that the increase in AIRI can predict coronary artery events in non-diabetic patients with acute chest pain. Multiple coronary vessel involvement is common in such cases and an appropriate planned invasive therapy strategy should be considered [18]. Another study conducted by Abdulla *et al.* proved an increase in AIRI in patients with ACS [19].

In this study, we divided the AIRI value and the TyG index value based on the tertile. Tertile values 1 and 2 AIRI ≤ 31.41 tertile 3 >31.41 . On the other hand, Rosalina *et al.* in Makassar concluded that in obese patients, the highest percentage of AIRI tertile 3 (70%) was found compared to AIRI tertile 2 (20%) and AIRI tertile 1 (15%). Meanwhile, those who were not obese were found to have the highest levels of 1 AIRI (85%) compared to 2 AIRI (80%) and 3 AIRI (30%) [9]. Another study by Refaie and Elewa found that AIRI value significantly increased in acute myocardial infarction (3.9 ± 0.1) and UAP (3.01 ± 0.2) when compared with the stable angina and control groups [18].

The average value of the TyG index in this study was 8.86 ± 0.39 and the tertile value of TyG index tertile 1 and 2 ≤ 9.06 tertile 3 >9.06 . Research conducted by Aman *et al.* in Makassar got a value of $4.79 \pm 0.25.40$ [11]. The other research conducted

by Wang *et al.* in South Korea divided the TyG index value into three tertiles, tertile 1 (8.45 ± 0.29), tertile 2 (9.09 ± 0.15), and tertile 3 (9.94 ± 0.52) [20].

Conclusion

There was a significant concordance between the TyG index and the AIRI in non-diabetic subjects of ACS patients. TyG index in predicting AIRI was 3141, with a sensitivity of 88.98% and specificity of 59.09%. The OR value revealed that subjects with a TyG index of >9.06 indicated a 11-time greater risk for IR than subjects with a TyG index ≤ 9.06 .

Study limitation

The limitation of the study was at the blood sample was taken only twice., first at the time of admission to the hospital and when patient was fasting. The sensitivity and specificity of the TyG index that was compared only with AIRI, which is not the gold standard in diagnosing IR.

References

- Jan S, Lee SW, Sawhney JP. Catastrophic health expenditure on acute coronary events in Asia: A prospective study. *Bull World Health Organ.* 2016;94(3):193-200. <https://doi.org/10.2471/blt.15.158303>
PMid:26966330
- Ormazabal V, Nair S, Elfeky O, Aguayo C, Salomon C, Zuñiga FA. Association between insulin resistance and the development of cardiovascular disease. *Cardiovasc Diabetol.* 2018;17(1):1- 14. <https://doi.org/10.1186/s12933-018-0762-4>
- Juzar DA, Dany SS, Irmalita I, Tobing DPL, Firdaus F, Widyantor B. *Pedoman Tatalaksana Sindrom Koroner Akut. Perhimpun Dokter Spesialis Kardiovaskuler Indonesia.* Jakarta: Interna Publishing; 2015.
- Lim J, Kim J, Kwon GC, Koo SH. Comparison of triglyceride glucose index, and related parameters to predict insulin resistance in Korean adults: An analysis of the 2007-2010 Korean National Health and Nutrition Examination Survey. *PLoS One.* 2019;14(3):e0212963. <https://doi.org/10.1371/journal.pone.0212963>
PMid:30845237
- Abel ED, O'Shea KM, Ramasamy R. Insulin resistance: Metabolic mechanisms and consequences in the heart. *Arterioscler Thromb Vasc Biol.* 2012;32(9):2068-76. <https://doi.org/10.1161/atvbaha.111.241984>
PMid:22895668
- Chen L, Chen R, Wang H, Liang F. Mechanisms linking inflammation to insulin resistance. *Int J Endocrinol.* 2015;2015:508409.
PMid:26136779
- De Souza AL, Batista GA, Alegre SM. Assessment of insulin sensitivity by the hyperinsulinemic euglycemic clamp: Comparison with the spectral analysis of photoplethysmography. *J Diabetes Complications.* 2017;31(1):128-33. <https://doi.org/10.1016/j.jdiacomp.2016.10.018>
PMid:27839921
- Conwell LS, Trost SG, Brown WJ, Batch JA. Indexes of insulin resistance and secretion in obese children and adolescents: A validation study. *Diabetes Care.* 2004;27(2):314-9. <https://doi.org/10.2337/diacare.27.2.314>
PMid:14747206
- Rosalina R, Umar H, Tandean P, Kasim H, Aman AM, Rasyid H, *et al.* The comparison of AIRI values in obese and non obese acute coronary syndrome patients. *Eurol J Mol Clin Med.* 2020;7(8):190-7.
- Luo E, Wang D, Yan G, Qiao Y, Liu B, Hou J, *et al.* High triglyceride-glucose index is associated with poor prognosis in patients with acute ST-elevation myocardial infarction after percutaneous coronary intervention. *J Cardiovasc Diabetol.* 2019;18(1):2-12. <https://doi.org/10.1186/s12933-019-0957-3>
PMid:31722708
- Aman M, Resnawita D, Rasyid H, Kasim H, Bakri S, Umar H, *et al.* The concordance of triglyceride glucose index (TyG index) and homeostatic model assessment for insulin resistance (Homa-IR) in non-diabetic subjects of adult Indonesian males. *Clin Epidemiol Glob Health.* 2021;9:227-30. <https://doi.org/10.1016/j.cegh.2020.09.003>
- Mozaffarian D, Benjamin JE, Amett KD. Heart Disease an stroke statistics. *Circ Aha J.* 2015;131(3):1-323.
- Bonura F, Vitale G. Insulin resistance and acute coronary syndrome. *Atherosclerosis.* 2010;211(2):672-5.
PMid:20466373
- Kim MK, Ahn CW, Kang S, Nam JS, Kim KR, Park JS. Relationship between the triglyseride glucose index and coronary artery calcification in Korean adults. *J Cardiovasc Diabetol.* 2017;16(1):1-7. <https://doi.org/10.1186/s12933-017-0589-4>
PMid:28830471
- Mao Q, Zhou D, Li Y, Wang Y, Wang Y, Xu SC, Zhao XH. The triglyceride-glucose index predicts coronary artery disease severity and cardiovascular outcomes in patients with non-ST-segment elevation acute coronary syndrome. *Dis Markers.* 2019;2019:6891537. <https://doi.org/10.1155/2019/6891537>
PMid:31281548
- Lee EY, Yang HK, Lee J, Kang B, Yang Y, Lee SH, *et al.* Triglyceride glucose index, a marker of insulin resistance, is associated with coronary artery stenosis in asymptomatic subjects with Type 2 diabetes. *Lipids Health Dis.* 2016;15(1):155. <https://doi.org/10.1186/s12944-016-0324-2>
PMid:27633375
- Hong S, Han K, Park CY. The triglyceride glucose index is a simple and low-cost marker associated with atherosclerotic cardiovascular disease: A population-based study. *BMC Med.* 2020;18(1):1-8. <https://doi.org/10.1186/s12916-020-01824-2>
- Refaie W, Elewa A. Admission insulin resistance index in non diabetic patients with acute coronary syndrome; clinical and angiographic features. *Egypt Hear J.* 2013;65(4):301-5. <https://doi.org/10.1016/j.ehj.2013.02.004>
- Abdullah I, Hasan G, Altunkeser BB, Ozdemir K, Gürbilek M, Gederet YT, *et al.* Evaluation of "admission index of insulin resistance (AIRI)" as an early stage risk predictor in non-diabetic acute coronary syndromes. *Anatol J Cardiol.* 2002;2(3):194-201.
PMid:12223324
- Wang S, Shi J, Peng Y, Fang Q, Mu Q, Gu W, *et al.* Stronger association of triglyceride glucose index than the HOMA-IR with arterial stiffness in patients with Type 2 diabetes: A real-world single-centre study. *Cardiovasc Diabetol.* 2021;20(1):1-10. <https://doi.org/10.1186/s12933-021-01274-x>

ORIGINALITY REPORT

25%
SIMILARITY INDEX

20%
INTERNET SOURCES

19%
PUBLICATIONS

3%
STUDENT PAPERS

PRIMARY SOURCES

1	atvb.ahajournals.org Internet Source	3%
2	academic-accelerator.com Internet Source	2%
3	insights.ovid.com Internet Source	2%
4	Submitted to iGroup Student Paper	1%
5	journals.plos.org Internet Source	1%
6	www.science.gov Internet Source	1%
7	Longlong Hu, Huihui Bao, Xiao Huang, Wei Zhou, Tao Wang, Lingjuan Zhu, Xi Liu, Minghui Li, Xiaoshu Cheng. "Relationship Between the Triglyceride Glucose Index and the Risk of First Stroke in Elderly Hypertensive Patients",	1%

International Journal of General Medicine, 2022

Publication

8

www.ncbi.nlm.nih.gov

Internet Source

1 %

9

Haipeng Yao, Zhen Sun, Wei Yuan, Chen Shao, Honghua Cai, Lihua Li, Yongjiang Qian, Zhongqun Wang. "Relationship Between the Triglyceride-Glucose Index and Type 2 Diabetic Macroangiopathy: A Single-Center Retrospective Analysis", Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2022

Publication

1 %

10

Valeska Ormazabal, Soumyalekshmi Nair, Omar Elfeky, Claudio Aguayo, Carlos Salomon, Felipe A. Zuñiga. "Association between insulin resistance and the development of cardiovascular disease", Cardiovascular Diabetology, 2018

Publication

1 %

11

www.nature.com

Internet Source

1 %

12

Qi Mao, Denglu Zhou, Youmei Li, Yuqing Wang, Shang-Cheng Xu, Xiao-Hui Zhao. "The Triglyceride-Glucose Index Predicts Coronary Artery Disease Severity and Cardiovascular Outcomes in Patients with Non-ST-Segment

1 %

Elevation Acute Coronary Syndrome", Disease Markers, 2019

Publication

13

Submitted to Aspen University

Student Paper

1 %

14

repository.unam.edu.na

Internet Source

1 %

15

worldwidescience.org

Internet Source

1 %

16

M. M. Nur, I. M. Newman, L. M. Siqueira.
"Glucose Metabolism in Overweight Hispanic
Adolescents With and Without Polycystic
Ovary Syndrome", PEDIATRICS, 2009

Publication

1 %

17

www.hcplive.com

Internet Source

1 %

18

Dominika Drwiła, Paweł Rostoff, Grzegorz
Gajos, Jadwiga Nessler, Ewa Konduracka.
"Prognostic value of the triglyceride-glucose
index among non-diabetic patients with acute
myocardial infarction at one-year follow-up",
Kardiologia Polska, 2021

Publication

<1 %

19

pure.eur.nl

Internet Source

<1 %

20 Da-Hye Son, Hye Sun Lee, Yong-Jae Lee, Jun-Hyuk Lee, Jee-Hye Han. "Comparison of triglyceride-glucose index and HOMA-IR for predicting prevalence and incidence of metabolic syndrome", Nutrition, Metabolism and Cardiovascular Diseases, 2021
Publication

21 bmccardiovascdisord.biomedcentral.com
Internet Source

22 lipidworld.biomedcentral.com
Internet Source

23 www.scienceopen.com
Internet Source

24 Yue Zhang, Xiaosong Ding, Bing Hua, Qingbo Liu, Hui Gao, Hui Chen, Xue-Qiao Zhao, Weiping Li, Hongwei Li. "High triglyceride-glucose index is associated with adverse cardiovascular outcomes in patients with acute myocardial infarction", Nutrition, Metabolism and Cardiovascular Diseases, 2020
Publication

25 elearning.medistra.ac.id
Internet Source

26 media.neliti.com
Internet Source

27

Chengping Hu, Jianwei Zhang, Jinxing Liu, Yan Liu, Ang Gao, Yong Zhu, Yingxin Zhao. "Discordance between the triglyceride glucose index and fasting plasma glucose or HbA1C in patients with acute coronary syndrome undergoing percutaneous coronary intervention predicts cardiovascular events: a cohort study from China", *Cardiovascular Diabetology*, 2020

Publication

<1 %

28

Luis E. Simental-Mendía, Martha Rodríguez-Morán, Fernando Guerrero-Romero. "The Product of Fasting Glucose and Triglycerides As Surrogate for Identifying Insulin Resistance in Apparently Healthy Subjects", *Metabolic Syndrome and Related Disorders*, 2008

Publication

<1 %

29

Wen Guo, Jing Lu, Pei Qin, Xiaona Li, Wenfang Zhu, Juan Wu, Nianzhen Xu, Qun Zhang. "The triglyceride-glucose index is associated with the severity of hepatic steatosis and the presence of liver fibrosis in non-alcoholic fatty liver disease: a cross-sectional study in Chinese adults", *Lipids in Health and Disease*, 2020

Publication

<1 %

30

Wenqin Guo, Lingyue Zhao, Fanrui Mo, Changnong Peng, Lang Li, Yan Xu, Wenyu

<1 %

Guo, Aimei Sun, Hongbing Yan, Lili Wang. "The prognostic value of the triglyceride glucose index in patients with chronic heart failure and type 2 diabetes: A retrospective cohort study", *Diabetes Research and Clinical Practice*, 2021

Publication

31

Yaling Li, Rui Zheng, Jie Li, Shuyi Feng, Li Wang, Zhiming Huang. "Association between triglyceride glucose-body mass index and non-alcoholic fatty liver disease in the non-obese Chinese population with normal blood lipid levels: a secondary analysis based on a prospective cohort study", *Lipids in Health and Disease*, 2020

Publication

<1 %

32

ouci.dntb.gov.ua

Internet Source

<1 %

33

users.humboldt.edu

Internet Source

<1 %

34

M. J. J. Finken, M. G. Keijzer-Veen, F. W. Dekker, M. Frölich, E. T. M. Hille, J. A. Romijn, J. M. Wit. "Preterm birth and later insulin resistance: effects of birth weight and postnatal growth in a population based longitudinal study from birth into adult life", *Diabetologia*, 2006

Publication

<1 %

35 Stalin Viswanathan, Yeshwanth Mohan Yalavarthy. "Burden and outcomes of chronic kidney disease in patients presenting with hypertensive crisis", Research Square Platform LLC, 2023

Publication

<1 %

36 www.karger.com

Internet Source

<1 %

37 Jihyun Yoon, Donghyuk Jung, Yongjae Lee, Byoungjin Park. "The Metabolic Score for Insulin Resistance (METS-IR) as a Predictor of Incident Ischemic Heart Disease: A Longitudinal Study Among Korean Without Diabetes", Journal of Personalized Medicine, 2021

Publication

<1 %

38 Yunfan Tian, Jingyu Sun, Ming Qiu, Yan Lu, Xuesong Qian, Wei Sun, Xiangqing Kong. "Association between the triglyceride-glucose index and albuminuria in hypertensive individuals", Clinical and Experimental Hypertension, 2022

Publication

<1 %

39 Hanna Luze, Anna Schwarz, Sebastian Philipp Nischwitz, Dagmar Kolb et al. "Autologous Fat Grafting in Reconstructive Breast Surgery: Clinically Relevant Factors Affecting the Graft Take", Aesthetic Surgery Journal, 2022

<1 %

40

Hye Jin Joo, Gyu Ri Kim, Eun-Cheol Park, Sung-In Jang. "Association between Frequency of Breakfast Consumption and Insulin Resistance Using Triglyceride-Glucose Index: A Cross-Sectional Study of the Korea National Health and Nutrition Examination Survey (2016–2018)", *International Journal of Environmental Research and Public Health*, 2020

<1 %

Publication

41

Ki-Woong Nam, Hyung-Min Kwon, Han-Yeong Jeong, Jin-Ho Park, Hyuktae Kwon, Su-Min Jeong. "High triglyceride-glucose index is associated with subclinical cerebral small vessel disease in a healthy population: a cross-sectional study", *Cardiovascular Diabetology*, 2020

<1 %

Publication

42

Xiang Wang, Wei Xu, Qirui Song, Zinan Zhao, Xuyang Meng, Chenxi Xia, Yibo Xie, Chenguang Yang, Pengfei Jin, Fang Wang. "Association between the triglyceride–glucose index and severity of coronary artery disease", *Cardiovascular Diabetology*, 2022

<1 %

Publication

43

Yingkai Li, Songyuan He, Zheng Wu, Wenzheng Li, Wen Jian, Zichao Cheng, Cong Wang, Yuchen Shi, Jinghua Liu. "The predictive

<1 %

value of the triglyceride—glucose index for
cardiovascular events in patients with
coronary chronic total occlusion",
Cardiovascular Diabetology, 2022

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On