

3._Isra_genomic.pdf

by Isra Wahid

FILE	3._ISRA_GENOMIC.PDF (2.52M)	WORD COUNT	11474
TIME SUBMITTED	22-AUG-2019 07:23AM (UTC+0700)	CHARACTER COUNT	57780
SUBMISSION ID	1162163047		



Contents lists available at ScienceDirect

Infection, Genetics and Evolution

journal homepage: www.elsevier.com/locate/meegid



Genomic analysis and growth characteristic of dengue viruses from Makassar, Indonesia

R. Tedjo Sasmono^{a,b,*}, Isra Wahid^{a,c,2}, Hidayat Trimarsanto^b, Benediktus Yohan^{a,b,1}, Sitti Wahyuni^{a,c,2}, Martin Hertanto^{a,3}, Irawan Yusuf^c, Halim Mubin^c, Idham J. Ganda^c, Rachmat Latief^c, Pablo J. Bifani^d, Pei-Yong Shi^d, Mark J. Schreiber^{d,4}

^a Novartis-Eijkman-Hasanuddin Clinical Research Initiative (NEHCRI), Indonesia
^b Eijkman Institute for Molecular Biology, Jl. Diponegoro 69, Jakarta 10430, Indonesia
^c Faculty of Medicine, Hasanuddin University, Jl. Perintis Kemerdekaan 10, Tamalanrea, Makassar 90245, Indonesia
^d Novartis Institute for Tropical Diseases (NITD), 10 Biopolis Road, Chromos #05-01, Singapore 138670, Singapore

ARTICLE INFO

Article history:
Received 7 January 2015
Received in revised form 5 March 2015
Accepted 6 March 2015
Available online xxxx

Keywords:
Dengue
Genotype
Indonesia
Genome
Growth kinetic

ABSTRACT

Dengue fever is currently the most important mosquito-borne viral disease in Indonesia. In South Sulawesi province, most regions report dengue cases including the capital city, Makassar. Currently, no information is available on the serotypes and genotypes of the viruses circulating in the area. To understand the dynamic of dengue disease in Makassar, we carried out dengue fever surveillance study during 2007–2010. A total of 455 patients were recruited, in which antigen and serological detection revealed the confirmed dengue cases in 43.3% of patients. Molecular detection confirmed the dengue cases in 27.7% of patients, demonstrating that dengue places a significant disease burden on the community. Serotyping revealed that dengue virus serotype 1 (DENV-1) was the most predominant serotype, followed by DENV-2, -3, and -4. To determine the molecular evolution of the viruses, we conducted whole-genome sequencing of 80 isolates. Phylogenetic analysis grouped DENV-2, -3 and -4 to the Cosmopolitan genotype, Genotype I and Genotype II, respectively. Intriguingly, each serotype paints a different picture of evolution and transmission. DENV-1 appears to be undergoing a clade replacement with Genotype IV being supplanted by Genotype I. The Cosmopolitan DENV-2 isolates were found to be regionally endemic and is frequently being exchanged between countries in the region. By contrast, DENV-3 and DENV-4 isolates were related to strains with a long history in Indonesia although the DENV-3 strains appear to have been following a distinct evolutionary path since approximately 1998. To assess whether the various DENV serotypes/genotypes possess different growth characteristics, we performed growth kinetic assays on selected viruses. We observed the relatively higher rate of replication for DENV-1 and -2 compared to DENV-3 and -4. Within the DENV-1, viruses from Genotype I grow faster than that of Genotype IV. This higher replication rate may underlie their ability to replace the circulation of Genotype IV in the community.

© 2015 Published by Elsevier B.V.

1. Introduction

Dengue fever is an acute febrile disease caused by dengue virus (DENV) infection, transmitted through the mosquito vector *Aedes aegypti*. The symptoms of the DENV infection vary from classic Dengue Fever (DF) to severe forms such as Dengue Hemorrhagic Fever (DHF) and the potentially fatal Dengue Shock Syndrome (DSS). DENV is an enveloped flavivirus with a 10.7 kb single-stranded positive-sense RNA genome. There are four antigenically distinct serotypes of the virus (DENV-1, -2, -3, and -4), with around 65% genome similarity, circulating throughout the tropical and sub-tropical regions of the world (Gubler, 1998; Guzman et al.,

* Corresponding author at: Eijkman Institute for Molecular Biology, Jakarta 10430, Indonesia. Tel.: +62 21 3917131; fax: +62 21 3147982.

E-mail address: sas@eijkman.go.id (R.T. Sasmono).

¹ Present address: Eijkman Institute for Molecular Biology, Jakarta 10430, Indonesia.

² Present address: Faculty of Medicine, Hasanuddin University, Makassar 90245, Indonesia.

³ Present address: Cordlife Persada, Jakarta, Indonesia.

⁴ Present address: Merck, 470 Atlantic Avenue, Boston, United States.

<http://dx.doi.org/10.1016/j.meegid.2015.03.006>
1567-1348/© 2015 Published by Elsevier B.V.

2010). The disease caused by the four serotypes is symptomatically identical.

Indonesia is a dengue endemic country that experiences periodic major outbreaks of the disease, such as those in 1998 (Corwin et al., 2001) and 2004 (Suwandono et al., 2006). In addition, annual and sporadic occurrences regularly afflict all the 33 provinces of the vast Indonesian archipelago (Sumarmo, 1987). While there have been studies describing the molecular genetics of dengue viruses isolated from Western part of the Indonesian archipelago (Sumatra and Java Islands) (Ong et al., 2008; Raekiansyah et al., 2005), no data is available for other major islands in the Eastern part of Indonesia, such as Sulawesi island. Data from the South Sulawesi provincial health authority (www.dinkes-sulsel.go.id) described the increasing number of dengue incidents in South Sulawesi province in 2007, in which 5333 patients were admitted to hospitals. In the capital city, Makassar, the dengue incidence was recorded at 425 cases with 5 fatalities. Dengue diagnosis in Makassar is commonly performed based on clinical symptoms only as the lack of resources and relatively high cost of laboratory tests prohibit serological confirmation. Because of this, the reported incidence may not represent the true disease burden.

Comparative analysis of dengue genomes has been used widely to study the genetic diversity of the dengue viruses (Holmes and Burch, 2000; Holmes and Twiddy, 2003) and previous studies have also suggested a correlation between viral pathogenicity with its genetic structures in which mutations in viral genome enhances the viral pathogenesis (Leitmeyer et al., 1999; Pandey and Igarashi, 2000). To understand the dengue disease dynamic in Makassar city, Indonesia, we studied the molecular evolution of circulating dengue viruses by analyzing the genomes of the viruses isolated in the area. We observed the diversity of serotypes and genotypes of dengue viruses and hypothesized that spatial isolation may contribute to the diversity of dengue viruses circulating in Makassar, South Sulawesi, Indonesia. To complement the viral genetic data, we performed viral growth characterization of various serotypes/genotypes. This study provides information in the genetic and biological characteristics of the diverse dengue virus strains circulating in Makassar, Indonesia.

2. Materials and methods

2.1. Study site and patient recruitment

We conducted surveillance of febrile patients in Makassar, the capital city of South Sulawesi province, Indonesia. The city resides in an area of about 175.77 km², inhabited by about 1.3 million people (<http://makassarkota.go.id>). Makassar city is located about 1400 km east of the Indonesian capital of Jakarta. Dengue-suspected febrile patients admitted to Dr. Wahidin Sudirohusodo, Labuang Baji and Daya Hospitals and primary health care centers in Makassar were enrolled in the study upon obtaining written consent. For the minors/children participants, written consents were obtained from parents/legal guardians. We obtained ethical clearance for this study from the Hasanuddin University Medical Research Ethics Committee and the Eijkman Institute Research Ethics Committee. All patients aged between 5 and 100 years old with fever >38.0 °C for less than 72 h and without concurrent signs of upper respiratory infection or obvious alternate diagnoses to dengue were invited to participate in the study. In total, we obtained 455 serum samples during the patient recruitment period (May 2007–August 2010).

Initial screening of dengue-suspected patients was determined serologically using anti-dengue IgM and IgG (Panbio Dengue Duo IgM/IgG enzyme-linked immunosorbent assay (ELISA) kit

(Panbio, Brisbane, Australia)). Dengue confirmation by detection of NS1 antigen was done using NS1 rapid test kit (Standard Diagnostics, Korea). Primary versus secondary dengue infection was determined using the ELISA results according to manufacturer's protocol. Briefly, the positive IgM (>11 Panbio Units) and negative IgG (<22 Panbio Units) indicated primary infection while positive IgG (>22 Panbio Units), which may be accompanied by elevated IgM levels, indicated secondary infection. The presence of viral RNA in NS1-positive samples was further confirmed by reverse transcription-polymerase chain reaction (RT-PCR) using method described by Lanciotti et al. (1992) (Lanciotti et al., 1992), with modification according to Harris et al. (1998). All dengue positive cases were categorized either dengue fever (DF) or Dengue Hemorrhagic Fever (DHF) according to criteria described by the WHO (WHO, 1997).

2.2. Virus propagation, cell lines, RNA extraction and serotyping

Viruses were isolated from RT-PCR positive samples, either directly from patients' sera or after one passage (or maximum of two passages for later isolates) of virus propagation in C6/36 *Aedes albopictus* cell lines. Baby hamster kidney (BHK-21) and C6/36 cell lines are maintained in RPMI 1640 medium (Gibco, Carlsbad, CA). African green monkey kidney (Vero76) cell line was maintained in Minimum Essential Medium (MEM) with Earle's salt (Gibco). Media were supplemented with 10% heat-inactivated fetal bovine serum (FBS; Gibco), 2 mM glutamine, 100 U/ml penicillin, and 100 µg/ml streptomycin (Gibco) for cell maintenance. BHK-21 and Vero76 cell culture was maintained in 37 °C incubator (ESCO) supplemented with 5% CO₂. C6/36 cell culture was maintained in 28 °C incubator. Virus stock titer was measured using modified standard plaque assay method (Fink et al., 2002) and recorded as plaque forming unit (PFU) equivalent/ml.

QIAamp Viral RNA Mini Kits (Qiagen, Hilden, Germany) were used to extract viral genomic RNAs from patients' sera or cell culture supernatant according to the manufacturer's instructions. Dengue viral RNA was reverse-transcribed into cDNA using Superscript III reverse transcriptase (RT) (Invitrogen, Carlsbad, CA) and DENV-specific primers (Lanciotti et al., 1992). Subsequently, cDNA was amplified using Taq DNA polymerase (Roche, Mannheim, Germany). The four dengue serotypes were distinguished by PCR product size.

2.3. Whole genome sequencing

Five overlapping PCR fragments covering the whole dengue genome were amplified using methods described previously (Ong et al., 2008; Schreiber et al., 2009) with slight modifications. Briefly, genomic RNAs were reverse-transcribed into cDNA using primers specific for each serotype. The primers used for DENV-4 sequencing were listed in Supplementary Table 1. Fragment amplification was performed using high-fidelity Pfu Turbo DNA polymerase (Stratagene, La Jolla, CA). Typical PCR conditions were as follows: templates were denatured for 2 min at 95 °C, followed by 45 cycles of 30 s denaturation at 95 °C, 1 min annealing at 55 °C, 4.5 min extension at 72 °C. Templates were allowed a 10 min final extension step at 72 °C followed by storage at 4 °C. PCR products were purified from 0.8% agarose gel using QIAquick gel extraction kit (Qiagen) and used as templates for cycle sequencing. DNA sequencing was performed using BigDye Dideoxy Terminator sequencing kits v3.1 (Applied Biosystems, Foster City, CA) using the method described in the kit. In addition to the sequencing and amplification primers previously described by Ong et al. (2008), a new set of primers for DENV-4 were used. Primer sequences and predicted genome binding positions of RT-PCR primers for DENV-4 are listed in Supplementary Tables 2 and 3,

191 respectively. ³ purified sequencing reactions were then run on
192 ABI3130xI (Applied Biosystems) sequencer at the Eijkman
193 Institute sequencing facility. Contig assembly was performed using
194 SeqSc⁴ 2.5 (Applied Biosystem) with additional manual adjust-
195 ment performed when manual inspection of ²⁵ assembly showed
196 some discrepancies. All genome sequences obtained in this study
197 have been deposited in GenBank (Table 1).

2.4. Sequence, phylogenetic and evolutionary analysis

198
199 We downloaded all DNA sequences for the DENV-1, -2, -3 and -
200 4 from NCBI GenBank database as of 13 April 2012. All sequences
201 from respected dengue serotype which had 90% identities with
202 any of the samples for 90% coverage of their sequences were
203 searched using USEARCH (Edgar, 2010). The obtained sequences

Table 1
Characteristics of the Makassar DENV samples with whole-genomes sequenced.

No	Sample ID	Accession no.	Sex	Isolation time	Genotype	Genotype	Infection status
1.	D1/ID/MKS-0055/2007	KC762649	F	5 2007	DENV-1	I	Primary
2.	D1/ID/MKS-0056/2007	KC762625	F	Jun 2007	DENV-1	I	Primary
3.	D1/ID/MKS-0077/2007	KC762654	F	Jul 2007	DENV-1	I	Primary
4.	D1/ID/MKS-0080/2007	KC762636	F	Jul 2007	DENV-1	I	Primary
5.	D1/ID/MKS-0088/2007	KC762635	M	Oct 2007	DENV-1	I	Primary
6.	D1/ID/MKS-0109/2007	KC762650	F	Jun 5 07	DENV-1	I	Primary
7.	D1/ID/MKS-0206/2007	KC762623	F	Jul 2007	DENV-1	I	Primary
8.	D1/ID/MKS-0248/2007	KC762632	M	Aug 2007	DENV-1	I	Primary
9.	D1/ID/MKS-0344/2007	KC762646	M	Aug 2007	DENV-1	I	Primary
10.	D1/ID/MKS-0352/2007	KC762637	F	Dec 2007	DENV-1	I	Primary
11.	D1/ID/MKS-0390/2007	KC762620	F	Dec 2007	DENV-1	IV	Primary
12.	D1/ID/MKS-0395/2008	KC762631	M	Feb 2008	DENV-1	I	Primary
13.	D1/ID/MKS-0397/2008	KC762621	F	Feb 2008	DENV-1	I	Primary
14.	D1/ID/MKS-0398/2008	KC762643	5	Feb 2008	DENV-1	I	Primary
15.	D1/ID/MKS-0483/2007	KC762651	M	Dec 2007	DENV-1	IV	Primary
16.	D1/ID/MKS-0486/2008	KC762653	M	Feb 2008	DENV-1	IV	Primary
17.	D1/ID/MKS-2029/2008	KC762640	M	Mar 2008	DENV-1	IV	Primary
18.	D1/ID/MKS-2040/2008	KC762645	F	Feb 2008	DENV-1	I	Primary
19.	D1/ID/MKS-2058/2008	KC762628	39	Feb 2008	DENV-1	IV	Primary
20.	D1/ID/MKS-2082/2008	KC762624	M	Feb 2008	DENV-1	I	Primary
21.	D1/ID/MKS-2094/2008	KC762644	M	Feb 2008	DENV-1	I	Primary
22.	D1/ID/MKS-2097/2008	KC762630	F	Feb 2008	DENV-1	IV	Primary
23.	D1/ID/MKS-2138/2008	KC762626	F	Feb 2008	DENV-1	I	Primary
24.	D1/ID/MKS-2147/2008	KC762641	F	Feb 2008	DENV-1	I	Primary
25.	D1/ID/MKS-2194/2008	KC762648	M	Apr 2008	DENV-1	I	Primary
26.	D1/ID/MKS-2200/2008	KC762622	M	Apr 2008	DENV-1	I	Primary
27.	D1/ID/MKS-2201/2008	KC762634	F	Apr 2008	DENV-1	I	Primary
28.	D1/ID/MKS-2216/2008	KC762633	M	Apr 2008	DENV-1	I	Primary
29.	D1/ID/MKS-1001/2008	KC762652	33	Apr 2008	DENV-1	I	Primary
30.	D1/ID/MKS-2004/2008	KC762638	F	Feb 2008	DENV-1	I	Primary
31.	D1/ID/MKS-IF039/2008	KC762629	F	Mar 2008	DENV-1	I	Primary
32.	D1/ID/MKS-IF062/2008	KC762627	F	Apr 2008	DENV-1	I	Secondary
33.	D1/ID/MKS-WS72/2010	KC762647	M	Mar 2010	DENV-1	50	Primary
34.	D1/ID/MKS-WS81/2010	KC762639	M	Apr 2010	DENV-1	I	Primary
35.	D1/ID/MKS-WS88/2010	KC762642	M	Jul 5 10	DENV-1	I	Primary
36.	D2/ID/MKS-0068/2007	KC762665	M	Jul 2007	DENV-2	Cosmopolitan	Primary
37.	D2/ID/MKS-0071/2007	KC762669	M	Jul 2007	DENV-2	Cosmopolitan	Primary
38.	D2/ID/MKS-0082/2007	KC762658	M	Aug 2007	DENV-2	Cosmopolitan	Secondary
39.	D2/ID/MKS-0084/2007	KC762660	M	Nov 2007	DENV-2	Cosmopolitan	Primary
40.	D2/ID/MKS-0091/2007	KC762670	F	Dec 2007	DENV-2	Cosmopolitan	Primary
41.	D2/ID/MKS-0099/2007	KC762676	M	Jul 2007	DENV-2	Cosmopolitan	Primary
42.	D2/ID/MKS-0297/2007	KC762662	M	Nov 2007	DENV-2	Cosmopolitan	Primary
43.	D2/ID/MKS-0412/2007	KC762661	F	Sep 2007	DENV-2	Cosmopolitan	Secondary
44.	D2/ID/MKS-0417/2007	KC762655	F	Sep 2007	DENV-2	Cosmopolitan	Primary
45.	D2/ID/MKS-0502/2007	KC762656	M	Dec 2007	DENV-2	Cosmopolitan	Primary
46.	D2/ID/MKS-2032/2008	KC762675	33	Mar 2008	DENV-2	Cosmopolitan	Primary
47.	D2/ID/MKS-2108/2008	KC762671	F	Feb 2008	DENV-2	Cosmopolitan	Primary
48.	D2/ID/MKS-2145/2008	KC762664	F	Mar 2008	DENV-2	Cosmopolitan	Primary
49.	D2/ID/MKS-2167/2008	KC762674	F	Mar 2008	DENV-2	Cosmopolitan	Primary
50.	D2/ID/MKS-2198/2008	KC762672	F	Apr 2008	DENV-2	Cosmopolitan	Primary
51.	D2/ID/MKS-2204/2008	KC762667	M	Apr 2008	DENV-2	Cosmopolitan	Primary
52.	D2/ID/MKS-2210/2008	KC762659	F	Apr 2008	DENV-2	Cosmopolitan	Secondary
53.	D2/ID/MKS-2234/2008	KC762657	M	Apr 2008	DENV-2	Cosmopolitan	Primary
54.	D2/ID/MKS-3007/2008	KC762666	39	Feb 2008	DENV-2	Cosmopolitan	Primary
55.	D2/ID/MKS-2024/2008	KC762663	M	Feb 2008	DENV-2	Cosmopolitan	Primary
56.	D2/ID/MKS-2018/2008	KC762673	M	Mar 2008	DENV-2	Cosmopolitan	Primary
57.	D2/ID/MKS-IF011/2008	KC762677	M	Feb 2008	DENV-2	Cosmopolitan	Primary
58.	D2/ID/MKS-IF014/2008	KC762668	F	Mar 2008	DENV-2	Cosmopolitan	Primary
59.	D2/ID/MKS-WS73/2010	KC762678	M	Mar 2010	DENV-2	Cosmopolitan	Primary
60.	D2/ID/MKS-WS79a/2010	KC762679	F	Mar 2010	DENV-2	Cosmopolitan	Primary
61.	D2/ID/MKS-WS80/2010	KC762680	M	Apr 2010	DENV-2	Cosmopolitan	Primary
62.	D3/ID/MKS-0057/2007	KC762681	F	Jun 2007	DENV-3	I	Primary
63.	D3/ID/MKS-0076/2007	KC762682	F	Jul 2007	DENV-3	I	Primary

(continued on next page)

Table 1 (continued)

No	Sample ID	Accession no.	Sex	Isolation time	Serotype	Genotype	Infection status
64.	D3/ID/MKS-0079/2007	KC762683	F	Jul 2007	DENV-3	I	Primary
65.	D3/ID/MKS-0098/2007	KC762684	M	Jul 2007	DENV-3	I	Secondary
66.	D3/ID/MKS-0172/2007	KC762686	M	Jul 2007	DENV-3	I	Secondary
67.	D3/ID/MKS-0388/2008	KC762685	F	Feb 2008	DENV-3	I	Primary
68.	D3/ID/MKS-0396/2008	KC762687	M	Feb 2008	DENV-3	I	Primary
69.	D3/ID/MKS-2065/2008	KC762689	F	Feb 2008	DENV-3	I	Primary
70.	D3/ID/MKS-3000/2008	KC762690	M	Aug 2008	DENV-3	I	Primary
71.	D3/ID/MKS-2006/2008	KC762688	M	Apr 2008	DENV-3	I	Primary
72.	D3/ID/MKS-IF058/2008	KC762691	M	Apr 2008	DENV-3	I	Primary
73.	D3/ID/MKS-WS78/2010	KC762692	F	Mar 2010	DENV-3	I	Secondary
74.	D3/ID/MKS-WS79b/2010	KC762693	F	Mar 2010	DENV-3	I	Primary
75.	D4/ID/MKS-0033/2007	KC762694	M	May 2007	DENV-4	II	Primary
76.	D4/ID/MKS-0070/2007	KC762695	M	Jul 2007	DENV-4	II	Primary
77.	D4/ID/MKS-0252/2008	KC762696	M	Aug 2007	DENV-4	II	Primary
78.	D4/ID/MKS-0706/2008	KC762697	-	Feb 2008	DENV-4	II	Primary
79.	D4/ID/MKS-2139/2008	KC762699	M	Mar 2008	DENV-4	II	Primary
80.	D4/ID/MKS-2007/2008	KC762698	M	Apr 2008	DENV-4	II	Primary

together with the samples were then aligned using MUSCLE (Edgar, 2004). The alignments were trimmed for their 5'-UTR and 3'-UTR leaving only the coding regions. A maximum likelihood phylogenetic tree using GTR substitution model and gamma distribution for varying rate for each serotype was constructed by FastTree (Price et al., 2010). The phylogenetic trees were used to collect 60 most-closely related published sequences from each of the samples as reference sequences by calculating the patristic distance as implemented in DendroPy (Sukumaran and Holder, 2010).

MUSCLE (Edgar, 2004) was utilized to create the multiple sequence alignment of each dengue virus serotype for the untrimmed sample and the collected reference sequences. We partitioned the alignments into 5-UTR, coding and 3-UTR regions accordingly. We determined the best model of evolution for each region by sampling across substitution model space in Bayesian MCMC analysis as implemented in MrBayes (Huelsenbeck and Ronquist, 2001). For all serotypes, we selected the GTR with invariable sites and 4 classes of gamma-distribution (GTR + I + Γ_4) with codon model and TrN model with invariable sites and 4 classes of gamma-distribution (TrN + I + Γ_4) for the coding region and 3'-UTR, respectively, as these were the models implemented in BEAST that were close enough to the models suggested by MrBayes. Bayesian Monte Carlo Markov Chain (MCMC) simulations with the selected evolution models as partitioned models and relaxed molecular clock were generated by BEAST (Drummond and Rambaut, 2007) and used to estimate the evolutionary parameters of each serotype. A Bayesian skyline plot was utilized as a coalescent prior as described in previous publications (Ong et al., 2008; Schreiber et al., 2009). For each type of dengue virus, we conducted a run of 100,000,000 replications with sampling of every 1000th, resulting in 100,000 chains. MCC (Maximum Clade Credibility) phylogenetics trees and evolutionary parameters were summarized with 25% burn-in. Tree visualization and sample labeling was performed using FigTree software (<http://tree.bio.ed.ac.uk/software/figtree>).

2.5. Viral growth kinetic assay

Vero76 cells were seeded at a density of 2×10^5 cells and allowed to attach overnight in 24 well plates (Nunc). Cells were then infected with viruses in duplicate using a multiplicity of infection of 0.1 in $1 \times$ MEM supplemented with 2% FBS and incubated at 37 °C, 5% CO₂ for 1 h. Three to five different virus isolates for each serotype were assayed. For comparison between Genotypes of DENV-1, four different isolates were used for each genotype. Inoculation media were aspirated and replenished with

fresh medium following incubation. Culture supernatants were collected at 12 h intervals (i.e., 12, 24, 36, 48, 60, and 72 h post inoculation) and stored directly in -80 °C freezer. Viral titers from each time point were quantified using a standard plaque assay on BHK-21 cell lines as described previously (Fink et al., 2007) and plotted as titration curves showing the mean titer and standard deviation. Statistical analysis was done using SPSS Statistics software version 17.0 (SPSS Inc., Chicago, IL). One-way analysis of variance (ANOVA) was used to analyze and compare means of viral titer among DENV serotypes. Comparison of viral growth kinetic between genotypes of DENV-1 was done using independent-samples Student's t-test. A probability value of $p < 0.05$ was considered statistically significant.

3. Results

3.1. Dengue incidence in Makassar city

Four hundred and fifty-five febrile patients were recruited during May 2007–August 2010. Of these, 197 (43.3%) patients were confirmed dengue as determined by NS1 rapid test kit. RT-PCR detections were performed on NS1-positive samples and a total of 126 sera (27.7% of all dengue-suspected patients) were detected as dengue positive. The male to female ratio was 1.14, with an average age of 12.5 ± 10.2 years old. Cases occurred predominantly in those aged between 11 and 20 years old (40%). Most dengue positive cases ($n = 165$ or 83.8%) were secondary infections, as determined by IgM and IgG ELISA ratio.

3.2. Dengue virus serotype distribution

We performed serotyping on RT-PCR-confirmed dengue cases using a standard RT-PCR serotyping method (Lanciotti et al., 1992) and obtained results in 126 cases, that demonstrated the circulation of all four dengue serotypes in Makassar. The predominant serotype was DENV-1 with 51 isolates (41%), followed by DENV-2 with 39 isolates (31%). Twenty-five isolates (20%) were DENV-3; and the remaining nine isolates (7%) were DENV-4. Two samples were mixed-infection of DENV-1 and -2 and DENV-2 and -3. We compared previous publications on serotype distributions in other parts on Indonesia (Corwin et al., 2001; Graham et al., 1999; Porter et al., 2005; Sukri et al., 2003; Suwandono et al., 2006; Yamanaka et al., 2011) with our results. We observed a distinct serotype distribution of DENV in Makassar compared to other cities throughout the Indonesian archipelago although we cannot be certain that

288 these historical reports reflect current serotype distributions in
289 those cities.

290 3.3. Phylogenetic analysis of dengue viruses

291 To determine the predominant genotypes in Makassar, we per-
292 formed full-length genome sequencing of the isolated viruses.
293 Virus genomic RNAs were extracted either from patient sera or
294 after one passage of virus propagation in tissue culture. We
295 obtained the complete genome sequences of 80 viruses covering
296 all four serotypes. Table 1 describes the characteristics of the
297 sequenced isolates including the length of the sequenced genomes.

298 Using a fast maximum-likelihood method as implemented in
299 FastTree, we performed an approximate phylogenetic analysis of
300 all dengue serotypes to select 60 most-closely related sequences
301 of each dengue sample for reference sequences for consecutive
302 analysis.

303 The MCC (Maximum Clade Credibility) phylogenetic tree of
304 DENV-1 isolates and their closest reference sequences was shown
305 in Fig. 1. Based on the DENV-1 genotype classification by
306 Goncalvez et al. (2002), we observed the circulation of two gen-
307 otypes of DENV-1 in Makassar. The majority of the viruses, including
308 all recent 2010 samples (29 isolates) belong to Genotype I. The 28
309 isolates of Genotype I were from single ancestor and had the time
310 to the most recent common ancestor (TMRCA) around 2005. One of
311 the recent samples (WS88) probably followed an independent
312 route of introduction as it had similar lineage with samples from
313 Singapore. All of Genotype I samples were clustered together with
314 viruses from China, Thailand, and Singapore, including the
315 Singapore dengue outbreak isolates of 2005. The rest of the
316 Makassar's DENV-1 viruses (6 isolates) were grouped into
317 Genotype IV. Three of the isolates were clustered together with
318 samples from Brunei (in 2006), Hawaii (in 2001), Philippines (in
319 2004) and China (in 1991), and their ancestor was likely to be
320 the same as the West Pacific sample of 1974 isolate. The other
321 three isolates were clustered with China isolates of 2002/2003
322 and Indonesian city of Palembang in 1998. It should be noted that
323 DENV-1 isolates from other Indonesian region were not included in
324 the analysis as they are more likely to distantly related to Makassar
325 samples and were not selected during the FastTree/patristic dis-
326 tance filtering process. This observation contributes to new infor-
327 mation on the DENV-1 genotype distribution in Indonesia.

328 We determined 26 DENV-2 genome sequences from Makassar
329 isolates and constructed an MCC phylogenetic analysis of those
330 viruses together with previously published DENV-2 genomes.
331 Based on the DENV-2 genotype classification reported by Twiddy
332 et al. (2002), all of our Makassar isolates were grouped into the
333 Cosmopolitan genotype (Fig. 2). This genotype is widely disseminated
334 from India to South East Asia, Africa, the Middle East, and
335 Australia. Within the Cosmopolitan genotype, the Makassar iso-
336 lates grouped together with two isolates from Jakarta in 2004
337 and many isolates from Singapore (in 2004, 2005 and 2008). The
338 phylogenetic trees suggested the presence of spatial isolation
339 among DENV-2 strains from cities in the western and eastern parts
340 of Indonesia, but the absence of spatial isolation between Makassar
341 and Singapore. The tree also indicated that DENV-2 viruses
342 currently circulating in Makassar were of multiple entries.

343 Phylogenetic analysis of 13 DENV-3 genome sequences from
344 Makassar identified them as belonging to Genotype I, based on
345 classification described by Lanciotti et al. (1994). This genotype
346 consists of viruses from Indonesia, Malaysia, the Philippines and
347 the South Pacific islands (Lanciotti et al., 1994). Indonesian viruses
348 collected in 1973, 1978 and 1985 also grouped into this genotype
349 (Lanciotti et al., 1994), as did recent isolates from Jakarta in 2004
350 and Palembang in 1998 (Fig. 3) and (Ong et al., 2008). Nine
351 Makassar isolates were clustered with Indonesian isolates of

1998 and 1988 from Jakarta and were from single common ances-
tor with TMRCA of 1990. Four other Makassar isolates, including
the recent 2010 samples, were clustered with the rest of
Indonesian isolates from 1998 and 2004, as well as from Timor
Leste in 2005 and 2005/2009 from Singapore. This may indicate
that all the Makassar samples were originated from and spread
in Indonesia and neighboring countries such as Singapore and
Timor Leste. TMRCA of all Makassar isolates together with other
Indonesian isolates was 1965. Together this suggests the presence
of strong temporal structure and that the Makassar's DENV-3
viruses appeared to be part of an older lineage which was indeed
endemic in the Indonesian region and had been circulating for
more than four decades.

We performed phylogenetic analysis on the genomes of 6
DENV-4 viruses isolated from Makassar and compared them with
available DENV-4 genome in the GenBank. As shown in Fig. 4,
the Makassar isolates fell into Genotype II based on Lanciotti clas-
sification (Lanciotti et al., 1997). This genotype is commonly found
in South-east Asia and America (Lanciotti et al., 1997). The
Makassar isolates grouped closely with other Indonesian isolates
from Jakarta that were isolated in 2004. Four of the Makassar sam-
ples in the same lineage with the Jakarta samples were originated
from a lineage which has higher mutation rate relative to other
samples. The TMRCA of these 4 Makassar samples and the related
Jakarta samples was around year 1990. This clustering suggests the
endemicity of this DENV-4 genotype in Indonesia, and the presence
of strong temporal link among DENV-4 viruses in Indonesia.

349 3.4. Evolutionary dynamics and amino acid comparison of Makassar
350 dengue viruses

351 The estimated evolutionary rates for substitution dynamics of
all serotypes, as calculated by BEAST, were 1.1×10^{-3} substitution/
site/year (95% highest probability density (HPD), 0.6×10^{-3} –
 1.8×10^{-3} substitution/site/year) and 1.5×10^{-3} substitution/
site/year (95% HPD, 0.7×10^{-3} – 1.7×10^{-3} substitution/site/year)
for DENV-1 and DENV-2, respectively. Similar estimates,
 1.1×10^{-3} substitution/site/year (95% HPD, 6.2×10^{-4} – $1.5 \times$
 10^{-3} substitution/site/year) and 0.2×10^{-3} substitution/site/year
(95% HPD, 1.5×10^{-4} – 0.8×10^{-3} substitution/site/year), were
made for DENV-3 and DENV-4, respectively.

352 The estimated ages of the most recent common ancestors
(MRCA) for all serotypes were 62 months (95% HPD, 53–91
months), 58 months (95% HPD, 51–86 months), 56 months (95%
HPD, 49–82 months) and 55 months (95% HPD, 45–78 months)
for DENV-1, DENV-2, DENV-3 and DENV-4, respectively.

353 A relatively low ratio of non-synonymous to synonymous sub-
stitution per site (dN/dS) in all genes indicated that there was no
evidence for positive selection in any gene and no evidence for
site-specific positive selection of the Makassar viruses (Table 2).
Several genes had ratio of 1.0 for dN/dS in DENV-4, however due
to small number of DENV-4 isolates, we feel that the dN/dS for
DENV-4 cannot be determined confidently.

354 Comparative analysis at the amino acid levels were illustrated
in Fig. 5 for DENV-1 Genotype I (A), DENV-1 Genotype IV (B),
DENV-2 (C), DENV-3 (D) and DENV-4 (E). Although DENV-1
Genotype I had the most sample number, the differences were
the smallest among the sample set, and substantially much smaller
than the DENV-1 genotype IV. In DENV-1 Genotype I residue 123 of
E gene was observed to be either Serine (S) or Threonine (T) in many
2007 samples, but only observed to be T in all 2008 samples.
Likewise, residue 222 of NS1 gene in some 2007 samples were
observed to be S or T, while the rest of 2008 samples
were observed to be T. However, residue 829 of NS5 gene which
were observed to be Isoleucine (I) for all 2007 samples, started to
have Valine (V) in some of 2008 samples. The rest of changes in




Fig. 1. MCC (Maximum Clade Credibility) phylogenetic tree of DENV-1 Genotype I and Genotype IV. The tree consisted of 35 complete genomes of DENV-1 isolated from Makassar, Indonesia together with their close-distanced genomes available in the GenBank database and was constructed using Bayesian inference as implemented in BEAST. All Makassar isolates were denoted with red labels, while samples from Indonesian cities of Palembang and Jakarta were denoted with green labels. Posterior probabilities of all major branches had values of more than 0.75, indicating the significant of the tree topology. The red line for a branch indicated that the rate of mutation of that particular branch was relatively faster compared to the overall rate of mutation, while the blue line indicated slower rate of mutation. The MCC tree branched into 2 main branches, the genotype IV (upper branch) and I (lower branch). Six of the Makassar isolates grouped into genotype IV, while the rest of Makassar isolates grouped into genotype I. Labels indicating the origin of isolates are as follow: BN (Brunei), CL (Chile), CN (China), ID (Indonesia), JP (Japan), MY (Malaysia), NR (Nauru Island), PF (Polynesian French), PH (Philippines), RE (Reunion Island), SG (Singapore), SC (Seychelles Island), TH (Thailand), and US (United States). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. MCC phylogenetic tree of DENV-2 genotype Cosmopolitan. The tree consisted of 26 complete genomes of DENV-2 viruses isolated from Makassar, Indonesia, together with their close-distanced genomes available in the GenBank database and was constructed using Bayesian inference as implemented in BEAST. All Makassar isolates were denoted with red labels, while samples from Indonesian cities of Palembang and Jakarta were denoted with green labels. Posterior probabilities of all major branches had values of more than 0.75, indicating the significant of the tree topology. The red line for a branch indicated that the rate of mutation of that particular branch was relatively faster compared to the overall rate of mutation, while the blue line indicated slower rate of mutation. Labels indicating the origin of isolates are as follow: AU (Australia), BN (Brunei), CN (China), GU (Guam), ID (Indonesia), SG (Singapore), TW (Taiwan) and VN (Vietnam). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. MCC phylogenetic tree of DENV-3 Genotype I. The tree consisted of 13 complete genomes of DENV-3 viruses isolated from Makassar, Indonesia, together with their close-distanced genomes available in the GenBank database and was constructed using Bayesian inference as implemented in BEAST. All Makassar isolates were denoted with red labels, while samples from other Indonesian regions were denoted with green labels. Magenta labels indicated Indonesian wild-type samples that have undergone serial passages in Vero cells. Posterior probabilities of all major branches had values of more than 0.75, indicating the significant of the tree topology. The red line for a branch indicated that the rate of mutation of that particular branch was relatively faster compared to the overall rate of mutation, while the blue line indicated slower rate of mutation. Labels indicating the origin of isolates are as follow: BR (Brazil), CK (Cook Islands), CN (China), ID (Indonesia), LK (Sri Lanka), MZ (Mozambique), PF (Polynesian French), PH (Philippines), SG (Singapore), TH (Thailand), TL (East Timor), TW (Taiwan), and WS (Western Samoa). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

416 the amino acids of the Makassar samples were unlikely to be
417 significant.
418 In DENV-1 Genotype IV, the most diverse samples, we
419 observed more significant differences in the amino acid changes.
420 Some sites, notably residue 217 of E, 115 of NS2B, 119 of NS3,
421 426 of NS3, 123 of NS4A genes, contained amino acid Proline
422 (P) which structurally different than that of other amino acids
423 as it had cyclic group in its backbone chain, which might bend
424 the backbone slightly (Fig. 5B).

In DENV-2, similar differences also occurred in the amino acid
changes. Residue 217, 332 and 384 of E, 320 of NS1, and 837 of
NS5 contained a mix of amino acid Proline (P) and Leucine (L).
Changes that were likely to be significant occurred in M (residue
57 and 71), E (residue 346), and NS5 (residue 563 and 570) as they
involved changes of amino acids with different physicochemical
properties (Fig. 5C).

Both DENV-3 and DENV-4 comparative analysis showed that
most differences came from 1 or 2 amino acid changes in most of




Fig. 4. MCC phylogenetic tree of DENV-4 Genotype II. The tree consisted of 6 complete genomes of DENV-4 viruses isolated from Makassar, Indonesia, together with their close-distanced genomes available in the GenBank database and was constructed using Bayesian inference as implemented in BEAST. All Makassar isolates were denoted with red labels, while samples from other Indonesian regions were denoted with green labels. Posterior probabilities of all major branches had values of more than 0.75, indicating the significant of the tree topology. The red line for a branch indicated that the rate of mutation of that particular branch was relatively faster compared to the overall rate of mutation, while the blue line indicated slower rate of mutation. Labels indicating the origin of isolates are as follow: BR (Brazil), CO (Colombia), CN (China), DM (Dominica), HT (Haiti), PF (Polynesian French), PH (Philippines), PR (Puerto Rico), SG (Singapore), TH (Thailand), TW (Taiwan), US (United States), and VE (Venezuela). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Table 2
Evolutionary patterns among the proteins of DENV-1, -2, -3 and -4 sampled from Makassar, Indonesia during 2007–2010.

Protein	Length (bp)	DENV-1		DENV-2		DENV-3		DENV-4	
		IS	dN/dS	IS	dN/dS	IS	dN/dS	IS	dN/dS
Capsid (C)	342	24	0.25	12	0.17	1	0	1	1.0
Membrane (M)	498	52	0.13	19	0.11	2	0.5	0	0
Envelope (E)	1479	167	0.11	53	0.10	11	0.1	0	0
NS1	1056	102	0.18	40	0.17	5	0.2	1	1.0
NS2A	654	75	0.19	27	0.19	7	0.29	3	0
NS2B	390	36	0.19	13	0.08	3	0	3	0.67
NS3	1857	187	0.11	67	0.03	15	0.2	2	1.0
NS4A	450	49	0.10	14	0	0	0	0	0
2K	68	8	0	4	0	0	0	1	0
NS4B	744	73	0.04	27	0.07	4	0	2	0.5
NS5	2679	230	0.10	103	0.18	10	0	8	0
Genome	10170	1003	NA	379	NA	58	NA	21	NA

* Coding region only; IS, number of informative sites; NA, not applicable.


434 the sites. Unfortunately, the sample sizes of both DENV-3 and
435 DENV-4 are too small to confidently detect any significant changes
436 or evidence of positive selection within these clades (Fig. 5D and
437 E).

3.5. Growth characteristics of Makassar DENV

Our study observed the circulation of all four DENV serotypes in Makassar. Phylogenetic analyses revealed the genotypes of viruses

10

R.T. Sasmono et al. / Infection, Genetics and Evolution xxx (2015) xxx–xxx



36

28

Fig. 5. Comparative analyses of amino acid substitutions among Makassar's DENV-1 viruses genotype I (A) and IV (B), DENV-2 viruses (C), DENV-3 viruses (D), and DENV-4 (E). Genes in which the substitutions present are shown above the amino-acid numbers. Amino acid coloring scheme is according to AESNN3 (Lin et al., 2002). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

441 isolated in this study and grouped the Makassar virus isolates into
442 several clades within each genotype (Figs. 1–4). It has been sug-
443 gested that particular genotypes of DENV possess different rates
444 of transmission and that some genotypes have a tendency to cause

more severe forms of dengue disease and are thus more virulent
than others (Rico-Hesse, 2010). One way to estimate the virulence
is by assessing the replication kinetics of the virus (Rico-Hesse,
2010).

445
446
447
448




Fig. 6. Replication kinetic assays of selected DENV from Makassar. (A) Replication kinetic comparison of four DENV serotypes. (B) Replication kinetic comparison of DENV-1 Genotype I and Genotype IV strains. Data were the mean \pm SD of three to five virus isolates representing each serotype/genotype performed in duplicate.

13

449 To determine the growth characteristics of Makassar's DENV,
450 we measured the replication kinetics of all four serotypes and
451 genotypes of DENV grown in mammalian Vero76 cell line. Three
452 to four viruses representing each serotype/genotype were assayed
453 for their replication kinetics. Growth kinetic comparison of DENV-
454 1, -2, -3, and -4 is shown in Fig. 6. We observed serotype-specific
455 differences in replication at 36 and 48 h post-infection, with
456 DENV-1 and -2 replicating to higher levels compared with DENV-
457 3 and -4 (Fig. 6A).

458 Next, we compared the replication kinetics of viruses belong to
459 two different genotypes of DENV-1, i.e., Genotype I and IV (Fig. 6B).
460 Genotype I apparently replicated at higher level compared to
461 Genotype IV (p value < 0.05). This suggests the relatively better fit-
462 ness of Genotype I which may underlie the ability of this Genotype
463 to replace the Genotype IV in the community. Similar kinetic pat-
464 terns were observed when other cell line (C6/36) was used in
465 growth kinetic experiment (data not shown). Replication kinetic
466 assays on four viruses representing four clades of DENV-2 did
467 not show any significant difference (data not shown).

468 4. Discussion

469 We conducted the first molecular-typing study of dengue in
470 Makassar-Indonesia to investigate dengue disease dynamics in
471 the area from 2007 to 2010. As confirmed by NS1 antigen

472 detection, we observed dengue viruses infected a remarkable
473 43.3% of febrile patients presenting to hospital. This data reflects
474 the importance of the disease in Makassar and the considerable
475 burden it must place on the local economy and health infrastruc-
476 ture. In terms of infection status, based on IgM and IgG ELISA
477 detection, our data showed a high number of secondary dengue
478 infections in hospitalized patients who participated in this study
479 (83.8%), indicating a sustained disease intensity over a number of
480 years. The age distribution of the hospitalized patients was found
481 to be highest in young patients (10–20 years old), similar to other
482 parts of Indonesia (Corwin et al., 2001). It is likely that this pre-
483 dominant age group may correlate with the higher proportion of
484 secondary infections observed.

485 To date, no serotype distribution data has been available for
486 Makassar. Our study demonstrated the circulation of all four den-
487 gue serotypes. We observed the predominance of DENV-1 (41%)
488 followed by DENV-2, -3, -4. This distribution is distinct from his-
489 torical reports in other parts of Indonesia, especially in the
490 Western regions of the archipelago, e.g., in the cities of Jakarta
491 and Palembang in 2004, in which DENV-3 was the predominant
492 serotype in the area and correlated with severe symptoms. It
493 would be interesting to know if the current serotype distributions
494 in Jakarta and Palembang match their historical values or if they
495 are now closer to what we have found in Makassar.

496 To study the evolutionary dynamics and phylogeny of
497 Makassar's dengue viruses, we performed whole-genome

sequencing of 80 virus isolates. Most of the genomes were sequenced from primary infection samples due to difficulties in virus isolation from secondary infection samples, as previously reported (Jarman et al., 2011). Based on the published classification of dengue virus's genotypes, we identified the co-circulation of Genotypes I and IV of the DENV-1 serotype. The nearest geographical areas that have been reported as harboring Genotype I are Singapore (Schreiber et al., 2009), Thailand (Ong et al., 2008) and Surabaya (Yamanaka et al., 2011). The clustering of all Genotype I samples together with viruses from China, Thailand, and Singapore may suggest that the Genotype I isolates were originated from China, migrated to Thailand and passed Singapore before entering Makassar (Fig. 1). Genotype IV was predominant in Jakarta in 2004 (Schreiber et al., 2009) and was also present in Makassar in smaller numbers. The close genetic relationship between the Makassar and Singapore Genotype I isolates suggests a recent importation has occurred and that the incumbent clade is being replaced. While the Genotype IV strains from Makassar are quite diverse the Genotype I strains appear to be relatively uniform further suggesting a recent introduction. We will continue to survey viruses from the region to see if Genotype I succeed in displacing Genotype I.

Makassar's DENV-2 viruses grouped into the Cosmopolitan genotype. This genotype circulates widely in India, South East Asia, Africa, the Middle East, and Australia (Twiddy et al., 2002). This genotype was also circulating in the cities of Jakarta in 2004 and Palembang in 1998. Phylogenetic analysis showed that closely related Cosmopolitan genotype strains were isolated in Jakarta in 2004, Brunei and Singapore in 2005, Vietnam in 2006 and Makassar in 2007 and 2008. It appears that the Cosmopolitan genotype is very successful in spreading, regionally endemic and continually re-distributing in South East Asia. The phylogenetic trees suggested the presence of spatial isolation among DENV-2 strains from cities in the western and eastern parts of Indonesia, but the absence of spatial isolation between Makassar and Singapore (Fig. 2). The tree also indicated that DENV-2 viruses currently circulating in Makassar were of multiple entries. Presumably, frequent direct air and sea travel between Makassar and Singapore is rapidly distributing the viruses between the two cities. Another explanation of the possible spatial isolation among Indonesian cities is that there is no other whole genome sequence data that has been reported and submitted to GenBank.

Our phylogenetic analysis revealed that Makassar's DENV-3 viruses grouped into Genotype I, similar to isolates from Jakarta in 2004 and Palembang in 1998, and same genotype as Indonesian strains from 1978, 1988, 1996 and 1998 isolates. This genotype has been described as causing four epidemics in the region in the past (Ong et al., 2008), and is most likely a local, Indonesian endemic strain that has circulated for more than three decades. It is interesting that the isolates we found in Makassar appear to be more closely related to strains from 1996 and 1998 than the strains more recently found in Timor-Leste, Jakarta, Singapore and Palembang suggesting DENV-3 in Makassar has been following a separate evolutionary history since 1988.

The DENV-4 viruses we isolated clustered into Genotype II (SE Asia and America), the same genotype as those isolates in Jakarta in 2004. Four of the Makassar samples were originated from a lineage with higher mutation rates relative to other samples, and were direct descendant from the ancestor of those Jakarta samples with TMRCA around year 1990. Higher mutation rates might indicate that this particular lineage was spreading rather quickly, as in outbreak cases. As such, those particular Makassar isolates were assumed to be descendant of Jakarta outbreak in 1998. Partial sequences of viruses from 1973, 1976, 1977 isolates also cluster into this clade (not shown). This genotype appears to be endemic in Indonesia and has been circulating for over 30 years.

A Bayesian coalescent approach was used to infer the evolutionary dynamics for all serotypes of isolates within 2007–2010 in Makassar (Diamond and Rambaut, 2007). We observed the mean of evolution rates for DENV-1, DENV-2 and DENV-3 were similar to those estimated previously for all DENV serotypes (Twiddy et al., 2003). The mean age of the common ancestor for DENV-1, DENV-2 and DENV-3 indicated that each of these diverse lineages had common ancestors already present in Makassar since approximately 2003. Since the number of samples for DENV-4 was small, we could not confidently estimate the mean age of the common ancestor of these isolates. That we did not see any convincing evidence for positive selection in any of the serotypes indicates that these strains are already well adapted to their environment.

The analysis based on the amino acid sequences of the Indonesian samples showed that some positions were undergoing amino acid substitutions from 2007 to 2008. That we did not see any convincing evidence for positive selection in any of the serotypes indicates that these strains are already well adapted to their environment, and that the amino acid substitutions might not have affected the characteristics of viruses substantially.

Genetic variations in dengue viruses have been described as being risk factors for severe disease (Balmaseda et al., 2006; Messer et al., 2003; Rico-Hesse et al., 1997) and particular serotypes/genotypes are suggested to have a tendency to cause more severe forms of dengue (i.e., DHF and ILL) and larger outbreaks. For example, DENV-2 and DENV-3 have been associated with severe disease and DENV-4 is claimed to lead to a milder illness (Nisalak et al., 2003; Vaughn et al., 2000). One of the ways to estimate the virulence or fitness of a virus *in vitro* is by looking at the virus replication rate (Rico-Hesse, 2010). In this study, we compared the replication kinetics of various serotype/genotypes of DENV to determine whether particular serotype/genotypes replicate better than others do.

The co-circulation of two DENV-1 genotypes (i.e., Genotype I and IV) prompted us to compare their replication rates. Isolates grouped into Genotype I apparently replicated at higher level compared to Genotype IV. This suggests the relatively better fitness of Genotype I which may underlie the ability of this genotype to replace Genotype IV in the community. Thus, this observation suggests that different genotypes of DENV may exhibit different phenotypic characteristics, as in the case of American and South East Asian genotypes of DENV-2 (Leitmeyer et al., 1999; Rico-Hesse et al., 1997). Continuous surveillance is currently underway to confirm the predominant circulation of Genotype I.

The DENV-2 American and South East Asian genotypes are known to have differences in their replication ability. American genotypes replicated less well than the South East Asian viruses in *in vitro* experiments (Pryor et al., 2001). As all of our DENV-2 viruses were grouped into Cosmopolitan genotype (Fig. 3), we could not perform replication comparisons against other genotypes. However, within the Cosmopolitan genotype, we observed the grouping of viruses into clades. To determine whether there are differences in growth, four viruses representing each clade were assayed for their replication kinetics. We observed no significant differences among those four viruses (data not shown), suggesting that members of the genotype are quite homogenous in their growth rates. Similar observation is also present in DENV-3 and -4 (data not shown). A typical DENV growth kinetic profile was also observed in other virus isolates from other cities in Indonesia (data not shown).

In summary, we have revealed the dengue serotype and genotype distribution in Makassar and observed that although these viruses have very similar evolutionary dynamics they have remarkably different phylogeographic histories. We observed the different replication rates of DENV-1 Genotype I and Genotype

630 IV, which may underlie the ability of Genotype I in replacing the
631 Genotype IV in Makassar, Indonesia.

632 **Acknowledgements**

633 This study was funded by Novartis Institute for Tropical
634 Diseases (NITD), Singapore, through research collaboration
635 between NITD, Eijkman Institute and Hasanuddin University
636 (NEHCRI). NEHCRI laboratories were located in Eijkman Institute
637 for Molecular Biology, Jakarta and in Hasanuddin University,
638 Makassar, Indonesia. We would like to thank Profs. A. Matter
639 and S. Marzuki for their support in the establishment of NEHCRI,
640 as well as to Dr. J. Leong for his management and ethics advice.
641 We thank all clinicians in Makassar hospitals and health centres
642 that provided patients access for the study. Drs. S. Vasudevan
643 and M. Hibberd were involved in the initial stages of the study.
644 We are grateful for the helps of Drs. S. Ong, Q. Wang, L. Wei, and
645 A. Taslim and technical assistance from Ms. F. Meutiawati and
646 R.F. Hayati. The services of the Eijkman Institute's sequencing facil-
647 ity were greatly appreciated.

648 **Appendix A. Supplementary data**

649 Supplementary data associated with this article can be found, in
650 the online version, at <http://dx.doi.org/10.1016/j.meegid.2015.03.006>.
651

652 **References**

653 Balmaseda, A., Hammond, S.N., Perez, L., Tellez, Y., Saborio, S.J., Mercado, J.C.,
654 Cuadra, R., Rocha, J., Perez, M.A., Silva, S., Rocha, C., Harris, E., 2006. Serotype-
655 specific differences in clinical manifestations of dengue. *Am. J. Trop. Med. Hyg.*
656 *74*, 449–456.
657 Corwin, A.L., Larasati, R.P., Bangs, M.J., Wuryadi, S., Arjoso, S., Sukri, N., Listryaningsih,
658 E., Hartati, S., Namursa, R., Anwar, Z., Chandra, S., Loho, B., Ahmad, H., Campbell,
659 J.R., Porter, K.R., 2001. Epidemic dengue transmission in southern Sumatra,
660 Indonesia. *Trans. R. Soc. Trop. Med. Hyg.* *95*, 257–265.
661 Drummond, A.J., Rambaut, A., 2007. BEAST: bayesian evolutionary analysis by
662 sampling trees. *BMC Evol. Biol.* *7*, 214.
663 Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and
664 high throughput. *Nucleic Acids Res.* *32*, 1792–1797.
665 Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST.
666 *Bioinformatics* *26*, 2460–2461. <http://dx.doi.org/10.1093/bioinformatics/btq461>.
667 Fink, J., Gu, F., Ling, L., Tolfvenstam, T., Olfat, F., Chin, K.C., Aw, P., George, J.,
668 Kuznetsov, V.A., Schreiber, M., Vasudevan, S.G., Hibberd, M.L., 2007. Host gene
669 expression profiling of dengue virus infection in cell lines and patients. *PLoS*
670 *Negl. Trop. Dis.* *1*, e86.
671 Goncalvez, A.P., Escalante, A.A., Pujol, F.H., Ludert, J.E., Tovar, D., Salas, R.A., Liprandi,
672 F., 2002. Diversity and evolution of the envelope gene of dengue virus type 1.
673 *Virology* *303*, 110–119.
674 Graham, R.R., Juffrie, M., Tan, R., Hayes, C.G., Laksono, I., Ma'roef, C., Erlin, Sutaryo.,
675 Porter, K.R., Halstead, S.B., 1999. A prospective seroepidemiologic study on
676 dengue in children four to nine years of age in Yogyakarta, Indonesia I. *Studies*
677 *in 1995–1996. Am. J. Trop. Med. Hyg.* *61*, 412–419.
678 Gubler, D.J., 1998. Dengue and dengue hemorrhagic fever. *Clin. Microbiol. Rev.* *11*,
679 480–496.
680 Guzman, M.G., Halstead, S.B., Artsob, H., Buchy, P., Farrar, J., Gubler, D.J.,
681 Hunsperger, E., Kroeger, A., Margolis, H.S., Martínez, E., Nathan, M.B.,
682 Pelegrino, J.L., Simmons, C., Yoksan, S., Peeling, R.W., 2010. Dengue: a
683 continuing global threat. *Nat. Rev. Microbiol.* *8*, 7–16.
684 Harris, E., Roberts, T.G., Smith, L., Selle, J., Kramer, L.D., Valle, S., Sandoval, E.,
685 Balmaseda, A., 1998. Typing of dengue viruses in clinical specimens and
686 mosquitoes by single-tube multiplex reverse transcriptase PCR. *J. Clin.*
687 *Microbiol.* *36*, 2634–2639.
688 Holmes, E.C., Burch, S.S., 2000. The causes and consequences of genetic variation in
689 dengue virus. *Trends Microbiol.* *8*, 74–77.
690 Holmes, E.C., Twiddy, S.S., 2003. The origin, emergence and evolutionary genetics of
691 dengue virus. *Infect Genet Evol.* *3*, 19–28.
692 Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: bayesian inference of phylogenetic
693 trees. *Bioinformatics* *17*, 754–755.
694 Jarman, R.G., Nisalak, A., Anderson, K.B., Klungthong, C., Thaisomboonsuk, B.,
695 Kaneechit, W., Kalayanarooj, S., Gibbons, R.V., 2011. Factors influencing dengue
696 virus isolation by C6/36 cell culture and mosquito inoculation of nested PCR-
697 positive clinical samples. *Am. J. Trop. Med. Hyg.* *84*, 218–223. <http://dx.doi.org/10.4269/ajtmh.2011.09-0798>.

Lanciotti, R.S., Calisher, C.H., Gubler, D.J., Chang, G.J., Vorndam, A.V., 1992. Rapid
700 detection and typing of dengue viruses from clinical samples by using reverse
701 transcriptase-polymerase chain reaction. *J. Clin. Microbiol.* *30*, 545–551.
702 Lanciotti, R.S., Gubler, D.J., Trent, D.W., 1997. Molecular evolution and phylogeny of
703 dengue-4 viruses. *J. Gen. Virol.* *78* (Pt 9), 2279–2284.
704 Lanciotti, R.S., Lewis, J.G., Gubler, D.J., Trent, D.W., 1994. Molecular evolution and
705 epidemiology of dengue-3 viruses. *J. Gen. Virol.* *75* (Pt 1), 65–75.
706 Leitmeyer, K.C., Vaughn, D.W., Watts, D.M., Salas, R., Villalobos, I., de, C., Ramos, C.,
707 Rico-Hesse, R., 1999. Dengue virus structural differences that correlate with
708 pathogenesis. *J. Virol.* *73*, 4738–4747.
709 Lin, K., May, A.C., Taylor, W.R., 2002. Amino acid encoding schemes from protein
710 structure alignments: multi-dimensional vectors to describe residue types. *J.*
711 *Theor. Biol.* *216*, 361–365.
712 Messer, W.B., Gubler, D.J., Harris, E., Sivananthan, K., de Silva, A.M., 2003. Emergence
713 and global spread of a dengue serotype 3, subtype III virus. *Emerg. Infect. Dis.* *9*,
714 800–809.
715 Nisalak, A., Endy, T.P., Nimmannitya, S., Kalayanarooj, S., Thisayakorn, U., Scott, R.M.,
716 Burke, D.S., Hoke, C.H., Innis, B.L., Vaughn, D.W., 2003. Serotype-specific dengue
717 virus circulation and dengue disease in Bangkok, Thailand from 1973 to 1999.
718 *Am. J. Trop. Med. Hyg.* *68*, 191–202.
719 Ong, S.H., Yip, J.T., Chen, Y.L., Liu, W., Harun, S., Lystiyanningsih, E., Heriyanto, B.,
720 Beckett, C.G., Mitchell, W.P., Hibberd, M.L., Suwandono, A., Vasudevan, S.G.,
721 Schreiber, M.J., 2008. Periodic re-emergence of endemic strains with strong
722 epidemic potential—a proposed explanation for the 2004 Indonesian dengue
723 epidemic. *Infect. Genet. Evol.* *8*, 191–204.
724 Pandey, B.D., Igarashi, A., 2000. Severity-related molecular differences among
725 nineteen strains of dengue type 2 viruses. *Microbiol. Immunol.* *44*, 179–188.
726 Porter, K.R., Beckett, C.G., Kosasih, H., Tan, R.I., Alisjahbana, B., Rudiman, P.I.,
727 Widjaja, S., Listryaningsih, E., Ma'Roef, C.N., McArdle, J.L., Parwati, I., Sudjana, P.,
728 Jusuf, H., Yuwono, D., Wuryadi, S., 2005. Epidemiology of dengue and dengue
729 hemorrhagic fever in a cohort of adults living in Bandung, West Java, Indonesia.
730 *Am J Trop Med Hyg* *72*, 60–66.
731 Price, M.N., Dehal, P.S., Arkin, A.P., 2010. FastTree 2—approximately maximum-
732 likelihood trees for large alignments. *PLoS One* *5*, e9490. <http://dx.doi.org/10.1371/journal.pone.0009490>.
733 Pryor, M.J., Carr, J.M., Hocking, H., Davidson, A.D., Li, P., Wright, P.J., 2001.
734 Replication of dengue virus type 2 in human monocyte-derived macrophages:
735 comparisons of isolates and recombinant viruses with substitutions at amino
736 acid 390 in the envelope glycoprotein. *Am. J. Trop. Med. Hyg.* *65*, 427–434.
737 Raekiansyah, M., Pramesyanti, A., Bela, B., Kosasih, H., Ma'roef, C.N., Tobing, S.Y.,
738 Rudiman, P.I., Alisjahbana, B., Endi, T.P., Green, S., Kalayanarooj, S., Rothman,
739 A.L., Sudiro, T.M., 2005. Genetic variations and relationship among dengue virus
740 type 3 strains isolated from patients with mild or severe form of dengue disease
741 in Indonesia and Thailand. *Southeast Asian J. Trop. Med. Public Health* *36*,
742 1187–1197.
743 Rico-Hesse, R., 2010. Dengue virus virulence and transmission determinants. *Curr.*
744 *Trop. Microbiol. Immunol.* *338*, 45–55.
745 Rico-Hesse, R., Harrison, L.M., Salas, R.A., Tovar, D., Nisalak, A., Ramos, C., Boshell, J.,
746 de Mesa, M.T., Nogueira, R.M., da Rosa, A.T., 1997. Origins of dengue type 2
747 viruses associated with increased pathogenicity in the Americas. *Virology* *230*,
748 244–251.
749 Schreiber, M.J., Holmes, E.C., Ong, S.H., Soh, H.S., Liu, W., Tanner, L., Aw, P.P., Tan,
750 H.C., Ng, L.C., Leo, Y.S., Low, J.G., Ong, A., Ooi, E.E., Vasudevan, S.G., Hibberd, M.L.,
751 2009. Genomic epidemiology of a dengue virus epidemic in urban Singapore. *J.*
752 *Virol.* *2009*.
753 Sukri, N.C., Laras, K., Wandra, T., Didi, S., Larasati, R.P., Rachdyatmaka, J.R., Osok, S.,
754 Tjia, P., Saragih, J.M., Hartati, S., Listryaningsih, E., Porter, K.R., Beckett, C.G.,
755 Prawira, I.S., Punjabi, N., Suparmanto, S.A., Beecham, H.J., Bangs, M.J., Corwin,
756 A.L., 2003. Transmission of epidemic dengue hemorrhagic fever in easternmost
757 Indonesia. *Am. J. Trop. Med. Hyg.* *68*, 529–535.
758 Sukumaran, J., Holder, M.T., 2010. DendroPy: a Python library for phylogenetic
759 computing. *Bioinformatics* *26*, 1569–1571. <http://dx.doi.org/10.1093/bioinformatics/btq228>.
760 Sumarmo, 1987. Dengue haemorrhagic fever in Indonesia. *Southeast Asian J. Trop.*
761 *Med. Public Health* *18*, 269–274.
762 Suwandono, A., Kosasih, H., Nurhayati, Kusriastuti, R., Harun, S., Ma'roef, C.,
763 Wuryadi, S., Herianto, B., Yuwono, D., Porter, K.R., Beckett, C.G., Blair, P.J., 2006.
764 Four dengue virus serotypes found circulating during an outbreak of dengue
765 fever and dengue haemorrhagic fever in Jakarta, Indonesia, during 2004. *Trans.*
766 *R. Soc. Trop. Med. Hyg.* *100*, 855–862.
767 Twiddy, S.S., Farrar, J.J., Vinh Chau, N., Wills, B., Gould, E.A., Gritsun, T., Lloyd, G.,
768 Holmes, E.C., 2002. Phylogenetic relationships and differential selection
769 pressures among genotypes of dengue-2 virus. *Virology* *298*, 63–72.
770 Twiddy, S.S., Holmes, E.C., Rambaut, A., 2003. Inferring the rate and time-scale of
771 dengue virus evolution. *Mol. Biol. Evol.* *20*, 122–129.
772 Vaughn, D.W., Green, S., Kalayanarooj, S., Innis, B.L., Nimmannitya, S., Suntayakorn,
773 S., Endy, T.P., Raengsakulrach, B., Rothman, A.L., Ennis, F.A., Nisalak, A., 2000.
774 Dengue viremia titer, antibody response pattern, and virus serotype correlate
775 with disease severity. *J. Infect. Dis.* *181*, 2–9.
776 WHO, 1997. *Dengue Haemorrhagic Fever: Diagnosis, Treatment, Prevention and*
777 *Control.* World Health Organization, Geneva.
778 Yamanaka, A., Mulyatno, K.C., Susilowati, H., Hendrianto, E., Ginting, A.P., Sary, D.D.,
779 Rantam, F.A., Soegijanto, S., Konishi, E., 2011. Displacement of the predominant
780 dengue virus from type 2 to type 1 with a subsequent genotype shift from IV to I
781 in Surabaya, Indonesia 2008–2010. *PLoS One* *6*, e27322. <http://dx.doi.org/10.1371/journal.pone.0027322>.

3._Isra_genomic.pdf

ORIGINALITY REPORT

% **18**
SIMILARITY INDEX

% **14**
INTERNET SOURCES

% **13**
PUBLICATIONS

% **10**
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to West Coast University % **1**
Student Paper

2 M. J. Schreiber. "Genomic Epidemiology of a Dengue Virus Epidemic in Urban Singapore", Journal of Virology, 02/11/2009 % **1**
Publication

3 www.biomedcentral.com % **1**
Internet Source

4 www.jidc.org % **1**
Internet Source

5 www.ausstats.abs.gov.au % **1**
Internet Source

6 Sneha Singh, Birendra P. Gupta, Anoop Manakkadan, Krishna Das Manandhar, Easwaran Sreekumar. "Phylogenetic study reveals co-circulation of Asian II and Cosmopolitan genotypes of Dengue virus serotype 2 in Nepal during 2013", Infection, Genetics and Evolution, 2015 % **1**
Publication

7	eprints.gla.ac.uk Internet Source	% 1
8	www.greengazette.co.za Internet Source	% 1
9	Benediktus Yohan, Puspa Wardhani, Aryati, Hidayat Trimarsanto, R. Tedjo Sasmono. "Production of recombinant dengue non-structural 1 (NS1) proteins from clinical virus isolates", Protein Expression and Purification, 2017 Publication	% 1
10	eprints.qut.edu.au Internet Source	<% 1
11	Massimo Giuliani, Maria M. Santoro, Alessandra Lo Presti, Eleonora Cella et al. "Circulation of HIV-1 CRF02_AG among MSM Population in Central Italy: A Molecular Epidemiology-Based Study", BioMed Research International, 2013 Publication	<% 1
12	Dewi Megawati, Sri Masyeni, Benediktus Yohan, Asri Lestarini et al. "Dengue in Bali: Clinical characteristics and genetic diversity of circulating dengue viruses", PLOS Neglected Tropical Diseases, 2017 Publication	<% 1
13	Submitted to Universiti Malaysia Sabah	

<% 1

14

Endang S. Kusmintarsih, Rahma F. Hayati, Oktaviani N. Turnip, Benediktus Yohan et al. "Molecular characterization of dengue viruses isolated from patients in Central Java, Indonesia", Journal of Infection and Public Health, 2018

Publication

<% 1

15

Ong, S.H.. "Periodic re-emergence of endemic strains with strong epidemic potential-A proposed explanation for the 2004 Indonesian dengue epidemic", Infection, Genetics and Evolution, 200803

Publication

<% 1

16

Submitted to National University of Singapore

Student Paper

<% 1

17

arca.icict.fiocruz.br

Internet Source

<% 1

18

aos.wisc.edu

Internet Source

<% 1

19

hal.archives-ouvertes.fr

Internet Source

<% 1

20

124.16.173.210

Internet Source

<% 1

21

www.ajtmh.org

Internet Source

<% 1

22

R. Tedjo Sasmono, Lily Pertiwi Kalalo, Suryani Trismiasih, Dionisius Denis et al. "Multiple introductions of dengue virus strains contribute to dengue outbreaks in East Kalimantan, Indonesia, in 2015–2016", *Virology Journal*, 2019

Publication

<% 1

23

paduaresearch.cab.unipd.it

Internet Source

<% 1

24

Juan Fidel Osuna-Ramos, José Manuel Reyes-Ruiz, Patricia Bautista-Carbajal, Margot Cervantes-Salazar et al. "Ezetimibe inhibits dengue virus infection in Huh-7 cells by blocking the cholesterol transporter Niemann–Pick C1-like 1 receptor", *Antiviral Research*, 2018

Publication

<% 1

25

academic.oup.com

Internet Source

<% 1

26

R. Tedjo Sasmono, Anne-Frieda Taurel, Ari Prayitno, Hermin Sitompul et al. "Dengue virus serotype distribution based on serological evidence in pediatric urban population in Indonesia", *PLOS Neglected Tropical Diseases*, 2018

Publication

<% 1

27	www.mdpi.com Internet Source	<% 1
28	tsukuba.repo.nii.ac.jp Internet Source	<% 1
29	Submitted to Universiti Putra Malaysia Student Paper	<% 1
30	discovery.ucl.ac.uk Internet Source	<% 1
31	www.nbb.be Internet Source	<% 1
32	www.frontiersin.org Internet Source	<% 1
33	Submitted to Fakultas Ekonomi Universitas Indonesia Student Paper	<% 1
34	www.biologie.uni-regensburg.de Internet Source	<% 1
35	epdf.tips Internet Source	<% 1
36	Shubham Shrivastava, Divya Tiraki, Arundhati Diwan, Sanjay K. Lalwani, Meera Modak, Akhilesh Chandra Mishra, Vidya A. Arankalle. "Co-circulation of all the four dengue virus serotypes and detection of a novel clade of	<% 1

DENV-4 (genotype I) virus in Pune, India during 2016 season", PLOS ONE, 2018

Publication

37 www.scielo.sa.cr <% 1
Internet Source

38 www.dengue-qsnich.org <% 1
Internet Source

39 Submitted to University of Strathclyde <% 1
Student Paper

40 Submitted to University of Stirling <% 1
Student Paper

41 www.bioscirep.org <% 1
Internet Source

42 www.citeulike.org <% 1
Internet Source

43 Submitted to University of Sheffield <% 1
Student Paper

44 mobile.repository.ubn.ru.nl <% 1
Internet Source

45 Submitted to Michigan State University <% 1
Student Paper

46 med.unhas.ac.id <% 1
Internet Source

www.g3journal.org

47	Internet Source	<% 1
48	www.mdpi.org Internet Source	<% 1
49	robdunnlab.com Internet Source	<% 1
50	cswb.gov.in Internet Source	<% 1
51	tampub.uta.fi Internet Source	<% 1
52	elfosscientiae.cigb.edu.cu Internet Source	<% 1
53	www.journals.elsevier.com Internet Source	<% 1
54	digital.bl.fcen.uba.ar Internet Source	<% 1
55	Juthamas Phadungsombat, Marco Yung-Cheng Lin, Narinee Srimark, Atsushi Yamanaka et al. "Emergence of genotype Cosmopolitan of dengue virus type 2 and genotype III of dengue virus type 3 in Thailand", PLOS ONE, 2018 Publication	<% 1
56	Submitted to University College London Student Paper	<% 1

57	www.coursehero.com Internet Source	<% 1
58	www.chem.ucla.edu Internet Source	<% 1
59	Panyupa Pankhong. "Molecular Genetic Relationship of the 3' Untranslated Region Among Thai Dengue-3 Virus, Bangkok Isolates, During 1973–2000", DNA and Cell Biology, 06/29/2009 Publication	<% 1
60	intl-jvi.asm.org Internet Source	<% 1
61	repository.tudelft.nl Internet Source	<% 1
62	jmscr.igmpublication.org Internet Source	<% 1
63	pdfs.semanticscholar.org Internet Source	<% 1
64	eprints.soton.ac.uk Internet Source	<% 1
65	media.neliti.com Internet Source	<% 1
66	Submitted to London School of Hygiene and Tropical Medicine Student Paper	<% 1

67 vir.sgmjournals.org Internet Source <% 1

68 A. P. Goncalvez. "Chimpanzee Fab Fragments and a Derived Humanized Immunoglobulin G1 Antibody That Efficiently Cross-Neutralize Dengue Type 1 and Type 2 Viruses", Journal of Virology, 12/01/2004
Publication <% 1

69 dspace.lib.cranfield.ac.uk Internet Source <% 1

70 biotaxa.org Internet Source <% 1

71 journal.frontiersin.org Internet Source <% 1

72 oro.open.ac.uk Internet Source <% 1

73 scholarbank.nus.edu.sg Internet Source <% 1

74 bmcinfectdis.biomedcentral.com Internet Source <% 1

75 theses.gla.ac.uk Internet Source <% 1

76 Submitted to Heriot-Watt University Student Paper <% 1

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5
WORDS