

Aldian_Irma_Amaruddin_Microbiota.pdf

by Firdaus Hamid

FILE	ALDIAN_IRMA_AMARUDDIN_MICROBIOTA.PDF (2.24M)	WORD COUNT	6481
TIME SUBMITTED	23-JUL-2020 08:35PM (UTC+0700)	CHARACTER COUNT	40756
SUBMISSION ID	1361187803		



Article 11

The Bacterial Gut Microbiota of Schoolchildren from High and Low Socioeconomic Status: A Study in an Urban Area of Makassar, Indonesia

Aldian I. Amaruddin ^{1,2,*}, Firdaus Hamid ³, Jan Pieter R. Koopman ¹,
Munawir Muhammad ³, Eric A.T. Brienen ¹, Lisette van Lieshout ¹, Anoecim R. Geelen ^{4,5},
Sitti Wahyuni ², Ed J. Kuijper ^{4,5}, Erliyani Sartono ¹, Maria Yazdanbakhsh ¹ and
Romy D. Zwartink ^{4,5,*}

¹ Department of Parasitology, Leiden University Medical Center, 2333ZA Leiden, The Netherlands; J.P.R.Koopman@lumc.nl (J.P.R.K.); E.A.T.Brienen@lumc.nl (E.A.T.B.); E.A.van_Lieshout@lumc.nl (L.v.L.); Erliyani Sartono@lumc.nl (E.S.); M.Yazdanbakhsh@lumc.nl (M.Y.)

² Department of Parasitology, Faculty of Medicine, Hasanuddin University, Makassar 90245, Indonesia; Sitti Wahyuni@gmail.com

³ Department of Microbiology, Faculty of Medicine, Hasanuddin University, Makassar 90245, Indonesia; Firdaus Hamid@gmail.com (F.H.); dr.munawirmuhammad@yahoo.co.id (M.M.)

⁴ Experimental Bacteriology, Department of Medical Microbiology, Leiden University Medical Center, 3720 XZ Leiden, The Netherlands; A.R.Geelen@lumc.nl (A.R.G.); E.J.Kuijper@lumc.nl (E.J.K.)

⁵ Center for Microbiome Analyses and Therapeutics, Leiden University Medical Center, 2333ZA Leiden, The Netherlands

* Correspondence: A.I.Amaruddin@lumc.nl (A.I.A.); R.D.Zwartink@lumc.nl (R.D.Z.)

Received: 30 May 2020; Accepted: 22 June 2020; Published: 26 June 2020



Abstract: To understand the relationship between the gut microbiota and the health profile of Indonesians, it is important to elucidate the characteristics of the bacterial communities that prevail in this population. To this end, we profiled the faecal bacterial community of 140 Indonesian schoolchildren in urban Makassar. The core microbiota of Indonesian schoolchildren consisted of *Bifidobacterium*, *Collinsella*, and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families, but the relative abundance of these taxa varied greatly among children. Socioeconomic status (SES) was the main driver for differences in microbiota composition. Multiple bacterial genera were differentially abundant between high and low SES children, including *Bifidobacterium*, *Lactobacillus*, *Prevotella*, and *Escherichia-Shigella*. In addition, the microbiota of high SES children was less diverse and strongly associated with body mass index (BMI). In low SES children, helminth infection was prevalent and positively associated with *Penella*, *Enterohabdus*, *Lactobacillus*, and *Mogibacterium* abundance, while negatively associated with relative abundance of *Prevotella*. Protozoa infection was also prevalent, and positively associated with *Rikenellaceae*, while it was negatively associated with the relative abundance of *Romboutsia* and *Prevotella*. In conclusion, Indonesian schoolchildren living in urban Makassar share a core microbiota, but their microbiota varies in diversity and relative abundance of specific bacterial taxa depending on socioeconomic status, nutritional status, and intestinal parasites infection.

Keywords: gut microbiota; socioeconomic status; intestinal parasites; nutritional status; schoolchildren

1. Introduction

In the past decades, several studies have established the role of the gut microbiota in maintaining host physiological states, including immune responses, metabolism, mental, and physical

development [1]. The imbalance of gut microbiota composition may aggravate inflammation, metabolic diseases, or other health problems [2–5].

Gut microbiota composition is predominantly driven by environmental factors, such as diet, physical activity level, hygiene, disease, and medication use, instead of genetics [6,7]. Previous studies revealed notable differences in gut microbiota profiles among different ethnic and geographical areas, sometimes representing socioeconomic inequalities between groups [8–11]. A study involving European and African children has shown that the microbiota of African children is more diverse and contains more fibre-degrading, short-chain fatty acid producing bacteria than European children, attributable to differences in diet and lifestyle [12].

Indonesia is a developing country with enormous economic growth, but with great socioeconomic disparities in its population. This inequality is also reflected by the wide gap in health status between people from high and low socioeconomic status (SES). Low SES has been associated with lack of sanitation and bad hygiene and, as a consequence, higher exposure to soil-transmitted-helminths or other intestinal parasites, especially in schoolchildren. Recently, we investigated gut microbiota composition in a helminth endemic area in rural Indonesia. When comparing helminth infected with uninfected individuals, we observed no difference in bacterial composition or diversity of the gut microbiota [13]. Since there are no large disparities in lifestyle and SES in that area, these factors were not studied [27].

Given the role of the gut microbiota in health and susceptibility to diseases, it is important to investigate the gut microbiota in a specific population and determine what factors affect its composition. Here, we studied the bacterial gut microbiota, and its association with environmental factors, of Indonesian schoolchildren living in urban Makassar.

36

2. Methods

2.1. Study Design and Ethics Approval

This study was a cross-sectional study involving Indonesian schoolchildren living in an urban area of Makassar. Children from two public schools, which were distinct in socioeconomic level, were included in this study. The study was approved by the ethics committee for medical research of the Faculty of Medicine, Universitas Hasanuddin Indonesia (approval number: 1504/H04.8.4.5.31/PP36-KOMETIK/2016). Written, signed, and dated informed consent was obtained from the parents or guardian of each child prior to the study.

57

2.2. Study Population and Data Collection

One hundred and forty children were recruited from one high ($n = 74$) and one low SES ($n = 66$) school. The high SES school is located in the city centre and is considered of high status, with a majority of the parents working as high-skilled workers or professionals with higher education. Meanwhile, the low SES school is located near a landfill and port area, where most of the parents are low-educated and work on low-skilled labour jobs.

A standardised questionnaire was used to gather demographic information of the children, including age and sex. To determine the nutritional status, anthropometric measurement was performed. To assess the intestinal parasitic infection and microbiota composition, stool samples were collected using a stool container with enclosed spoon (Sarstedt AG&Co.KG, Nümbrecht, Germany). As soon as samples were collected by research staff, samples were stored inside an ice bag and transported to the Laboratory of Parasitology Department at Hasanuddin University to be aliquoted and stored at $-80\text{ }^{\circ}\text{C}$. The children's characteristics are presented in Table 1.

Table 1. Characteristics of study populations.

Characteristics	All Children (n = 140)	High SES Children (n = 74)	Low SES Children (n = 66)	p-Value
Age, years, mean ± SD	10.33 ± 0.85	10.40 ± 0.56	10.25 ± 1.04	0.294
Female: n (%)	85 (60.7)	41 (62.1)	44 (59.5)	0.748
zBMI, mean ± SD	−0.31 ± 1.46	0.33 ± 1.5	−0.89 ± 1.14	<0.001
Nutritional status (WHO): n (%)				<0.001
Thinness	13 (9.3)	2 (3.0)	11 (14.9)	
Normal	99 (70.7)	39 (59.1)	60 (81.1)	
Overweight and Obese	28 (20.0)	25 (37.9)	3 (4.1)	
Intestinal parasitic infection: n (%)				
Any helminth	47 (33.6)	2 (3.0)	45 (60.8)	<0.001
<i>Ascaris lumbricoides</i>	34 (24.3)	0	34 (45.9)	
<i>Trichuris trichiura</i>	25 (17.9)	2 (3.0)	23 (31.1)	<0.001
<i>Hymenolepis diminuta</i>	1 (0.7)	0	1 (1.4)	
Any protozoa	66 (47.1)	18 (27.3)	48 (65.8)	<0.001
<i>Entamoeba histolytica</i>	14 (10.0)	0	14 (19.2)	
<i>Dientamoeba fragilis</i>	30 (21.4)	10 (15.2)	20 (27.4)	0.080
<i>Giardia lamblia</i>	39 (27.9)	9 (13.6)	30 (41.1)	<0.001
<i>Cryptosporidium parvum</i>	2 (1.4)	0	2 (2.7)	

The number of positives (n) of the total population examined (n = 140). Statistical testing was performed using the student t-test for continuous variables and using the chi-square test for categorical variables. SD: standard deviation, SES: socioeconomic status.

2.3. Anthropometric Measurements

The height and weight of the participant were measured using a portable stadiometer (SECA GmbH & Co., Hamburg, Germany) and digital scale (GEA, Megapratama Medikalindo, Indonesia). The weighing scale was calibrated using standardised weight as part of routine care. Age-for-age (zBMI) was calculated according to the World Health Organization (WHO) references value to determine the nutritional status of the participants [14]. Children were categorised as thinness if zBMI < −2 SD, normal if zBMI was between −2 SD to 1 SD, and overweight/obese if zBMI > 1SD.

2.4. Parasitological Examination

A single-slide of each freshly-collected stool sample was examined by the Katokatz methods for the detection of soil transmitted helminth eggs. DNA was extracted from cryopreserved stool for intestinal protozoa identification using a QIAamp Spin Columns/Mini Kit (Qiagen, Germany) [15]. In each sample, a fixed amount of Phocine Herpes 1 was included within the isolation lysis buffer as an internal control [16]. A panel of multiplex real-time Polymerase Chain Reaction (PCR) was used to detect and quantify intestinal protozoa, targeting *Entamoeba histolytica*, *Dientamoeba fragilis*, *Giardia lamblia*, and *Cryptosporidium parvum*. The procedure has been described elsewhere [15,17].

2.5. Microbiota Analysis

DNA was extracted from approximately 0.1 g of cryopreserved stool using the Quick-DNA™ Fecal/Soil Microbe Miniprep Kit (Zymo Research, CA, USA) according to manufacturer instructions. Quality control, library preparation, and 16S rRNA gene amplicon sequencing were performed by BaseClear (Leiden, The Netherlands), targeting the V3-V4 region (F: CCTACGGGNGGCWGCAG, R: GACTACHVGGGTCTAATCC), and using the Illumina MiSeq platform (300 bp, paired-end, Illumina, CA, USA). Raw sequencing data are available in the European Nucleotide Archive (<https://www.ebi.ac.uk/ena>) under study accession PRJEB38465.

Read filtering, operational taxonomic unit (OTU)-picking, and taxonomic assignment were performed using the NG-Tax 0.4 pipeline with the following settings: forward and reverse read length of 120, ratio OTU abundance of 2.0, classify ratio of 0.9, minimum threshold of 1×10^{-7} , entity level of 100%, and an error correction of 98.5, using the Silva_132_SSU Ref database [18,19]. The obtained

OTU-table was filtered for OTUs with a number of sequences less than 0.005% of the total number of sequences [20].

Six positive controls (ZymoBiomics Microbial Community Standard, ZymoResearch, Leiden, The Netherlands) and six negative controls (empty extractions) were taken along from DNA extraction onwards, meaning one control per DNA extraction kit. In addition, four positive controls (ZymoBiomics Microbial Community DNA Standard, ZymoResearch, Leiden, The Netherlands) were taken along from Quality Control onwards, meaning one control per sequencing run. The eight bacterial species present in the included positive controls were all identified. The relative abundance of these species was on average 1.04 ± 0.20 and 1.06 ± 0.30 -fold different from the theoretical abundances for the sequencing control and DNA extraction control, respectively. This indicates that minor variation is induced by the sequencing and DNA extraction procedures. Four additional bacterial taxa were identified in the positive controls, namely *Collinsella*, *Bifidobacterium*, *Enterobacteriaceae*, and *Catenibacterium*, but their relative abundance only accounted for $0.017 \pm 0.013\%$ of total relative abundance. The included negative controls resulted in non-quantifiable DNA concentrations using the Qubit™ dsDNA HS Assay Kit (Thermo Fisher, Landsmeer, The Netherlands) on a Qubit 3.0 Fluorometer (Invitrogen, Breda, The Netherlands), but they were sequenced nevertheless, resulting in approximately ten times less reads than the stool samples, with *Delftia* and *Staphylococcus* as the most abundant contaminants.

2.6. Statistical Analysis

Prevalence rates were calculated as percentage of collected data and compared between schools using Pearson chi-square test. Comparisons between groups for continuous data were analysed using the Student *t*-test for normally distributed data, and using the Mann-Whitney-U test if the data were not normally distributed. This analysis was performed using IBM SPSS Statistics version 25. (IBM-SPSS Inc., Chicago, IL, USA).

Statistical analysis and data visualisation for microbiota data were performed in R (v3.6.1) using the packages phyloseq (v1.30.0) [21], vegan (v2.5-6) [22], ggplot2 (v3.2.1) [23], DESeq2 (v1.22.2) [24], and microbiome (v1.8.0) [25]. All analyses were performed on genus-level, except for alpha-diversity measures (Shannon diversity and observed richness). For differential abundance testing by DESeq2, genera present in less than 25% of the samples were removed to minimise zero-variance errors and spurious significance. Outcomes were considered significant when the Benjamini–Hochberg corrected *p*-value was ≤ 0.05 . Bacterial taxa were considered part of the core microbiota when present in 95% of the samples from the specified group. Permutational multivariate analysis of variance (PERMANOVA) was performed using the *adonis* function with 999 permutations and Bray–Curtis dissimilarity to determine associations between microbiota composition and clinical variables. Outcomes were considered significant when the *p*-value was ≤ 0.05 .

3. Results

3.1. The Bacterial Gut Microbiota of Indonesian Schoolchildren

To obtain a comprehensive overview of the bacterial gut microbiota of Indonesian school children, microbiota composition, as well as factors influencing their microbiota composition, were determined. The core microbiota consisted of *Bifidobacterium*, *Collinsella* and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families (Table 2), together constituting an average relative abundance of $56.3 \pm 19.8\%$. With an average relative abundance of 23.5% and 13.4%, *Bifidobacterium* and *Collinsella* were the most abundant members of the bacterial community (Table 3). However, their relative abundance varied greatly between children, ranging from 0.8 to 78.6% and 0.0 to 65.2%, respectively (Table 3). Large variation in relative abundance was also observed for the other most abundant bacterial taxa (Table 3). Variation in overall microbiota composition was significantly associated with socioeconomic status, as determined by the Bray–Curtis dissimilarity-based multivariate analysis (PERMANOVA; $R^2 = 0.049$; $p = 0.001$).

65
Table 2. Core microbiota in all, high SES, and low SES children.

Family	Genus	All Children	High SES Children	Low SES Children
Bifidobacteriaceae	Bifidobacterium	x	x	x
Coriobacteriaceae	Collinsella	x	x	x
Ruminococcaceae	Faecalibacterium	x	x	x
Ruminococcaceae	Subdoligranulum	x	x	x
Lachnospiraceae	-	x	x	x
Lachnospiraceae	Blautia	x	x	x
Lachnospiraceae	Dorea	x	x	x
Lachnospiraceae	Fusicatenibacter	x	x	x
Lachnospiraceae	Eubacterium hallii group	x	x	x
Lachnospiraceae	Ruminococcus torques group	x		x
Peptostreptococcaceae	Romboutsia		x	
Clostridiaceae 1	Clostridium sensu stricto 1		x	
Erysipelotrichaceae	Holdemanella			x
Eggerthellaceae	Senegalimassilia			x

Core taxa (x) are considered those present in 95% of the samples from the specified group.

34
Table 3. The ten most abundant bacterial taxa in all, high SES, and low SES children.

	Family	Genus	Average ± SD	Min–Max
All children	Coriobacteriaceae	Collinsella	0.235 ± 0.188	0.009–0.786
	Bifidobacteriaceae	Bifidobacterium	0.134 ± 0.139	0.000–0.652
	Erysipelotrichidae	Catenibacterium	0.077 ± 0.090	0.000–0.425
	Lachnospiraceae	Blautia	0.049 ± 0.047	0.001–0.208
	Prevotellaceae	Prevotella 9	0.044 ± 0.118	0.000–0.638
	Ruminococcaceae	Faecalibacterium	0.043 ± 0.056	0.000–0.322
	Erysipelotrichaceae	Holdemanella	0.031 ± 0.055	0.000–0.337
	Ruminococcaceae	Subdoligranulum	0.030 ± 0.035	0.000–0.224
	Lachnospiraceae	-	0.024 ± 0.038	0.000–0.255
	Coriobacteriaceae	Olsenella	0.024 ± 0.077	0.000–0.729
High SES children	Coriobacteriaceae	Collinsella	0.246 ± 0.185	0.013 ± 0.786
	Bifidobacteriaceae	Bifidobacterium	0.198 ± 0.152	0.004 ± 0.652
	Erysipelotrichidae	Catenibacterium	0.065 ± 0.082	0.000 ± 0.416
	Lachnospiraceae	Blautia	0.055 ± 0.050	0.001 ± 0.208
	Ruminococcaceae	Subdoligranulum	0.033 ± 0.038	0.000 ± 0.224
	Lachnospiraceae	-	0.030 ± 0.047	0.000 ± 0.255
	Ruminococcaceae	Faecalibacterium	0.029 ± 0.044	0.000 ± 0.266
	Ruminococcaceae	Ruminococcus 2	0.029 ± 0.059	0.000 ± 0.292
	Erysipelotrichaceae	Holdemanella	0.028 ± 0.059	0.000 ± 0.337
	Coriobacteriaceae	Olsenella	0.025 ± 0.064	0.000 ± 0.456
Low SES children	Coriobacteriaceae	Collinsella	0.225 ± 0.190	0.009 ± 0.737
	Erysipelotrichidae	Catenibacterium	0.088 ± 0.095	0.000 ± 0.425
	Bifidobacteriaceae	Bifidobacterium	0.077 ± 0.095	0.000 ± 0.403
	Prevotellaceae	Prevotella 9	0.071 ± 0.150	0.000 ± 0.638
	Ruminococcaceae	Faecalibacterium	0.055 ± 0.062	0.000 ± 0.322
	Lachnospiraceae	Blautia	0.043 ± 0.043	0.001 ± 0.170
	Erysipelotrichaceae	Holdemanella	0.035 ± 0.051	0.000 ± 0.267
	Ruminococcaceae	Subdoligranulum	0.029 ± 0.031	0.001 ± 0.183
	Coriobacteriaceae	Olsenella	0.023 ± 0.087	0.000 ± 0.729
	Lachnospiraceae	-	0.019 ± 0.026	0.000 ± 0.119

Relative abundances (average, min, max) are presented as a fraction of the total relative abundance. SD: standard deviation, SES: socioeconomic status.

To explore the relation between SES and the bacterial gut microbiota, microbiota composition and bacterial richness/diversity were compared between high and low SES children (Figure 1). Taxonomic profiles and principal coordinate analysis confirmed the difference, but also showed

the similarity, in overall microbiota composition between high and low SES children (Figure 1A,B). Differential abundance analysis revealed that, among others, *Bifidobacterium*, *Lactobacillus*, and various *Lachnospiraceae* and *Ruminococcaceae* members were more abundant in high SES children, while *Prevotella*, *Mogibacterium*, *Escherichia-Shigella*, and other members of the *Lachnospiraceae* and *Ruminococcaceae* families were more abundant in low SES children (Figure 1C). While bacterial richness was comparable, diversity was significantly higher in low SES children (Figure 1D).




Figure 1. Association between the bacterial gut microbiota and SES. (A) Taxonomic profiles of the gut microbiota of high and low SES children. The 20 most abundant bacterial taxa are shown. Relative abundance of all other taxa is summed and labelled as ‘other.’ (B) The Bray–Curtis dissimilarity-based principal coordinate analysis plot labelled according to SES. Ellipses indicate confidence assuming a multivariate t-distribution (solid line) or a multivariate normal distribution (dotted line). (C) Differential abundance of bacterial taxa between high and low SES children. Taxa with Benjamini–Hochberg corrected p -value below 0.05 are shown. (D) Alpha diversity in high and low SES children. Boxplots show the median, 25th and 75th percentiles, and minimal and maximal values with the exception of outliers (black circles).

3.2. The Bacterial Gut Microbiota of High SES Indonesian Schoolchildren

To obtain a comprehensive overview of the bacterial gut microbiota of high SES children ($n = 66$), microbiota composition, as well as factors influencing their microbiota composition, were determined. The core microbiota of high SES children consisted of *Bifidobacterium*, *Collinsella*, *Clostridium sensu stricto 1*, *Romboutsia*, and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families (Table 2), constituting an average relative abundance of $65.6 \pm 15.2\%$, with *Bifidobacterium* and *Collinsella* being most abundant (Table 3). Variation in overall microbiota composition was significantly associated with nutritional status (zBMI; PERMANOVA; $R^2 = 0.036$; $p = 0.011$).

To explore the relation between nutritional status and the bacterial gut microbiota, microbiota composition and bacterial richness/diversity were compared between normal weight and overweight/obese high SES children (Figure 2). Despite the association of zBMI with overall microbiota composition (Figure 2A), bacterial richness and diversity were similar between normal weight and overweight/obese children (Figure 2B), and only *Eggerthella* was significantly differentially abundant (Log2FoldChange = -3.49 ; padj = 0.005), with higher relative abundance in normal weight children.




Figure 2. Association between the bacterial gut microbiota and nutritional status in high SES children. (A) The Bray–Curtis dissimilarity-based principal coordinate analysis plot labelled according to nutritional status. Ellipses indicate confidence assuming a multivariate t-distribution (solid line) or a multivariate normal distribution (dotted line). (B) Alpha diversity in high SES children with normal weight and overweight/obesity. Boxplots show the median, 25th and 75th percentiles, and minimal and maximal values with the exception of outliers (black circles).

3.3. The Bacterial Gut Microbiota of Low SES Indonesian Schoolchildren

To obtain a comprehensive overview of the bacterial gut microbiota of low SES children ($n = 74$), microbiota composition, as well as factors influencing their microbiota composition, were determined. The core microbiota of low SES children consisted of *Bifidobacterium*, *Collinsella*, *Senegalimassilia*, *Holdemanella*, and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families (Table 2), constituting an average relative abundance of $54.4 \pm 20.8\%$, with *Collinsella* being the most abundant (Table 3). None of the included variables were significantly associated with overall microbiota composition in low SES children. Nevertheless, the high prevalence of helminth infection (60.8%) and protozoa infection (65.8%) in this group prompted us to study their association with microbiota composition.

To explore the relation between helminth or protozoa infection and the bacterial gut microbiota, microbiota composition and bacterial richness/diversity were compared between low SES children with and without these infections (Figure 3). Principal coordinate analysis confirmed that overall microbiota composition was not associated with helminth or protozoa status (Figure 3A,C). In addition, the bacterial richness and diversity were similar between children with and without helminth or protozoa infection (Figure 3B,D). Despite these commonalities, various bacterial taxa were differentially abundant. Relative abundance of *Olsenella*, *Enterorhabdus*, *Lactobacillus*, and *Mogibacterium* was higher in helminth infected children, whereas the relative abundance of *Prevotella* and unclassified *Lachnospiraceae* was higher in children without helminth infection (Figure 3E). In addition, the relative abundance of the *Rikenellaceae* RC9 gut group was higher in protozoa infected children, whereas the relative abundance of *Prevotella* and *Romboutsia* was higher in children without protozoa infection (Figure 3F).




Figure 3. Association between the bacterial gut microbiota and helminth and protozoa infection in low SES children. (A) The Bray–Curtis dissimilarity-based principal coordinate analysis plot labelled according to helminth infection. Ellipses indicate confidence assuming a multivariate t-distribution (solid line) or a multivariate normal distribution (dotted line). (B) Alpha diversity in low SES children that are helminth negative (HN) and helminth positive (HP). Boxplots show the median, 25th and 75th percentiles, and minimal and maximal values with the exception of outliers (black circles). (C) The Bray–Curtis dissimilarity-based principal coordinate analysis plot labelled according to protozoa infection. Ellipses indicate confidence assuming a multivariate t-distribution (solid line) or a multivariate normal distribution (dotted line). (D) Alpha diversity in low SES children that are protozoa negative (PN) and protozoa positive (PP). Boxplots show the median, 25th and 75th percentiles, and minimal and maximal values with the exception of outliers (black circles). (E) Differential abundance of bacterial taxa between helminth positive and helminth negative low SES children. Taxa with Benjamini–Hochberg corrected p -value below 0.05 are shown. (F) Differential abundance of bacterial taxa between protozoa positive and protozoa negative low SES children. Taxa with Benjamini–Hochberg corrected p -value below 0.05 are shown.

4. Discussion

Gut microbiota composition is largely influenced by environmental factors. In this study, socioeconomic status was the main factor associated with bacterial gut microbiota composition of Indonesian schoolchildren. Bacterial diversity was lower in children from high socioeconomic status, and their microbiota contains a higher relative abundance of *Bifidobacterium* and *Lactobacillus*, while containing a lower relative abundance of *Prevotella* and *Escherichia-Shigella*, among others. The gut microbiome plays a key role in host metabolism and immune functioning, and microbial dysbiosis during early life has been associated with the development of several diseases in later life, including atopy, obesity, chronic inflammatory conditions, and infections [26]. Alterations in microbiota composition as observed in our study may, therefore, contribute to the inequalities in health status and life expectancy of children from low and high socioeconomic status. Vice versa, early life exposures are likely to be affected by socioeconomic status-associated differences in lifestyle, education and healthcare, all which influence gut microbiota composition and may explain the observed differences in gut microbiota composition between low and high SES children.

Similar to our study, Chong CW et al. showed lower microbial diversity in wealthier children as compared to economically deprived children living in the same rural area in Malaysia [27]. Such differences in microbiota structure most likely result from lifestyle differences. Children from low SES are expected to have higher exposure to microorganisms and parasites [28]. In addition, diet is considered a main driver of gut microbiota composition [12]. As such, observed differences in microbiota composition between high and low SES children may be reflecting differences in their diet. High SES children are more likely to be exposed to high-fat diet products as compared to low SES children [29]. The high abundance of *Bifidobacterium* and *Lactobacillus* in high SES children could be associated with higher probiotics intake [30]. High SES children consume dairy products regularly, and probiotic drinks were available in the canteen of the high SES school, but not in the low SES school. *Prevotella*, observed to be of higher relative abundance in low SES children, has been associated with a higher intake of vegetables and fibre [31]. However, the influence of diet on microbiota composition remains speculative herein, as food intake was not determined in our study.

In high SES children, *Bifidobacterium*, *Collinsella*, *Clostridium sensu stricto 1*, *Romboutsia*, and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families were prevalent. Overall, the microbiota composition was mostly associated with nutritional status, and despite the differential abundance of *Eggerthella*, relative abundance of individual bacterial taxa did not significantly differ between normal weight and overweight/obese children, nor did bacterial richness and diversity. A recent study investigating Asian children from five different countries also found *Bifidobacterium* as the most abundant bacterium, among others [32]. In addition, nutritional status has previously been associated with microbiota composition in Mexican children [33]. It is important to realise that these differences may be a result of underlying variations in diet and environmental, geographical, demographic, and clinical factors. After correction for such factors, however, a significant association between microbiota and BMI has been shown in children who were part of the American Gut Project [34].

In low SES children, the core microbiota consisted of *Bifidobacterium*, *Collinsella*, *Senegalimassilia*, *Holdemanella*, and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families. Helminth infection was prevalent in low SES children, and was positively associated with *Olsenella*, *Enterorhabdus*, *Lactobacillus*, and *Mogibacterium* abundance, while it was negatively associated with the relative abundance of *Prevotella*. In addition, protozoa infection was prevalent in low SES children, and was also negatively associated with the relative abundance of *Prevotella*. We observed no apparent association between helminth or protozoa infection and microbiota diversity. Recently, there has been much interest in the relationship between helminth infection and the gut microbiota as they shared the same niche in the human body. Infection with intestinal helminths can impact the intestinal microbiome, with important consequences for each player in the tripartite relationship between the host, helminth, and the microbiota [35]. In a previous study conducted in rural Flores, Indonesia, where helminth infection is highly prevalent, no association between helminth infection and gut microbiota diversity

was observed [13]. In contrast to our study, increased diversity of gut microbiota has been observed in helminth colonised people in rural Malaysia [36]. Regarding protozoa, a previous study conducted in rural Cameroon showed higher alpha diversity in people infected by *Entamoeba* (*E. histolytica*, *E. dispar*, or both). Furthermore, *Entamoeba* presence was also associated with gut microbiota composition [37]. Discrepancies in findings most likely result from differences in the included population, population size, and the specific intestinal parasite species someone is infected with. Further studies are needed to elucidate the complex “three-way” relationship between intestinal parasites, microbiota, and the human host.

5. Conclusions

We demonstrated that Indonesian schoolchildren living in urban Makassar share a core microbiota consisting of *Bifidobacterium*, *Collinsella*, and multiple members of the *Lachnospiraceae* and *Ruminococcaceae* families, but that their microbiota varies in the diversity and relative abundance of specific bacterial taxa depending on socioeconomic status, nutritional status, and helminth and protozoa infection. These findings contribute to increased understanding of the environmental factors shaping gut microbiota composition in Indonesian schoolchildren.

Author Contributions: Conceptualisation, M.Y. and R.D.Z.; Formal analysis, A.I.A. and R.D.Z.; Funding acquisition, F.H. and M.Y.; Investigation, A.I.A., F.H., J.P.R.K., E.A.T.B., A.R.G., M.M. and R.D.Z.; Methodology, A.I.A. and R.D.Z.; Project administration, A.I.A., J.P.R.K., F.H., S.W.; Software, R.D.Z.; Supervision, L.v.L., E.J.K., E.S., M.Y. and R.D.Z.; Visualisation, A.I.A. and R.D.Z.; Writing—original draft, A.I.A. and R.D.Z.; Writing—review and editing, A.I.A., F.H., J.P.R.K., M.M., E.A.T.B., L.v.L., A.R.G., S.W., E.J.K., E.S., M.Y. and R.D.Z. All authors contributed substantially; approved the submitted version; and agree to be personally accountable for their own contributions and for ensuring that questions related to the accuracy or integrity of any part of the work, even parts in which the author was not personally involved, are appropriately investigated, resolved, and documented in the literature. All authors have read and agreed to the submitted version of the manuscript.

Funding: This work was supported by the Ministry of Research, Technology and Higher Education Republic of Indonesia (Kemenristekdikti) grant numbers: 123/SP2H/PTNBH/DPRM/2018, 6284/UN 4.1.2/PL.00.00/2018, 1674/UN4.21/PL.00.00/2018] and Leiden University Medical Center, The Netherlands. The sponsor of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Acknowledgments: The authors would like to thank all participants involved in this study as well as all their parents, teachers and headmasters of schools involved in this study, local government, Hasanuddin University Medical Research Center (HUM-RC) for laboratory facilities, and Hasanuddin University for the support. IPHC team who were involved in this study as well as our field team are gratefully acknowledged for their tremendous efforts on samples. The authors thank the Joint Scholarship of Directorate General of Resources for Science Technology and Higher Education (DGRSTHE) of Indonesia and Leiden University for providing PhD scholarship to A.I.A.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Liang, D.; Leung, R.K.; Guan, W.; Au, W.W. Involvement of gut microbiome in human health and disease: Brief overview, knowledge gaps and research opportunities. *Gut Pathog.* **2018**, *10*, 3. [CrossRef] [PubMed]
2. Li, X.; Watanabe, K.; Kimura, I. Gut Microbiota Dysbiosis Drives and Implies Novel Therapeutic Strategies for Diabetes Mellitus and Related Metabolic Diseases. *Front. Immunol.* **2017**, *8*, 1882. [CrossRef] [PubMed]
3. Lobionda, S.; Sittipo, P.; Kwon, H.Y.; Lee, Y.K. The Role of Gut Microbiota in Intestinal Inflammation with Respect to Diet and Extrinsic Stressors. *Microorganisms* **2019**, *7*, 271. [CrossRef]
4. Chiu, C.Y.; Chan, Y.L.; Tsai, M.H.; Wang, C.J.; Chiang, M.H.; Chiu, C.C. Gut microbial dysbiosis is associated with allergen-specific IgE responses in young children with airway allergies. *World Allergy Organ. J.* **2019**, *12*, 100021. [CrossRef] [PubMed]
5. Liu, X.; Tao, J.; Li, J.; Cao, X.; Li, Y.; Gao, X.; Fu, Y. Dysbiosis of Fecal Microbiota in Allergic Rhinitis Patients. *Am. J. Rhinol. Allergy* **2020**. [CrossRef]

6. Rothschild, D.; Weissbrod, O.; Barkan, E.; Kurilshikov, A.; Korem, T.; Zeevi, D.; Costea, P.I.; Godneva, A.; Kalka, I.N.; Bar, N.; et al. Environment dominates over host genetics in shaping human gut microbiota. *Nature* **2018**, *555*, 210–215. [CrossRef] [PubMed]
7. Scepanovic, P.; Hodel, F.; Mondot, S.; Partulla, V.; Byrd, A.; Hammer, C.; Alanio, C.; Bergstedt, J.; Patin, E.; Touvier, M.; et al. A comprehensive assessment of demographic, environmental, and host genetic associations with gut microbiome diversity in healthy individuals. *Microbiome* **2019**, *7*, 130. [CrossRef] [PubMed]
8. Brooks, A.W.; Priya, S.; Blekhan, R.; Bordenstein, S.R. Gut microbiota diversity across ethnicities in the United States. *PLoS Biol.* **2018**, *16*, e2006842. [CrossRef]
9. Yatsunenko, T.; Rey, F.E.; Manary, M.J.; Trehan, I.; Dominguez-Bello, M.G.; Contreras, M.; Magris, M.; Hidalgo, G.; Baldassano, R.N.; Anokhin, A.P.; et al. Human gut microbiome viewed across age and geography. *Nature* **2012**, *486*, 222–227. [CrossRef]
10. Deschasaux, M.; Bouter, K.E.; Prodan, A.; Levin, E.; Groen, A.K.; Herrema, H.; Tremaroli, V.; Bakker, G.J.; Attaye, I.; Pintof-Sietsma, S.J.; et al. Depicting the composition of gut microbiota in a population with varied ethnic origins but shared geography. *Nat. Med.* **2018**, *24*, 1526–1531. [CrossRef]
11. He, Y.; Wu, W.; Zheng, H.M.; Li, P.; McDonald, D.; Sheng, H.F.; Chen, M.X.; Chen, Z.H.; Ji, G.Y.; Zheng, Z.D.; et al. Regional variation limits applications of healthy gut microbiome reference ranges and disease models. *Nat. Med.* **2018**, *24*, 1532–1535. [CrossRef] [PubMed]
12. De Filippo, C.; Cavalieri, D.; Di Paola, M.; Ramazzotti, M.; Poullet, J.B.; Massart, S.; Collini, S.; Pieraccini, G.; Lionetti, P. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 14691–14696. [CrossRef] [PubMed]
13. Martin, I.; Djuardi, Y.; Sartono, E.; Rosa, B.A.; Supali, T.; Mitreva, M.; Houwing-Duistermaat, J.J.; Yazdanbakhsh, M. Dynamic changes in human-gut microbiome in relation to a placebo-controlled anthelmintic trial in Indonesia. *PLoS Negl. Trop. Dis.* **2018**, *12*, e0006620. [CrossRef] [PubMed]
14. De Onis, M.; Onyango, A.W.; Borghi, E.; Siyam, A.; Nishida, C.; Siekmann, J. Development of a WHO growth reference for school-aged children and adolescents. *Bull. World Health Organ.* **2007**, *85*, 660–667. [CrossRef]
15. Verweij, J.J.; Blange, R.A.; Templeton, K.; Schinkel, J.; Brienens, E.A.; van Rooyen, M.A.; van Lieshout, L.; Polderman, A.M. Simultaneous detection of *Entamoeba histolytica*, *Giardia lamblia*, and *Cryptosporidium parvum* in fecal samples by using multiplex real-time PCR. *J. Clin. Microbiol.* **2004**, *42*, 1220–1223. [CrossRef]
16. Niesters, H.G.M. Clinical Virology in Real Time. *J. Clin. Virol.* **2002**, *25*, 3–12. [CrossRef]
17. Verweij, J.J.; Mulder, B.; Poell, B.; van Middelkoop, D.; Brienens, E.A.; van Lieshout, L. Real-time PCR for the detection of *Dientamoeba fragilis* in fecal samples. *Mol. Cell. Probes* **2007**, *21*, 400–404. [CrossRef]
18. Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glockner, F.O. The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* **2013**, *41*, D590–D596. [CrossRef]
19. Ramiro-Garcia, J.; Hermes, G.D.A.; Giatsis, C.; Sipkema, D.; Zoetendal, E.G.; Schaap, P.J.; Smidt, H. NG-Tax, a highly accurate and validated pipeline for analysis of 16S rRNA amplicons from complex biomes. *F1000Research* **2016**, *5*, 1791. [CrossRef]
20. Bokulich, N.A.; Subramanian, S.; Faith, J.J.; Gevers, D.; Gordon, J.I.; Knight, R.; Mills, D.A.; Caporaso, J.G. Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nat. Methods* **2013**, *10*, 57–59. [CrossRef]
21. McMurdie, P.J.; Holmes, S. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* **2013**, *8*, e61217. [CrossRef]
22. Oksanen, J.; Blanchet, F.G.; Friendly, M.; Kindt, R.; Legendre, P.; McGlenn, D.; Minchin, P.R.; O'Hara, R.B.; Simpson, G.L.; Solymos, P.; et al. Vegan: Community Ecology Package. Available online: <https://CRAN.R-project.org/package=vegan> (accessed on 1 January 2020).
23. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*; Springer: New York, NY, USA, 2009.
24. Love, M.I.; Huber, W.; Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **2014**, *15*, 550. [CrossRef] [PubMed]
25. Lahti, L.; Shetty, S. Tools for Microbiome Analysis in R. Available online: <http://microbiome.github.com/microbiome> (accessed on 1 January 2020).
26. Tamburini, S.; Shen, N.; Wu, H.C.; Clemente, J.C. The microbiome in early life: Implications for health outcomes. *Nat. Med.* **2016**, *22*, 713–722. [CrossRef]

27. Chong, C.W.; Ahmad, A.F.; Lim, Y.A.; Teh, C.S.; Yap, I.K.; Lee, S.C.; Chin, Y.T.; Loke, P.; Chua, K.H. Effect of ethnicity and socioeconomic variation to the gut microbiota composition among pre-adolescent in Malaysia. *Sci. Rep.* **2015**, *5*, 13338. [[CrossRef](#)]
28. Strunz, E.C.; Addiss, D.G.; Stocks, M.E.; Ogden, S.; Utzinger, J.; Freeman, M.C. Water, sanitation, hygiene, and soil-transmitted helminth infection: A systematic review and meta-analysis. *PLoS Med.* **2014**, *11*, e1001620. [[CrossRef](#)] [[PubMed](#)]
29. Dinsa, G.D.; Goryakin, Y.; Fumagalli, E.; Suhrcke, M. Obesity and socioeconomic status in developing countries: A systematic review. *Obes. Rev.* **2012**, *13*, 1067–1079. [[CrossRef](#)] [[PubMed](#)]
30. Savard, P.; Lamarche, B.; Paradis, M.E.; Thiboutot, H.; Laurin, E.; Roy, D. Impact of *Bifidobacterium animalis* subsp. *lactis* BB-12 and *Lactobacillus acidophilus* LA-5-containing yoghurt, on fecal bacterial counts of healthy adults. *Int. J. Food Microbiol.* **2011**, *149*, 50–57. [[CrossRef](#)]
31. Precup, G.; Vodnar, D.C. Gut *Prevotella* as a possible biomarker of diet and its eubiotic versus dysbiotic roles: A comprehensive literature review. *Br. J. Nutr.* **2019**, *122*, 131–140. [[CrossRef](#)]
32. Nakayama, J.; Watanabe, K.; Jiang, J.; Matsuda, K.; Chao, S.H.; Haryono, P.; La-Ongkham, O.; Sarwoko, M.A.; Sujaya, I.N.; Zhao, L.; et al. Diversity in gut bacterial community of school-age children in Asia. *Sci. Rep.* **2015**, *5*, 8397. [[CrossRef](#)]
33. Lopez-Contreras, B.E.; Moran-Ramos, S.; Villarruel-Vazquez, R.; Macias-Kauffer, L.; Villamil-Ramirez, H.; Leon-Mimila, P.; Vega-Badillo, J.; Sanchez-Munoz, F.; Llanos-Moreno, L.E.; Canizalez-Roman, A.; et al. Composition of gut microbiota in obese and normal-weight Mexican school-age children and its association with metabolic traits. *Pediatric Obes.* **2018**, *13*, 381–388. [[CrossRef](#)]
34. Bai, J.; Hu, Y.; Bruner, D.W. Composition of gut microbiota and its association with body mass index and lifestyle factors in a cohort of 7–18 years old children from the American Gut Project. *Pediatric Obes.* **2019**, *14*, e12480. [[CrossRef](#)] [[PubMed](#)]
35. Rapin, A.; Harris, N.L. Helminth-Bacterial Interactions: Cause and Consequence. *Trends Immunol.* **2018**, *39*, 724–733. [[CrossRef](#)] [[PubMed](#)]
36. Lee, S.C.; Tang, M.S.; Lim, Y.A.; Choy, S.H.; Kurtz, Z.D.; Cox, L.M.; Gundra, U.M.; Cho, I.; Bonneau, R.; Blaser, M.J.; et al. Helminth colonization is associated with increased diversity of the gut microbiota. *PLoS Negl. Trop. Dis.* **2014**, *8*, e2880. [[CrossRef](#)]
37. Morton, E.R.; Lynch, J.; Froment, A.; Lafosse, S.; Heyer, E.; Przeworski, M.; Blekhman, R.; Séguirel, L. Variation in Rural African Gut Microbiota Is Strongly Correlated with Colonization by *Entamoeba* and Subsistence. *PLoS Genet.* **2015**, *11*, e1005658. [[CrossRef](#)] [[PubMed](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

ORIGINALITY REPORT

% **23**

SIMILARITY INDEX

% **11**

INTERNET SOURCES

% **20**

PUBLICATIONS

% **12**

STUDENT PAPERS

PRIMARY SOURCES

- 1** research-information.bris.ac.uk % **1**
Internet Source

- 2** Esther J. d'Haens, Romy D. Zwitterink, Clara Belzer, Marieke A. C. Hemels et al. "Short-term changes of intestinal microbiota composition in preterm infants after two prophylactic doses of vancomycin", *Acta Paediatrica*, 2019 % **1**
Publication

- 3** Hamid, Firdaus, Sitti Wahyuni, Astrid van Leeuwen, Ronald van Ree, Maria Yazdanbakhsh, and Erliyani Sartono. "Allergic disorders and socioeconomic status: a study of school children in an urban area of Makassar, Indonesia", *Clinical & Experimental Allergy*, 2015. % **1**
Publication

- 4** Submitted to American Intercontinental University Online % **1**
Student Paper

- 5** Lynn Meurs, Anton M. Polderman, Natalie V. S.

Vinkeles Melchers, Eric A. T. Brienen et al.
"Diagnosing Polyparasitism in a High-
Prevalence Setting in Beira, Mozambique:
Detection of Intestinal Parasites in Fecal
Samples by Microscopy and Real-Time PCR",
PLOS Neglected Tropical Diseases, 2017

Publication

% 1

6

Romy D Zwitterink, Ingrid B Renes, Richard A van
Lingen, Diny van Zoeren-Grobbe et al.

"Association between duration of intravenous
antibiotic administration and early-life microbiota
development in late-preterm infants", European
Journal of Clinical Microbiology & Infectious
Diseases, 2018

Publication

% 1

7

Yaqun Liu, Liguang Shang, Yuhua Zhan, Min Lin,
Zhu Liu, Yongliang Yan. "Genome-Wide
Analysis of Sugar Transporters Identifies the
gtsA Gene for Glucose Transportation in
Pseudomonas stutzeri A1501", Microorganisms,
2020

Publication

% 1

8

provost.uni.edu

Internet Source

% 1

9

Pieter F. de Groot, Tatjana Nikolic, Sultan
Imangaliyev, Siroon Bekkering et al. "Oral
butyrate does not affect innate immunity and

% 1

islet autoimmunity in individuals with longstanding type 1 diabetes: a randomised controlled trial", Diabetologia, 2020

Publication

10

MARIA M. M. KAISAR, ERIC A. T. BRIENEN, YENNY DJUARDI, ERLIYANI SARTONO et al. "Improved diagnosis of by using a bead-beating procedure on ethanol preserved stool samples prior to DNA isolation and the performance of multiplex real-time PCR for intestinal parasites ", Parasitology, 2017

Publication

% 1

11

www.terkko.helsinki.fi

Internet Source

<% 1

12

repositorio.ual.es

Internet Source

<% 1

13

Lara Pérez-Etayo, David González, Ana Isabel Vitas. "The Aquatic Ecosystem, a Good Environment for the Horizontal Transfer of Antimicrobial Resistance and Virulence-Associated Factors Among Extended Spectrum β -lactamases Producing E. coli", Microorganisms, 2020

Publication

<% 1

14

pericles.pericles-prod.literatumonline.com

Internet Source

<% 1

15

Noer Bahry Noor, Ridwan Amiruddin, Muhammad Awal, Sukri Palutturi, Anwar Mallongi. "Proxy Model of Comorbidities with Stroke Incident in South Sulawesi", *Pakistan Journal of Nutrition*, 2017

Publication

<% 1

16

Submitted to University of Minnesota System

Student Paper

<% 1

17

Ivonne Martin, Hae-Won Uh, Taniawati Supali, Makedonka Mitreva, Jeanine J. Houwing-Duistermaat. "The mixed model for the analysis of a repeated-measurement multivariate count data", *Statistics in Medicine*, 2019

Publication

<% 1

18

Alexis Rapin, Nicola L. Harris. "Helminth–Bacterial Interactions: Cause and Consequence", *Trends in Immunology*, 2018

Publication

<% 1

19

rbej.biomedcentral.com

Internet Source

<% 1

20

Naoki Ozato, Shinichiro Saito, Tohru Yamaguchi, Mitsuhiro Katashima et al. "Blautia genus associated with visceral fat accumulation in adults 20–76 years of age", *npj Biofilms and Microbiomes*, 2019

Publication

<% 1

-
- 21 Ilse R. Dubbelboer, Natasa Pavlovic, Femke Heindryckx, Erik Sjögren, Hans Lennernäs. "Liver Cancer Cell Lines Treated with Doxorubicin under Normoxia and Hypoxia: Cell Viability and Oncologic Protein Profile", *Cancers*, 2019
Publication <% 1
-
- 22 Justinn Renelies-Hamilton, Marc Noguera-Julian, Mariona Parera, Roger Paredes et al. "Exploring interactions between *Blastocystis* sp., *Strongyloides* spp. and the gut microbiomes of wild chimpanzees in Senegal", *Infection, Genetics and Evolution*, 2019
Publication <% 1
-
- 23 jissn.biomedcentral.com
Internet Source <% 1
-
- 24 hal.archives-ouvertes.fr
Internet Source <% 1
-
- 25 Submitted to Universiteit van Amsterdam
Student Paper <% 1
-
- 26 Submitted to Adtalem Global Education
Student Paper <% 1
-
- 27 Tamaki Ito, Tsuyoshi Sekizuka, Norimi Kishi, Akifumi Yamashita, Makoto Kuroda. "Conventional culture methods with commercially available media unveil the

presence of novel culturable bacteria", Gut
Microbes, 2018

Publication

28

Firdaus Hamid, Abena S. Amoah, Ronald van
Ree, Maria Yazdanbakhsh. "Chapter 5
Helminth-Induced IgE and Protection Against
Allergic Disorders", Springer Science and
Business Media LLC, 2015

Publication

29

Klaudyna Borewicz, Fangjie Gu, Edoardo
Saccenti, Ilja C.W. Arts et al. "Correlating Infant
Fecal Microbiota Composition and Human Milk
Oligosaccharide Consumption by Microbiota of
1-Month-Old Breastfed Infants", Molecular
Nutrition & Food Research, 2019

Publication

30

Christine P. Zolnik, Richard C. Falco, Thomas J.
Daniels, Sergios-Orestis Kolokotronis.
"Transient influence of blood meal and natural
environment on blacklegged tick bacterial
communities", Ticks and Tick-borne Diseases,
2018

Publication

31

Irek I. Mukhamatdinov, Aliya R. Khaidarova,
Rumia D. Zaripova, Rezeda E.
Mukhamatdinova, Sergey A. Sitnov, Alexey V.
Vakhin. "The Composition and Structure of

<% 1

<% 1

<% 1

<% 1

Ultra-Dispersed Mixed Oxide (II, III) Particles and Their Influence on In-Situ Conversion of Heavy Oil", Catalysts, 2020

Publication

32

Ali Mirza, Jessica D. Forbes, Feng Zhu, Charles N. Bernstein et al. "The multiple sclerosis gut microbiota: A systematic review", Multiple Sclerosis and Related Disorders, 2019

Publication

<% 1

33

www.eurosurveillance.org

Internet Source

<% 1

34

Janet Currie, Mark Stabile. "Socioeconomic Status and Child Health: Why Is the Relationship Stronger for Older Children?", American Economic Review, 2003

Publication

<% 1

35

Hamid, Firdaus, Aprilianto E. Wiria, Linda J. Wammes, Maria M. M. Kaiser, Yenny Djuardi, Serge A. Versteeg, Sitti Wahyuni, Ronald van Ree, Erliyani Sartono, Taniawati Supali, and Maria Yazdanbakhsh. "Risk Factors Associated with the Development of Atopic Sensitization in Indonesia", PLoS ONE, 2013.

Publication

<% 1

36

Submitted to Swinburne University of Technology

Student Paper

<% 1

37

Allen Ross, Keren Papier, Ruby Luceres-Catubig, Thao Chau, Marianette Inobaya, Shu-Kay Ng. "Poverty, Dietary Intake, Intestinal Parasites, and Nutritional Status among School-Age Children in the Rural Philippines", *Tropical Medicine and Infectious Disease*, 2017

Publication

<% 1

38

Peyton Ginakes, Julie M. Grossman, John M. Baker, Thanwalee Sooksa-nguan. "Living Mulch Management Spatially Localizes Nutrient Cycling in Organic Corn Production", *Agriculture*, 2020

Publication

<% 1

39

doaj.org

Internet Source

<% 1

40

Natalia Fraija-Fernández, Marie-Catherine Bouquieaux, Anaïs Rey, Iñaki Mendibil et al. "Marine water environmental DNA metabarcoding provides a comprehensive fish diversity assessment and reveals spatial patterns in a large oceanic area", *Ecology and Evolution*, 2020

Publication

<% 1

41

worldwidescience.org

Internet Source

<% 1

42

Submitted to University of Bath

Student Paper

<% 1

43 tropmedhealth.biomedcentral.com <% 1
Internet Source

44 repository.ubn.ru.nl <% 1
Internet Source

45 Masita Fujiko, Maisuri T. Chalid, Turyadi, Susan I. Ie et al. "Chronic hepatitis B in pregnant women: is hepatitis B surface antigen quantification useful for viral load prediction?", *International Journal of Infectious Diseases*, 2015 <% 1
Publication

46 Weizhong Dong, Ying Wang, Shuaixiong Liao, Minghang Lai, Li Peng, Gang Song. "Reduction in the Choking Phenomenon in Elite Diving Athletes Through Changes in Gut Microbiota Induced by Yogurt Containing *Bifidobacterium animalis* subsp. *lactis* BB-12: A Quasi Experimental Study", *Microorganisms*, 2020 <% 1
Publication

47 Petra Zimmermann, Nigel Curtis. "Breast milk microbiota: A complex microbiome with multiple impacts and conditioning factors", *Journal of Infection*, 2020 <% 1
Publication

48 hdl.handle.net <% 1
Internet Source

- 49 Annelies Verlaet, Nieke van der Bolt, Ben Meijer, Annelies Breynaert et al. "Toll-Like Receptor-Dependent Immunomodulatory Activity of Pycnogenol®", *Nutrients*, 2019
Publication <% 1
-
- 50 biomedpharmajournal.org
Internet Source <% 1
-
- 51 Valeria Marzano, Livia Mancinelli, Giorgia Bracaglia, Federica Del Chierico et al. "'Omic' investigations of protozoa and worms for a deeper understanding of the human gut 'parasitome'", *PLOS Neglected Tropical Diseases*, 2017
Publication <% 1
-
- 52 E.M. Terveer, E.A.T. Brienen, M.A.A. Erkens, L. van Lieshout. "Late manifestation of a mixed *Plasmodium falciparum* and *Plasmodium malariae* infection in a non-immune toddler after traveling to Chad", *Travel Medicine and Infectious Disease*, 2016
Publication <% 1
-
- 53 Submitted to University of Glasgow
Student Paper <% 1
-
- 54 peerj.com
Internet Source <% 1
-
- 55 bmccgenomics.biomedcentral.com
Internet Source <% 1

56 Leon Michael Hartman, Madeleine Josephine Henriette van Oppen, Linda Louise Blackall. "Microbiota characterization of *Exaiptasia diaphana* from the Great Barrier Reef", *Animal Microbiome*, 2020

Publication

<% 1

57 Tomohisa Takagi, Yuji Naito, Ryo Inoue, Saori Kashiwagi et al. "Differences in gut microbiota associated with age, sex, and stool consistency in healthy Japanese subjects", *Journal of Gastroenterology*, 2018

Publication

<% 1

58 P. Smith-Brown, M. Morrison, L. Krause, P. S. W. Davies. "Dairy and plant based food intakes are associated with altered faecal microbiota in 2 to 3 year old Australian children", *Scientific Reports*, 2016

Publication

<% 1

59 Diping Zeng, Zhiyi Liu, Song Bai, Jian Wang. "Influence of Sealing Treatment on the Corrosion Resistance of PEO Coated Al-Zn-Mg-Cu Alloy in Various Environments", *Coatings*, 2019

Publication

<% 1

60 Chin-An Yang, Chao Liang, Chia-Li Lin, Chiung-Tzu Hsiao, Ching-Tien Peng, Hung-Chih Lin, Jan-Gowth Chang. "Impact of *Enterobius*

<% 1

vermicularis infection and mebendazole treatment on intestinal microbiota and host immune response", PLOS Neglected Tropical Diseases, 2017

Publication

61

Simoës, C. D., J. Maukonen, J. Kaprio, A. Rissanen, K. H. Pietilainen, and M. Saarela. "Habitual Dietary Intake Is Associated with Stool Microbiota Composition in Monozygotic Twins", Journal of Nutrition, 2013.

Publication

<% 1

62

serval.unil.ch

Internet Source

<% 1

63

Paola León-Mimila, Hugo Villamil-Ramírez, Blanca López-Contreras, Sofía Morán-Ramos et al. "Low Salivary Amylase Gene (AMY1) Copy Number Is Associated with Obesity and Gut Prevotella Abundance in Mexican Children and Adults", Nutrients, 2018

Publication

<% 1

64

Submitted to University of Lancaster

Student Paper

<% 1

65

Gonneke W. J. M. Stevens, Wilma A. M. Vollebergh, Trees V. M. Pels, Alfons A. M. Crijnen. "Parenting and Internalizing and Externalizing Problems in Moroccan Immigrant Youth in the Netherlands", Journal of Youth and

<% 1

Adolescence, 2006

Publication

66 Submitted to University of Hull <% 1
Student Paper

67 pt.scribd.com <% 1
Internet Source

68 environhealthprevmed.biomedcentral.com <% 1
Internet Source

69 Alexia F. P. Lépine, Prokopis Konstanti, Klaudyna Borewicz, Jan-Willem Resink et al. "Combined dietary supplementation of long chain inulin and Lactobacillus acidophilus W37 supports oral vaccination efficacy against Salmonella Typhimurium in piglets", Scientific Reports, 2019 <% 1
Publication

70 Submitted to The University of Manchester <% 1
Student Paper

71 Arlette Santacruz, Ascensión Marcos, Julia Wärnberg, Amelia Martí et al. "Interplay Between Weight Loss and Gut Microbiota Composition in Overweight Adolescents", Obesity, 2009 <% 1
Publication

72 L. Kleinjans, D.H. Veening-Griffioen, T. Wehkamp, J. van Berghenhenegouwen et al. <% 1

"Mice co-administrated with partially hydrolysed whey proteins and prebiotic fibre mixtures show allergen-specific tolerance and a modulated gut microbiota", Beneficial Microbes, 2019

Publication

73

Submitted to University of Warwick

Student Paper

<% 1

74

Submitted to Imperial College of Science,
Technology and Medicine

Student Paper

<% 1

75

Submitted to Sussex Downs College

Student Paper

<% 1

EXCLUDE QUOTES ON

EXCLUDE ON

BIBLIOGRAPHY

EXCLUDE MATCHES

< 5
WORDS