

Nainu_et_al._HVI_Deseember_20 20.pdf

by

FILE	NAINU_ET_AL._HVI_DESEMBER_2020.PDF (2.27M)	WORD COUNT	13114
TIME SUBMITTED	22-FEB-2021 03:05PM (UTC+0700)	CHARACTER COUNT	69251
SUBMISSION ID	1515076440		

12

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344461457>

SARS-CoV-2 reinfection and implications for vaccine development

Article in *Human Vaccines and Immunotherapeutics* · December 2020

DOI: 10.1080/21645515.2020.1830683

CITATIONS

0

READS

253

8 authors, including:



Firzan Nainu

Universitas Hasanuddin

41 PUBLICATIONS 174 CITATIONS

[SEE PROFILE](#)



Rufika Shari Abidin

The University of Queensland

7 PUBLICATIONS 24 CITATIONS

[SEE PROFILE](#)



Muh. Akbar Bahar

Universitas Hasanuddin

15 PUBLICATIONS 129 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



What are the risks of ⁶⁸COVID-19 infection in pregnant women, neonates and breastfeeding women in Bangladesh [View project](#)



REVIEW SPECIAL ISSUE 2020 (Veterinary Quarterly, IF 2.34): Advances in safeguarding health and enhancing production in animals, wildlife and poultry - Deadline for Submissions: April 30, 2020 <https://think.taylorandfrancis.com/veterinary/> ³[View project](#)

MINI REVIEW



11
SARS-CoV-2 reinfection and implications for vaccine development

Firzan Nainu^{a,1}, Rufika Shari Abidin^{b,1}, Muh. Akbar Bahar^a, Andri Frediansyah^{c,d}, Talha Bin Emran^e, Ali A Rabaan^f,
Kuldeep Dhama^g, and Harapan Harapan^{h,i,j}

^aFaculty of Pharmacy, Hasanuddin University, 90245, Tamalanrea, Makassar, Indonesia; ^bFaculty of Medicine, Hasanuddin University, 90245, Tamalanrea, Makassar, Indonesia; ^cResearch Division for Natural Product Technology (BPTBA), Indonesian Institute of Sciences (LIPI), 55861, Wonosari, Indonesia; ^dDepartment of Pharmaceutical Biology, Pharmaceutical Institute, University of Tübingen, 72076, Tübingen, Germany; ^eDepartment of Pharmacy, BGC Trust University Bangladesh, 4381, Chittagong, Bangladesh; ^fMolecular Diagnostic Laboratory, Johns Hopkins Aramco Healthcare, 31311, Dhahran, Saudi Arabia; ^gDivision of Pathology, ICAR-Indian Veterinary Research Institute, 243122, Izatnagar, Bareilly, Uttar Pradesh, India; ^hMedical Research Unit, School of Medicine, Universitas Syiah Kuala, 23111, Banda Aceh, Indonesia; ⁱTropical Disease Centre, School of Medicine, Universitas Syiah Kuala, 23111, Banda Aceh, Indonesia; ^jDepartment of Microbiology, School of Medicine, Universitas Syiah Kuala, 23111, Banda Aceh, Indonesia

11 **TRACT**

Coronavirus disease 2019 (COVID-19) pandemic continues to constitute a public health emergency of international concern. Multiple vaccine candidates for COVID-19, which is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), have entered clinical trials. However, some evidence suggests that patients who have recovered from COVID-19 can be reinfected. For example, in China, two discharged COVID-19 patients who had recovered and fulfilled the discharge criteria for COVID-19 were retested positive to a reverse transcription polymerase chain reaction (RT-PCR) assay for the virus. This finding is critical and can hamper COVID-19 vaccine development. This review offers literature-based evidence of reinfection with SARS-CoV-2, provides explanation for the possibility of SARS-CoV-2 reinfection both from the agent and host points of view, and discusses its implication for COVID-19 vaccine development.

6

ARTICLE HISTORY

Received 6 June 2020
Revised 7 September 2020
Accepted 25 September 2020

KEYWORDS

SARS-CoV-2; covid-19;
reinfection; reactivation;
vaccine

10 **Introduction**

A novel coronavirus disease (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), emerged in central China in December 2019.^{1,2} Since then, the virus has spread worldwide, and more than 30 million cases and more than one million deaths have been reported, based on the COVID-19 Dashboard database.³ SARS-CoV-2, an enveloped and positive-sense single-stranded RNA virus, is a member of the genus *Betacoronavirus*, together with two highly pathogenic human viruses – severe acute respiratory syndrome coronavirus (SARS-CoV) and Middle East respiratory syndrome coronavirus (MERS-CoV).² Different studies have suggested that bats^{4,5} and pangolin^{6,7} might be the original hosts of SARS-CoV-2. Because of the genomic similarity between SARS-CoV-2 and SARS-CoV,^{4,5} these viruses are also suggested to have similar immunopathology.⁸

With no current specific treatments for COVID-19,^{9–11} a vaccine is expected to provide protection against SARS-CoV-2 infection. In an animal model study in which rhesus macaques after recovering from SARS-CoV-2 infection were re-exposed to SARS-CoV-2, viral replication was not detected in anal and nasopharyngeal swabs, suggesting the

protective effect of the primary infection.¹² This promising result is important for vaccine development, as it suggests that vaccination could be an effective protective measure against SARS-CoV-2 infection. Currently, multiple vaccine candidates have entered clinical trials; more than 100 are in the vaccine development pipeline,^{13,14} and their consumer acceptance¹⁵ as well as the willingness to purchase¹⁶ the vaccine candidates have been assessed. However, recent studies have reported that some COVID-19 patients that have recovered and fulfilled the discharge criteria for COVID-19 continued to show a positive result to reverse transcription polymerase chain reaction (RT-PCR) test for the virus.^{17–21} Since these findings are critical to the design of a vaccine against the virus, scientists are still debating the authenticity of these results to determine whether reinfection is possible. In SARS-CoV and MERS-CoV infections, conflicting findings have been reported,^{22–29} and hence, no licensed vaccine is currently available against the viruses.^{30,31} In this review, we systematically review the evidence of repositive RT-PCR test results for those who have been declared free of COVID-19, explain the possibility of COVID-19 reinfection from the virus and host points of view, and discuss the implication of this reinfection possibility for the development of COVID-19 vaccine.

CONTACT Firzan Nainu firzannainu@unhas.ac.id of Pharmacy, Hasanuddin University Jl. Perintis Kemerdekaan Km.10, Tamalanrea 90245, Makassar, Indonesia; Harapan Harapan harapan@unsyiah.ac.id Medical Research Unit, School of Medicine, Universitas Syiah Kuala, Jl. T. Tanoeh Abe, Darussalam, Banda Aceh 23111.

¹co-first authors

© 2020 Taylor & Francis Group, LLC

SARS-CoV-2 reinfection: the evidence from animal models and patients

MERS-CoV has been reported to reinfect camels, and neutralizing antibodies (nAbs), while not providing full immunity, could still reduce the viral load.^{25,26} In humans, MERS-CoV infection induced immunoglobulin G (IgG) production and sustained antibody levels against spike protein to clear the viral load.^{22,23} For SARS-CoV infection, studies on mice, Syrian hamsters, as well as rhesus and cynomolgus monkeys reported that SARS-CoV reexposure conferred resistance and did not enhance the severity of the disease.^{27–29} In SARS patients, IgG antibodies produced after SARS-CoV infection could neutralize the virus and prevent reinfection by the same virus for up to 2 years.²⁴ In a small ⁶³ animal model study on rhesus macaques, animals that had ⁷⁹ previously tested positive for SARS-CoV-2 and then had a negative RT-PCR test result after treatment, showed no viral replication by RT-PCR assay of anal and nasopharyngeal swabs after SARS-CoV-2 reexposure.¹² Besides, none of the COVID-19 symptoms was observed, ¹³ is why scientists believe that people who recover from SARS-CoV-2 infection will produce antibodies and be immun ⁷² reinfection.

Currently, the detection of SARS-CoV-2 from patients' samples relies on the use of a molecular-based diagnostic approach, such as RT-PCR.³² With a high level of precision and definitive ⁶¹ to produce reliable results, this method is recognized as the current gold standard for SARS-CoV-2 detection.³³ However, despite its decisive role in the detection of viral genomes, the RT-PCR method has some limitations that may lead to misdiagnosis in the state of infection.³³ Therefore, other clinical characteristics need to be evaluated and chest computed tomography scan needs to be performed before patients are discharged from hospitals.^{20,34–36} According to China CDC,²⁰ a COVID-19 patient should meet the following discharge criteria before discharge from hospital: (1) afebrile state for more than three consecutive days; (2) improved respiratory ⁴³ symptoms (no cough and expectoration, normal ranges of interleukin-6 (IL-6) and C-reactive protein (CRP) as well as oxygenation index ≥ 350); (3) improved chest radiography; and (4) negative RT-PCR results for two consecutive tests with sampling interval of at least 24 h. Similar criteria are also issued by the US CDC,³⁷ which include negative real-time RT-PCR results of nasopharyngeal and throat swabs for at least two consecutive tests with sampling interval of at least 24 ⁶⁰ febrile state, and improvement in signs and symptoms of COVID-19.

The incidence of ⁵⁹ reinfection in COVID-19 patients who have recovered and had a negative RT-PCR test has been brought to ⁴² public attention. In China, two discharged COVID-19 patients (39-year-old woman and 50-year-old man) were retested positive to the RT-PCR assay for the virus.¹⁹ Both patients had been treated with lopinavir-ritonavir, provided with supportive care, and were discharged after meeting the discharge criteria set by CDC China. Some medical workers also showed positive RT-PCR test approximately 5–13 days after being discharged from hospital in the Hubei province of China.²¹ Another report from China showed that 14.5% (n = 172) patients tested positive by RT-PCR after having been discharged from hospital (the median

age was 28 years; patients included child ⁵⁸ below 12 years (n = 6)).²⁰ Korean CDC reported repositive RT-PCR test results for recovered COVID-19 patients in Sejong city (25.9%, n = 27), Daegu city (27.2%, n = 195), and Gyeongbuk province (48.9%, n = 47).¹⁸ Approximately 59.6% of the re-positive cases had no symptoms, while the rest presented symptoms such as sore throat and cough.¹⁸ A list of studies that have reported ⁵⁷ reinfection with COVID-19 is presented in Table 1. In short, there is clear evidence that reinfection with SARS-CoV-2 is possible in humans, which should be considered in the development of an effective vaccine.

While the concern of SARS-CoV-2 reinfection cases have been raised, it is important to note that misdiagnosis for repositivity may occur due to laboratory errors. Many steps during SARS-CoV-2 detection by RT-qPCR are prone to human and/or technical faults. Samples mishandling, problems in sensitivity and/or specificity of reagents and/or techniques used to detect the viral RNA, or even record keeping are examples of error sources that may cause atypical pattern of laboratory results, as suggested.³⁸ Indeed, the newly developed ⁷⁸ COVID-19 reagents or even primer sets currently used in the RT-qPCR detection of SARS-CoV-2 RNA are factors that cannot be ruled out from the error equation.

SARS-CoV-2 reinfection: evidence in favor of and against

As an obligate intracellular parasite, a virus heavily relies on robust activity ⁵⁶ host organelles, particularly ribosomes, to propagate and produce new virions that are ready to infect other healthy cells.³⁹ To facilitate efficient transmission and its continuous existence, a virus requires sequential interaction between the infected host and other potential healthy host(s). Unfortunately, human cells are among the victims of pathogenic organisms with insidious viral life cycle, which include SARS-CoV-2.^{40,41} To survive infection, humans are equipped with two arms of immune responses, innate and adaptive immunity, which provide protection from invasion by pathogenic microbes.⁴² Several studies have been performed world- ⁷ wide to reveal the molecular nature and characteristics of SARS-CoV-2 and the clinical manifestations of COVID-19.³² However, ⁵⁵ despite massive global cooperation, information about the host immune response characteristics against SARS-CoV-2 remains limited.

Fortunately, accumulative experience from other ⁴⁷ zoonotic human coronavirus infections, which include SARS-CoV and MERS-CoV as well as from other closely related animal coronaviruses, has provided valuable insights into the viral structure, replication, potential ways of transmission, organ targets, clinical features, and possible pathogenic mechanisms of SARS-CoV-2.^{32,43,44} This information is important in fostering our efforts to mitigate COVID-19 transmission, accelerate disease management, and promote empirical yet rational pharmacological as well as nonpharmacological interventions.

Cellular and humoral antiviral immune responses, particularly adaptive immunity, are important players in the continuous host protection against cytopathic viruses.^{45,46} The

Table 1. List of reinfection or reactivation cases of COVID-19 reported worldwide as of July 25, 2020.

Country	Number of cases	Age years (range)	Sex	Days after negative test/discharged (range)	Characteristics		Reference
					Sample	Sample	
China	25	Median: 28 (IQR: 16–42); 6 children under 12	8 (M), 17 (F)	Mean: 7.32 (±SD: 3.86)	Cloacal and nasopharyngeal swab	38	
China	4	30–36	2 (M), 2 (F)	5–13	Throat swabs	21	
China	7	NA	4 (M), 3 (F)	Mean: 11 (8–15)	Throat swab	39	
China	22	Median: 28 (2–55)	8 (M), 14 (F)	Mean: 4.7 (2–13)	Throat and anal swabs	40	
China	1	54	1 (M)	5	Throat swabs and sputum	41	
China	1	46	1 (F)	4	Oropharyngeal swab	32	
China	1	44	1 (M)	3	Throat swabs	42	
China	4	Median: 71.5 (37–73)	2 (M), 2 (F)	5–14	Sputum and fecal specimens	43	
China	5	Under 16	NA	NA	NA	44	
China	15	Median: 64 (IQR: 51–73)	9 (M), 6 (F)	NA	Throat swab samples or deep nasal cavity	45	
China	7	Median: 6 (1–35)	6 (M), 1 (F)	7–11	Rectal and throat swabs	46	
China	8	Median: 59 (26–72)	3 (M), 5 (F)	Mean: 15 (7–30)	Deep nasal cavity or throat swab	47	
China	5	Median: 31 (27–42)	2 (M), 3 (F)	Mean: 10.6 (4–17)	Throat swab	48	
China	3	Median: 36 (34–74)	1 (M), 2 (F)	Mean: 9.3 (7–12)	Nasopharyngeal swab	49	
China	1	8	1 (M)	10	Throat swab	50	
China	2	56 and 21	1 (M), 1 (F)	17 and 7	throat and anus swab	51	
China	1	41	1 (M)	18	Nasal swabs, sputum, and stool	52	
China	1	46	1 (F)	6	Throat swab	53	
China	2	40s and 20s	1 (M), 1 (F)	6 and 7	Throat swab and Stool	54	
China	11	Mean: 48 (33–72)	3 (M), 8 (F)	Mean: 16.00 (6–27)	Oropharyngeal swab	55	
Italy	1	48	1 (M)	30	Nasopharyngeal swab	56	
China	11	Median: 66 (34–80)	4 (M), 7 (F)	NA	Throat swab	57	
Italy	2	81 and 85	2 (F)	2 and 5	Nasopharyngeal swab	58	
China	3	NA	NA	7	Salivary and fecal samples	59	
China	6	Median: 45 (30–56)	6 (F)	Mean: 10 (9–15)	NA	60	
China	27	Mean: 41.3 ± 17.0	16 (M), 11 (F)	11	Nasopharyngeal and throat swabs	61	
Brunei Darussalam	264	NA	NA	1–60	Nasopharyngeal swab	62	
China	61	Median: 54.79	25 (M), 36 (F)	3–35	Nasal, pharyngeal, Stool and sputum	63	
China	24	NC	NC	NC	Respiratory specimen	64	
China	11	Median: 27 (4–58)	7 (M), 4 (F)	Median: 14 (9–17)	Nasopharyngeal swab	65	
China	69	0–86	28 (M), 41 (F)	5–25	Throat swabs	66	
China	11	Median: 49 (37–62)	6 (M), 5 (F)	NC	Pharyngeal swabs	67	
China	20	Mean: 37.2 (4–80)	12 (M), 8 (F)	7–47	Nasopharyngeal and anal swabs	68	
China	2	55 and 68	1 (M), 1 (F)	16–29	Nasopharyngeal swab, sputum	69	
China	10	NC	NC	4–24	Nasopharyngeal and anal swabs	70	
China	20	Median: 41.5 (1–72)	7 (M), 13 (F)	7–14	Nasopharyngeal and anal swabs	71	
China	17	Median: 54 (IQR: 44–63)	5 (M), 12 (F)	Median: 4 (IQR: 3–8.5)	Sputum and nasopharyngeal swabs	72	
China	53	Mean: 62.19	23 (M), 30 (F)	1–12	Throat swabs	73	

presence of memory T cells and B cells with the ability to produce antibodies immediately upon reintroduction of a pathogen provides steadfast response and a high level of protection to the host,^{39,45} which in turn leads to continuous protective immunity. However, in the event of reinfection Table 1, this particular immunological concept is profoundly challenged.

Recent reports showed that B cells play an important role in the clearance of SARS-CoV-2, mostly through the production of nAbs.^{47,48} However, the duration of protection by these nAbs remains unknown. With the detection of SARS-CoV-2 in discharged patients, which is currently considered infection,^{20,32,49–52} many have questioned whether prolonged immune protection against SARS-CoV-2 truly prevails. In this section, the results and evidence from different fields are discussed and possible explanations on the caveats of host defenses that may lead to relapse and/or reinfection are provided. The possible explanations or arguments are provided based on two factors: the host and the agent.

Host point of view

Accumulated evidence has demonstrated that COVID-19 patients with moderate to severe disease level suffer from

different types of symptoms, including decreased number of lymphocytes known as lymphopenia.^{51,53,54} Lymphocytes, which comprise T and B cells, are the sole players of human adaptive immunity,^{45,46} and therefore, any deleterious effect on the quality and quantity of these cells may cause serious consequences on the integrity of host anti-SARS-CoV-2 immune responses.

A recent report suggested that lymphopenia occurs because of the extensive killing of lymphocytes in COVID-19 patients, and this event is apparently linearly correlated with IL-6 level and Fas-FasL interactions.⁵⁵ Lymphopenia may occur as a collateral damage because of increasing pro-inflammatory cytokine levels in COVID-19 patients^{41,56,57} or because of direct consequence of SARS-CoV-2 infection on lymphocytes.⁵⁸ Both these proposed mechanisms in cell death-mediated lymphocyte reduction in COVID-19 patients are illustrated in Figure 1A. However, whether SARS-CoV-2 virions can infect lymphocytes and directly promote the destruction of infected lymphocytes remains an interesting aspect of investigation.

Lymphopenia in COVID-19 patients correlates with disease severity,^{51,56} probably owing to the important role of lymphocytes in providing adaptive protection against viral infections such as SARS-CoV-2.⁸ In general, patients with lymphopenia most likely have low levels of B and/or T memory cells.⁵⁹




Figure 1. Potential causes of lymphopenia in COVID-19 patients. (A) Lymphopenia is possible to occur as a result of T cell depletion due to either SARS-CoV-2-mediated or IL-6-mediated pyroptotic cell death. (B) Alternatively, lymphocyte count in circulation may be reduced due to massive infiltration of lymphocytes into infected tissues (e.g. alveolus) (created with BioRender).

Indeed, extensive infiltration of lymphocytes into infected tissues or organs, including the lungs, as suggested by recent reports,^{60,61} may result in low lymphocyte numbers **Figure 1B**. Nevertheless, owing to the roles of memory lymphocytes in providing prolonged antiviral protection,⁴⁵ the decreased quantity (as well as quality) of these cells will dampen the host immune responses against reinfection.⁵⁴ In the context of SARS-CoV-2 infection, lymphopenia may lead to suboptimal production of anti-SARS-CoV-2 nAbs and/or reduced activities of CD4 helper T cells as well as CD8 cytotoxic T cells.⁴¹ Taking that into account, it is possible that recovered COVID-19 patients with lymphopenia history may have increased vulnerability to SARS-CoV-2 reinfection.

As mentioned before, the occurrence of lymphopenia is associated with increased levels of proinflammatory cytokines such as IL-6.⁵⁵ Normally, proinflammatory cytokines are expressed as an initial response to the presence of foreign materials, including viral particles, and the expression decreases gradually during resolution of the inflammation.⁸ These foreign materials can be characterized as pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs).⁶² Both PAMPs and DAMPs are detected by pattern recognition receptors (PRRs) in a manner dependent on their unique structures⁶³ leading to the activation of a signaling cascade via NF- κ B and/or IRF pathways to produce antiviral molecules/effectors and/or pro-inflammatory cytokines.^{62,64} Elevated levels of pro-inflammatory cytokines have been suggested as one of the drivers of lymphocytes killing in COVID-19 patients.^{40,41,65}

As the expression of proinflammatory cytokines is one of the most tightly regulated processes in the human body,^{42,46} it

is important to investigate the reasons for the breach of such tightly regulated processes upon SARS-CoV-2 infection. Nonetheless, based on the well-known tripartite correlation between increased level of proinflammatory cytokines, cell death, and tissue injury,⁴⁶ investigational studies to answer these questions will probably uncover the fundamental aspects on why reinfections occur and patients that are likely to experience reinfection, if reinfection is possible in the first place.

Another striking report regarding the characteristics of COVID-19 patients and lymphopenia is the presence of PD1 and TIM3, which are markers of T cell exhaustion.⁵⁶ In general, T cell exhaustion is characterized by inadequate effector function, persistent expression of inhibitory markers such as PD1 and TIM3, and a distinctive transcriptional profile compared with that of normal effector and/or memory T cells.⁶⁶ It has been suggested that these exhaustion characteristics cause insufficient T cell-mediated control of chronic infection,^{66,67} which may provoke relapse and/or recurrent infection.

In general, host immune responses are rapidly activated in the event of viral infection,⁴⁵ and the induction of humoral and cellular immune responses has been documented during human coronavirus 229E,⁶⁸ SARS-CoV,^{69,70} and MERS-CoV infections.^{71,72} This adaptive immunity might persist in the recovered patients for up to several years,⁷³ thus implying that reinfection is plausible after the levels of memory cells and antibodies of adaptive immunity are diminished. However, in the context of COVID-19, this alternative scenario might not be the case as reinfections were reported soon after the patients were discharged from hospitals **Table 1**.^{20,32,49-52}

Thus, while all of our speculations **Figure 2** remain to be demonstrated experimentally, reintroduction of SARS-CoV-2




Fig 762. Possible cause(s) of re-positivity of COVID-19 patients. Status of SARS-CoV-2 negative patients are decided based on at least two consecutive negative results on SARS-CoV-2 presence in patients' samples using reverse transcriptase-quantitative polymerase chain reaction (RT-qPCR). Plausible assumptions of patients who turn positive RT-qPCR after discharged from hospital are based on the status of patients' adaptive immunity (exhaustive memory cells and/or depleted memory cells) and the origin of SARS-CoV-2 present in the patients' samples (either already present in a low copy number or obtained by reexposure). The scarce evidence is unable to conclude either this as reinfection or relapse cases (created with BioRender).

into convalescent individuals with lymphopenia history may not only cause reinfection but also provide clues about the unexplored potential of convalescent plasma therapy in the management of COVID-19.

Virus point of view

In response to the COVID-19 pandemic, rapid and reliable methods for the detection of SARS-CoV-2 are urgently needed. Currently, the RT-PCR technique can be adopted to vigorously assess samples within several hours. The fundamental principle of this method is the detection of SARS-CoV-2 genetic material in a manner dependent on the action of reverse transcriptase and optimal amplification by DNA polymerase.⁷⁴ However, false negative results can occur owing to a really low level of starting genetic materials present in the examined samples Figure 2, leading to misinterpretation of supposedly positive results as negative for SARS-CoV-2.

To this date, most of reinfection cases were reported from China despite the fact that COVID-19 pandemic has been occurred for several months in more than 200 countries. Many factors, including laboratory errors as described above, may play a role in the detection of false-positive reinfection cases. In addition to that, variation of swab sites for SARS-CoV-2 sample detection, oral, anal, sputum, saliva, and nasopharyngeal swabs, may yield different results, as suggested in recent publications.^{75,76} In respect to the immunogenicity data, there is a possibility that patients may yield different results; tested negative from samples collected in one site but potentially positive in samples taken from other swab sites. With this in mind, it might be important to compare laboratory data using the same swab sites and diagnostic parameters before jumping to conclusion.

SARS-CoV-2, with ssRNA as its genome, uses RNA-dependent RNA polymerase (RdRp) in its replication, and owing to its error-prone tendency, RdRp may produce progenies with slightly mutated genes.⁷⁷ A recent report has suggested the presence of mutations in SARS-CoV-2 genome, particularly in ORF1ab, ORF8, and N genes.⁷⁸ The RT-PCR method used for the detection of SARS-CoV-2 heavily relies on the amplification of particular sequences in SARS-CoV-2 genes; hence, mutation at primer and/or probe sites may cause false-negative detection, and SARS-CoV-2-infected persons may show negative results. However, with several uncertainties in the characteristics of the SARS-CoV-2 genome, it remains difficult to conclude whether the mutations identified in the viral genome are responsible for the improper detection of SARS-CoV-2 in the clinical setting.

Alternatively, certain mutations in the viral RNA genome may lead to changes in epitope structure and/or characteristics and thus result in the production of progenies with new antigenic determinants, as shown in the case of influenza virus.⁷⁹ If reinfection occurs, the antibodies produced in the previous (primary) infection may not be able to recognize the epitope of the new virus. In this case, host defense towards infection will start from the beginning (from innate to adaptive immune activation), which eventually will result in either a successful recovery or reinfection.

Previous studies have pointed out the possible involvement of viral evasion strategies in the pathogenesis of SARS-CoV-2 in humans.^{8,80} As seen in cases of HIV, HBV, and HCV, evasion of host defense is one of the strategies used by the virus to initiate a chronic state of infection.^{81,82} Chronic infection has been linked to the reemergence of infection or relapse state. Relapse is different from reinfection; relapse is described as a recurrent infection with the same type of pathogen that was present beforehand, whereas reinfection is defined as the emergence of infection with a different species or serologic strain of pathogen, as seen in other infectious diseases.^{83,84} It is yet unclear whether relapse or reinfection occurs in COVID-19 patients.

An experimental study to assess the possible occurrence of reinfection using rhesus macaque showed that primary SARS-CoV-2 infection could provide adequate protection against subsequent exposure.^{85,86} Nevertheless, while the current data from a SARS-CoV-2-infected animal model are likely to support protective immunity against reinfection, related studies are still in early stages. Based on experience from animal coronaviruses,⁸⁷ including other human coronaviruses,⁸⁸ it is tempting to speculate that reinfection is likely to occur in some people. This is practically possible in immunodeficient individuals with limited capacity to mount proper adaptive immunity and/or those that failed to produce sufficient memory cells.⁸⁹ However, recurrent infection may not be a general feature of COVID-19, as the cases reported so far are not particularly high.^{20,49,50} Further experiments are needed to provide a definitive answer.

Implication in vaccine development

Vaccine aims to stimulate the cellular and humoral immunity of the adaptive immune system to recognize a pathogen and mount sufficient numbers of memory T and B cells as well as long lasting nAbs against the pathogen. Therefore, it is expected to protect the vaccinated individual from severe disease when infected by the specific pathogen the vaccine was designed for. Unlike cancer vaccines, which have also been used for immunotherapy (postphylaxis), vaccines against infectious disease-related pathogens are usually administered as a means of prevention (prephylaxis).⁹⁰

Vaccine candidates currently developed against SARS-CoV-2 come in different forms, which include the classic whole-virus vaccines (both inactivated and live attenuated vaccines), genetic vaccines in the form of DNA or RNA vaccines, viral vector vaccines, and protein-based vaccines in the forms of virus-like particles (VLPs) or subunit vaccines.⁹¹ In order for a vaccine to be effective, several immunological factors have to be taken into account.

COVID-19 vaccine should induce protective and durable immunity

An effective vaccine ideally aims to prevent vaccinated individuals from target pathogens by stimulating the memory mechanism of protective humoral and cellular immunity; this protection should be long lasting. However, natural infection by coronaviruses does not usually result in long-lasting

protective immunity, and antibodies wane within a few years or even months.⁹² A similar phenomenon has been observed with immunity to SARS-CoV-2. Short duration of immune protection may allow reinfection by the same virus once the stimulated protective memory component of the adaptive immune system has depleted. Impaired immunity and disease severity in COVID-19 result from viral interference with type I interferon (IFN-I) synthesis.⁹³ IFN-I plays a critical role in the activation and maturation of the adaptive immune system,⁹⁴ and inhibition of IFN-I synthesis hinders the activation and maturation of B cells,^{95–97} dendritic cells (DC),⁹⁸ and T cells.⁹⁹

Consequently, vaccine design should not only mimic natural SARS-CoV-2 infection through the administration of appropriate antigens, but vaccine components should also be engineered to stimulate stronger immune response both in magnitude and durability than what is achievable by natural infection. One possible way is by enhancing the innate immune response, especially by pathways regulating the adaptive immune response, such as, but not limited to, IFN-I synthesis.¹⁰⁰

Vaccine adjuvants have been shown to increase the durability of the immune response elicited by a SARS-CoV whole virus vaccine candidate. Different adjuvants including alum, CpG, Adva, and delta-inulin-based polysaccharide increased serum nAb titers and reduced lung virus titers in mice.⁹² The use of alum in a yeast recombinant Hepatitis B (HBV), for example, had an increasing seroprotective effect in adults ≥40 years old.¹⁰¹

Several antigen delivery systems currently used in the development of SARS-CoV-2 vaccine candidates may also enhance the innate and adaptive immune responses. Liposomes, for example, are synthetic phospholipid bilayers mimicking the plasma membrane of living cells. Liposomal delivery systems may be engineered to constitute phospholipid types, which act as intracellular messengers in modulating innate and adaptive immunity. Liposome surfaces can also be decorated with adjuvants or PAMP entities such as polysaccharides (LPS) to stimulate the corresponding pattern recognition receptors (PRR), such as the toll-like receptors (TLR), which in return induces IFN-I expression.¹⁰²

DNA vaccine encoding SARS-CoV-2 protein utilizes the adenoviral vector (Ad) as a delivery system.¹⁰³ The Ad vector, for more or less than a decade, has been shown to activate or transduce DCs, macrophages, and natural killer (NK) cells. Ad vector also plays a role in IFN-I induction, which is important for the efficacy of Ad-based vaccines, and has a critical role in antigen presenting cell (APC) maturation and proinflammatory cytokine induction and regulation. IFN-I induction by Ad vector is mediated by the TLR9 and RIG-I receptors of the innate immune system.¹⁰⁴

VLP and other protein-based nanocages (PNC) are composed of an assembly of monomeric subunit proteins such as viral capsid proteins, or other protein subunits such as the small heat shock protein, forming a cage-like nanostructure.¹⁰⁵ VLP-based vaccines are either composed of native SARS-CoV-2 capsid proteins⁹¹ or a heterologous VLP platform presenting the SARS-CoV-2 spike protein on the surface.¹⁰⁶ The assembly of monomeric subunit proteins to form the VLP or PNC structure is advantageous in immune

response enhancement in several ways. Firstly, the structural assembly provides a repetitive antigen presentation motif with high spatio-geometric density, which is known to enhance the immunogenicity and responsiveness of B cells.¹⁰⁷ Secondly, the repetitive nature of VLP and NPC assembly provides a PAMP motif allowing recognition by PRR, such as the TLR, and leading to IFN-I synthesis.¹⁰⁸ Thirdly, VLPs can activate DCs and enter APCs, including DC, to present antigens via the MHC Class I and MHC Class II pathways and therefore mediate the cytotoxic T cell and helper T cell immune response, respectively.^{108–111} Small heat shock protein nanocages, without any additional antigen, induced the formation of bronchus-associated lymphoid tissue (iBALT) in the lung when inoculated intranasally and showed protection against lethal challenge induced by viral respiratory pathogens, including SARS-CoV.¹¹² A similar phenomenon was also observed when the inoculation with antigen-free papaya mosaic virus (PapMV) VLP showed protection against influenza or *Streptococcus pneumoniae* challenge in mice.¹¹³ The observed protective immune response was mediated by the innate immunity, most likely involving neutrophils and CD11c+ cells.¹¹³

HPV vaccines are made of virus-like particles (VLP) assembled from recombinant HPV coat proteins (L1). The fact that the vaccine is highly immunogenic and is very effective in preventing infection long-term^{114,115} may be due to the immunogenic properties of the close geometrical spacing of the repetitive antigen presented on the VLP.^{90,116} However, the L1-based VLP vaccine does not have an effective post-exposure therapeutic effect to clear HPV infected tumour cells. HPV-associated tumour progression is the result of HPV DNA integration into the host cell DNA, inactivating some HPV genes, including L1, and upregulating HPV E6 and E7 protein expression, which play a role in host cell transformation into tumour cells. The activation of L1 protein hinders their presentation via the MHC Class I and MHC Class II pathways on the infected host cells, and as a consequence, also the L1-based vaccine stimulated cytotoxic and helper T cell clearance.¹¹⁷ To circumvent this, efforts have been made to develop vaccines using the continuously expressed E6 and E7 as antigens.¹¹⁸ In the context of VLP-based vaccine, the encapsidation technology¹¹⁹ may be useful to ensure entry of the E6 and E7 antigens into APCs and to prevent extracellular antibody neutralisation.

The ability of pathogenic viral proteins to interfere with the innate and adaptive immune response in the form of altered gene expression conferring tolerance or other forms of immune evasion has been observed.^{120,121} For the coronaviruses, several structural and non-structural proteins which play a role in pathways leading to IFN-I and III expression have been identified,¹²² including nsp16,¹²³ membrane,^{124,125} nucleocapsid,¹²⁶ amongst others. Some of these immune evasive proteins have a high homology with SARS-CoV-2 proteins.¹²⁷ Recent literature suggested that the non-structural protein¹²⁸ and the ORF3 protein¹²⁹ of the virus may interfere with the immune response. Therefore, the inclusion of these proteins in a vaccine design should be evaluated.

In addition, a strong antibody response and T cell response to ensure protective and enduring immunity requires

prolonged antigen exposure to B and T cells in a dose-escalating manner.^{130,131} Accordingly, COVID-19 vaccine engineering may utilize a controlled-release delivery technology, and repeated vaccination in a dose-escalating manner may also provide a similar benefit.

COVID-19 vaccine should stimulate humoral and cellular immunity

Protective immunity against a pathogen usually depends on the availability of nAbs and effector T cells at the time of infection.¹³² To induce protective immunity, vaccines are primarily designed to induce nAb expression by the adaptive immune system and the corresponding formation of memory B cells that can be rapidly activated after antigen reexposure.¹³⁰ The function of nAbs is to prevent viral interaction with the host, and hence, much work has been done in vaccine development to evaluate the generation of antibodies capable of binding to the receptor-binding domain (RBD) of SARS-CoV-2 spike protein. The RBD plays a role in viral binding to the angiotensin-converting enzyme-2 (ACE-2) receptor in a variety of host cell types.^{133,134} Antibodies that specifically bind to this domain prevent the entry of the SARS-CoV-2 into the host cells and therefore exhibit neutralizing properties.¹³⁵

The generation of memory B cells and nondividing bone marrow-resident plasma cells for long-term antibody production protective for the next bout of infection requires antigenic multivalency and antigenic threshold. Repetitive antigen and a longer duration of antigen exposure during natural infection or vaccination may increase the activation of germinal centers and thus the number of activated T and B cells, and hence, increased amount of long-lived plasma cells as well as memory B and T cells.^{90,116}

A recent study also pointed out that SARS-CoV-2 could enter T cells with an alternative receptor, possibly CD147, through a yet unidentified region on its spike protein.¹³⁶ Hence, identification of this binding region and its corresponding nAbs would be important in preventing T cell infection and consequently T cell response impairment.

Stimulation of cellular immunity through corresponding T cell activation should be targeted concurrently with the generation of antibody-related responses against SARS-CoV-2. Single-cell RNA sequencing analysis of the bronchoalveolar lavage fluid (BALF) from COVID-19 patients demonstrated the crucial role of cytotoxic T cells (Tc) during recovery from COVID-19.¹³⁷ Clonally expanded effector Tc of homogenous transcriptional profile with tissue-resident characteristics and the upregulation of genes associated with activation, migration, and the cytokine pathway are the hallmarks of Tc population in the BALF from moderate COVID-19 patients. However, severe/critical patients display an ensemble of highly proliferative T cells with heterogeneous transcriptional profiles and upregulation of genes associated with translation initiation, cell homeostasis, and nucleoside metabolic pathway.¹³⁷ Moderate COVID-19 cases are also marked by elevated T cell recruitment into the lung, whereas chemokine recruitment favors inflammatory monocytes and neutrophils into the lungs in severe/critical cases.¹³⁷ Analysis of Tc responses in COVID-19 patients revealed that the majority of convalescent

individuals generate Tc responses, responding to Tc epitopes from the spike (26%), membrane (22%), nonstructural protein (15%), nucleocapsid protein (12%), ORF8 (10%), and ORF3a (7%).¹³⁸ Consequently, in addition to the spike protein, a vaccine designed to concurrently target Tc response should also consider the delivery of nonspike SARS-CoV-2 proteins related to Tc responses.

Analysis of helper T cell response revealed their important role in patient recovery from COVID-19, with T helper 1 (Th1) showing the prominent response.^{138,139} Unlike the non-spike Th response, spike-specific Th response correlated well with the magnitude of anti-RBD spike protein IgG and IgA responses. Th responses were mostly directed towards the spike (27%), membrane (21%), and nucleocapsid protein (11%) of SARS-CoV-2.¹³⁸ As Th response plays a significant role in B cell maturation to generate nAbs, the raising of Th response should be considered in vaccine design for COVID-19.

In order for a vaccine to generate a T cell response, T cell-associated antigens need to be delivered into the host cellular compartment to stimulate the cellular immune response. Moreover, it would be advantageous to minimize the generation of antibodies against T-cell associated antigens, which may prevent the corresponding antigens from reaching the host cellular compartment during natural infection. This may be achieved by encapsulating the T cell-associated antigen vaccine component inside the relevant delivery system (such as in the use of VLPs¹¹⁹ and liposomes) or engineer their expression inside the host cell (such as in the use of RNA and DNA vaccines). Moreover, T cell epitopes are associated with HLA, and therefore, T-cell based vaccine candidates should be designed to encompass the majority of the global population.¹⁴⁰

Immunity works by exerting both the humoral (antibody-based) and cellular (T cell-based) immunity. In vaccine development, the correlate of protection should be determined and the presence of immune-related adverse events should be tested and avoided.

The RNA polymerase of RNA vaccines such as SARS-CoV-2 are known to be error-prone, and therefore, may introduce frequent mutations.¹⁴¹ Although the mutational rate of SARS-CoV-2 is lower than influenza,¹⁴² a future mutation altering the conformational structure of nAb recognition sites of the SARS-CoV-2 protein may result in possible reinfection. Furthermore, a vaccine designed to stimulate cross-protective immunity would be a good strategy to prevent or reduce the severity of disease from other possible pandemic coronavirus strains in the future.¹⁴³

A cross-protective vaccine may utilize conserved T and B cell epitopes to stimulate cross-protective immunity.^{144,145} Conserved or partially conserved T cell epitopes are commonly easier to identify and engineer than conserved B cell epitopes due to the conformation-dependent nature of B cell epitopes.^{146,147}

A natural T cell response may not prevent infection but may reduce the severity of the disease.¹⁴⁸ Therefore, even if reinfection occurs in a state of reduced antibody levels, there is a possibility that pre-existing T cell immunity may prevent clinically severe disease. Moreover, conserved or partially

conserved T cell epitopes from other circulating coronaviruses including those that cause the common cold may have provided a broadly cross-protective T cell immunity against SARS-CoV-2 in unexposed individuals. This may explain why a significant number of SARS-CoV-2 positively tested individuals are asymptomatic/presymptomatic or show only mild symptoms or moderate disease in SARS-CoV-2.¹⁴⁹ However, it has to be kept in mind that during the period of natural SARS-CoV-2 infection, with or without symptoms, a patient may still transmit the virus to other people in their vicinity, who may be at higher risk in contracting a more severe disease.

A T cell-based SARS-CoV-2 vaccine utilizing conserved T cell epitopes shared amongst coronavirus strains (those causing common cold and also the more severe disease such as SARS and MERS), may not only confer some protection against SARS-CoV-2 in terms of reducing disease severity but may also protect from other potential coronavirus pandemic strains in the future. These T cell-based vaccines can be used as a pre-pandemic vaccine or administered in the early phase of a pandemic to reduce health and disease burden before an antibody-based vaccine becomes available.^{30,150}

Studies on Th responses have identified SARS-CoV-2 epitopes that have also been found previously circulating human coronaviruses,^{149,151} which may play a role in the immune protection of unexposed and recovered patients of COVID-19. Conserved antibody recognition site¹⁵² and cross-protective Th epitopes on the SARS-CoV-2 S protein^{149,151} have been identified and their use in vaccine design may help from severe re-infection of SARS-CoV-2 or infection from a future coronavirus related pandemic.

Nonetheless, T cell immunity may only protect to a certain extent, and therefore, a combination with antibody-based immunity will be the best approach in vaccine development. Moreover, there are additional factors to consider, such as ruling out immune-related adverse events (irAE)¹⁵³ and determining the correlates of protection of both T cell-based and antibody-based immunity.⁹⁰ Epidemiologically, a current estimate of 70% vaccinated individual in the population will result in herd immunity and thus reduce the transmission of the virus in the population to a negligible level.^{154–156}

Conclusion and future perspectives

The transmission of infectious viral particles from one host to another usually triggers vigorous host immune responses leading to the clearance of viral particles. However, in some cases, the virus cannot be cleared off, resulting in either the death of the infected host or the emergence of persistent infection. In another scenario, host immunity can remove the viral particles in the primary infection but fails to protect the host from reinfection. There is still limited knowledge about whether the reported reinfection cases of COVID-19 were accurately described as reinfection or whether they were simply due to the failure of currently available methods to detect the low copy number of SARS-CoV-2 in certain patients, which led to a relapse. To improve our understanding on this issue, suitable model organisms to investigate the nature of reinfection and elucidate its possible mechanistic basis are urgently required.

Due to its insidious life cycle in the host cells, SARS-CoV-2 can achieve continuous transmission among humans. For susceptible hosts such as immunodeficient, older people, and those with associated comorbidities, SARS-CoV-2 primary infection increases the risk of death. To avoid such outcome, effective and safe vaccination to achieve herd immunity in the given population is the best option to adopt. However, with the challenges associated with development of an effective COVID-19 vaccine and the probability of reinfection by SARS-CoV-2, if that is truly possible, the risk of death of susceptible hosts may persist. In such conditions, avoidance of reinfection is the only available option, however difficult it might be to achieve.

5 Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

This review received no external funding.

Human subjects approval statement

Not required.

19 ORCID

Ali A Rabaan <http://orcid.org/0000-0006-6774-9847>

Kuldeep Dhama <http://orcid.org/0000-0001-7469-4752>

Harapan Harapan <http://orcid.org/0000-0001-7630-8413>

References

- Rodriguez-Morales AJ, Cardona-Ospina JA, Gutierrez-Ocampo E, Villamizar-Pena R, Holguin-Rivera Y, Escalera-Antezana JP, Alvarado-Arnez LE, Bonilla-Aldana DK, Franco-Paredes C, Henao-Martinez AF, et al. Clinical, laboratory and imaging features of COVID-19: a systematic review and meta-analysis. *Travel Med Infect Dis.* 2020;34:101623. doi:10.1016/j.tmaid.2020.101623.
- Harapan H, Itoh N, Yufika A, Winardi W, Keamg S, Te H, Megawati D, Hayati Z, Wagner AL, Mudatsir M, et al. Coronavirus disease 2019 (COVID-19): a literature review. *J Infect Public Health.* 2020;13(5):667–73. doi:10.1016/j.jiph.2020.03.019.
- Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis.* 2020;20(5):533–34. doi:10.1016/S1473-3099(20)30120-1.
- Chan JF-W, Zhang AJ, Yuan S, Poon VK-M, Chan CC-S, Lee AC-Y, Chan W-M, Fan Z, Tsoi H-W, Wen L, et al. Simulation of the clinical and pathological manifestations of Coronavirus Disease 2019 (COVID-19) in golden Syrian hamster model: implications for disease pathogenesis and transmissibility. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa325.
- Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si H-R, Zhu Y, Li B, Huang C-L, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature.* 2020;579(7798):270–73. doi:10.1038/s41586-020-2012-7.
- Lam TT, Shum MH, Zhu HC, Tong YG, Ni XB, Liao YS, Tong Y-G, Shi Y-X, Ni X-B, Liao Y-S, et al. Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. *Nature.* 2020;583(7815):282–85. doi:10.1038/s41586-020-2169-0.

7. Zhang T, Wu Q, Zhang Z. Probable pangolin origin of SARS-CoV-2 associated with the COVID-19 outbreak. *Curr Biol.* 2020;30(8):1578. doi:10.1016/j.cub.2020.03.063.
8. Keam S, Megawati D, Patel S, Tiwari R, Dhama K, Harapan H. Immunopathology and immunotherapeutic strategies in SARS-CoV-2 infection. *Rev Medical Virol.* 2020;30(5). doi:10.1002/rmv.2123.
9. Frediansyah A, Nainu F, Dhamad K, Mudatsir M, Harapan H. Remdesivir and its antiviral activity against COVID-19: a systematic review. *Clin Epidemiol Glob Health.* 2020. (In press) doi:10.1016/j.cegh.2020.07.011
10. Frediansyah A, Tiwari R, Sharun K, Dhama K, Harapan H. Antivirals for COVID-19: a critical review. *Clin Epidemiol Global Health.* 2020. doi:10.1016/j.cegh.2020.07.006.
11. Sharun K, Tiwari R, Iqbal Yatoo M, Patel SK, Natesan S, Dhama J, Malik YS, Harapan H, Singh RK, Dhama K, et al. Antibody-based immunotherapeutics and use of convalescent plasma to counter COVID-19: advances and prospects. *Expert Opin Biol Ther.* 2020;20(9):1033–46. doi:10.1080/14712598.2020.1796963.
12. Bao L, Deng W, Gao H, Xiao C, Liu J, Xue J, Lv Q, Liu J, Yu P, Xu Y, et al. Reinfection could not occur in SARS-CoV-2 infected rhesus macaques. *BioRxiv.* 2020. doi:10.1101/2020.03.13.990226.
13. Le T T, Andreadakis Z, Kumar A, Gomez Roman R, Tollefsen S, Saville M, Mayhew S. The COVID-19 vaccine development landscape. *Nat Rev Drug Discov.* 2020;19:305–306. doi:10.1038/d41573-020-00073-5.
14. Lurie N, Saville M, Hatchett R, Halton J. Developing Covid-19 vaccines at pandemic speed. *N Engl J Med.* 2020;382(21):1969–73. (In press). doi:10.1056/NEJMp2005630.
15. Harapan H, Wagner AL, Yufika A, Winardi W, Anwar S, Gan AK, Setiawan AM, Rajamoorthy Y, Sofyan H, Mudatsir M, et al. Acceptance of a COVID-19 vaccine in Southeast Asia: a cross-sectional study in Indonesia. *Front Public Health.* 2020;8:381. doi:10.3389/fpubh.2020.00381.
16. Harapan H, Wagner A, Yufika A, Winardi W, Anwar S, Gan A, Setiawan AM, Rajamoorthy Y, Sofyan H, Vo TQ, et al. Willingness to pay for a COVID-19 vaccine in Indonesia. *Hum Vacc Immunother.* 2020;1–7. (Submitted). doi:10.1080/21645515.2020.1819741.
17. Smith J South Korea reports more recovered coronavirus patients testing positive again. April 13, 2020.
18. Korean CDC. Findings from investigation and analysis of re-positive cases. 19 May 2020.
19. Ni L, Ye F, Cheng M-L, Feng Y, Deng Y-Q, Zhao H, Wei P, Ge J, Gou M, Li X, et al. Detection of SARS-CoV-2-specific humoral and cellular immunity in COVID-19 convalescent individuals. *Immunity.* 2020;52(6):971–977.e3. doi:10.1016/j.immuni.2020.04.023.
20. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, Xie C, Ma K, Shang K, Wang W, et al. Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis.* 2020;71(15):762–68. doi:10.1093/cid/ciaa248.
21. Lan L, Xu D, Ye G, Xia C, Wang S, Li Y, Xu H. Positive RT-PCR test results in patients recovered from COVID-19. *Jama.* 2020;323(15):1502–03. doi:10.1001/jama.2020.2783.
22. Da Guan W, Mok CKP, Chen ZL, Feng LQ, Li ZT, Huang JC, Ke CW, Deng X, Ling Y, Wu SG. Characteristics of traveler with Middle East respiratory syndrome, China, 2015. *Emerg Infect Dis.* 2015;21(12):2278. doi:10.3201/eid2112.151232.
23. Liu Z-Y, Li T-S, Wang Z, Xu Z-J, Wang H-L, Yu Y, Du T-K, Bai Y, Qiu Z-F, Lu W, et al. Clinical features and therapy of 106 cases of severe acute respiratory syndrome. *Zhonghua Nei Ke Za Zhi.* 2003;42:373–77.
24. MO H, ZENG G, REN X, LI H, KE C, TAN Y, CAI C, LAI K, CHEN R, CHAN-YEUNG M, et al. Longitudinal profile of antibodies against SARS-CoV-2 in SARS patients and their clinical significance. *Respirology.* 2006;11(1):49–53. doi:10.1111/j.1440-1843.2006.00783.x.
25. Alagaili AN, Briese T, Mishra N, Kapoor V, Sameroff SC, Burbelo PD, de Wit E, Munster VJ, Hensley LE, Zalmout IS, et al. Middle East respiratory syndrome coronavirus infection in dromedary camels in Saudi Arabia. *MBio.* 2014;5(2):e00884–14. doi:10.1128/mBio.01002-14.
26. Hemida MG, Chu DK, Poon LL, Perera RA, Alhammadi MA, Ng H-Y, Siu LY, Guan Y, Alnaeem A, Peiris M, et al. MERS coronavirus in dromedary camel herd, Saudi Arabia. *Emerg Infect Dis.* 2014;20(7):1231. doi:10.3201/eid2007.140571.
27. Roberts A, Vogel L, Guarner J, Hayes N, Murphy B, Zaki S, Subarao K. Severe acute respiratory syndrome coronavirus infection of golden Syrian hamsters. *J Virol.* 2005;79(1):503–11. doi:10.1128/JVI.79.1.503-511.2005.
28. Subbarao K, McAuliffe J, Vogel L, Fahle G, Fischer S, Tatti K, Packard M, Shieh W-J, Zaki S, Murphy B, et al. Prior infection and passive transfer of neutralizing antibody prevent replication of severe acute respiratory syndrome coronavirus in the respiratory tract of mice. *J Virol.* 2004;78(7):3572–77. doi:10.1128/JVI.78.7.3572-3577.2004.
29. McAuliffe J, Vogel L, Roberts A, Fahle G, Fischer S, Shieh W-J, Butler E, Zaki S, St. Claire M, Murphy B, et al. Replication of SARS coronavirus administered into the respiratory tract of African Green, rhesus and cynomolgus monkeys. *Virology.* 2004;330(1):8–15. doi:10.1016/j.virol.2004.09.030.
30. Woodland DL. Progress towards a SARS vaccine. *Viral Immunol.* 2010;23(5):455. doi:10.1089/vim.2010.ed.23.5.
31. Excler JL, Delvecchio CJ, Wiley RE, Williams M, Yoon IK, Modjarrad K, Boujelal M, Moorthy VS, Hersi AS, Kim JH, et al. Toward developing a preventive MERS-CoV vaccine-report from a workshop organized by the Saudi Arabia ministry of health and the international vaccine institute, Riyadh, Saudi Arabia, November 14–15, 2015. *Emerg Infect Dis.* 2016;22(8):e160229. doi:10.3201/eid2208.160229.
32. Chen D, Xu W, Lei Z, Huang Z, Liu J, Gao Z, Peng L. Recurrence of positive SARS-CoV-2 RNA in COVID-19: a case report. *Int J Infect Dis.* 2020;93:297–99. doi:10.1016/j.ijid.2020.03.003.
33. Lippi G, Simundic A-M, Plebani M. Potential preanalytical and analytical vulnerabilities in the laboratory diagnosis of coronavirus disease 2019 (COVID-19). *Clin Chem Lab Med.* 2020;58(7):1070–1076. doi:10.1515/cclm-2020-0285.
34. Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, Xing F, Liu J, Yip CCY, Poon RWS, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet.* 2020;395(10223):514–23. doi:10.1016/S0140-6736(20)30154-9.
35. Menggi HYT, Wang S, Chen S, Zhou W, Chen D, Zhou L, Wang M, Zhao Y, Zeng W, Huang Q, et al. Clinical characteristics of laboratory confirmed positive cases of SARS-CoV-2 infection in Wuhan, China: a retrospective single center analysis. *Travel Med Infect Dis.* 2020;36:101606. doi:10.1016/j.tmaid.2020.101606.2020.
36. Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for typical 2019-nCoV pneumonia: relationship to negative RT-PCR testing. *Radiology.* 2020;296(2):E41–E45. doi:10.1148/radiol.202003043.
37. CDC. Coronavirus Disease 2019 (COVID-19). <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>, 2020.
38. Lippi G, Simundic A-M, Plebani M. Potential preanalytical and analytical vulnerabilities in the laboratory diagnosis of coronavirus disease 2019 (COVID-19). *Clin Chem Lab Med.* 2020;58(7):1070. doi:10.1515/cclm-2020-0285.
39. Pulendran B, Oh JZ, Nakaya HI, Ravindran R, Kazmin DA. Immunity to viruses: learning from successful human vaccines. *Immunity Rev.* 2013;255(1):243–55. doi:10.1111/immr.12099.
40. Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LFP. The trinity of COVID-19: immunity, inflammation and intervention. *Nat Rev Immunol.* 2020;20(6):363–74. doi:10.1038/s41577-020-03111-8.
41. Vabret N, Britton GJ, Gruber C, Hegde S, Kim J, Kuksin M, Levantovsky R, Malle L, Moreira A, Park MD, et al. Immunology of COVID-19: current state of the science. *Immunity.* 2020;52(6):910–41. doi:10.1016/j.immuni.2020.05.002.
42. Chaplin DD. Overview of the immune response. *J Allergy Clin Immunol.* 2010;125(2):S3–S23. doi:10.1016/j.jaci.2009.12.980.
43. Petrosillo N, Viceconte G, Ergonul O, Ippolito G, Petersen E. COVID-19, SARS and MERS: are they closely related? *Clin Microbiol Infect.* 2020;S1198–743X:30171–73.

44. Wang L, Wang Y, Ye D, Liu Q. Review of the 2019 novel coronavirus (SARS-CoV-2) based on current evidence. *Int J Antimicrob Agents.* 2020;55(6):105948. doi:10.1016/j.ijantimicag.2020.105948.
45. Aoshi T, Koyama S, Kobiyama K, Akira S, Ishii KJ. Innate and adaptive immune responses to viral infection and vaccination. *Curr Opin Virol.* 2011;1(4):226–32. doi:10.1016/j.coviro.2011.07.002.
46. Rouse BT, Sehrawat S. Immunity and immunopathology to viruses: what decides the outcome? *Nat Rev Immunol.* 2010;10(7):514–26. doi:10.1038/nri2802.
47. Ju B, Zhang Q, Ge X, Wang R, Yu J, Shan S, Zhou B, Song S, Tang X, Yu J, et al. Potent human neutralizing antibodies elicited by SARS-CoV-2 infection. *bioRxiv.* 2020;990770.
48. Viroj JBW. Hemorrhagic problem among the patients with COVID-19: clinical summary of 41 Thai infected patients. *J Sage: Clin Appl Thrombosis/Hemostasis.* 2020;26:1076029620918308. doi:10.1177/1076029620918308.
49. Peng J, Wang M, Zhang G, Lu E. Seven discharged patients turning positive again for SARS-CoV-2 on quantitative RT-PCR. *Am J Infect Control.* 2020;S0196–6553:30198–X.
50. Tang X, Zhao S, He D, Yang L, Wang MH, Li Y, Mei S, Zou X. Positive RT-PCR tests among discharged COVID-19 patients in Shenzhen, China. *Infect Control Hosp Epidemiol.* 2020;1–2. doi:10.1017/ice.2020.1232
51. Chen G, Wu D, Guo Y, Cao Y, Huang D, Wang H, Wang T, Zhang X, Chen H, Yu H, et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest.* 2020;130(5):2620–29. doi:10.1172/JCI137244.
52. Li Y, Hu Y, Yu Y, Zhang X, Li B, Wu J, Li J, Wu Y, Xia X, Tang H, et al. Positive result of Sars-Cov-2 in faeces and sputum from discharged patient with COVID-19 in Yiwu, China. *J Med Virol.* 2020;92:1938–1947. doi:10.1002/jmv.259050.
53. Zheng H-Y, Zhang M, Yang C-X, Zhang N, Wang X-C, Yang X-P, Dong X-Q, Zheng Y-T. Elevated exhaustion levels and reduced functional diversity of T cells in peripheral blood may predict severe progression in COVID-19 patients. *Cell Mol Immunol.* 2020;17(5):541–43. doi:10.1038/s41423-020-0401-3.
54. Tan L, Wang Q, Zhang D, Ding J, Huang Q, Tang Y-Q, Wang Q, Miao H. Lymphopenia predicts disease severity of COVID-19: a descriptive and predictive study. *Signal Transduction Targeted Ther.* 2020;5(1):33. doi:10.1038/s41392-020-0148-4.
55. Chen X, Zhao B, Qu Y, Chen Y, Xiong J, Feng Y, Men D, Huang Q, Liu Y, Yang B, et al. Detectable serum SARS-CoV-2 viral load (RNAemia) is closely correlated with drastically elevated interleukin 6 (IL-6) level in critically ill COVID-19 patients. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa449.
56. Diao B, Wang C, Tan Y, Chen X, Liu Y, Ning L, Chen L, Li M, Liu Y, Wang G, et al. Reduction and functional exhaustion of T cells in patients with coronavirus disease 2019 (COVID-19). *medRxiv.* 2020;20024364. doi:10.1101/2020.02.18.20024364.
57. Wan S, Yi Q, Fan S, Lv J, Zhang X, Guo L, Lang C, Xiao Q, Xiao K, Yi Z, et al. Characteristics of lymphocyte subsets and cytokines in peripheral blood of 123 hospitalized patients with 2019 novel coronavirus pneumonia (NCP). *medRxiv.* 2020 Feb 10. 20021832. doi:10.1101/2020.02.10.20021832.
58. Yang M. Cell pyroptosis, a potential pathogenic mechanism of 2019-nCoV infection. *SSRN.* 2020. doi:10.2139/ssrn.3527420.
59. Biggs JR, Zhang D-E. Chapter 16 - molecular basis of lymphoid and myeloid diseases. In: Coleman WB, Tsongalis GJ, editors. *Molecular pathology.* Seconded. Academic Press;2018. p.299–328.
60. Liao M, Liu Y, Yuan J, Wen Y, Xu G, Zhao J, Cheng L, Li J, Wang X, Wang F, et al. Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat Med.* 2020;26(6):842–44. doi:10.1038/s41591-020-0901-9.
61. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, Liu S, Zhao P, Liu H, Zhu L, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med.* 2020;8(4):420–22. doi:10.1016/S2213-2600(20)30076-X.
62. Newton K, Dixit VM. Signaling in innate immunity and inflammation. *Cold Spring Harb Perspect Biol.* 2012;4(3):a006049. doi:10.1101/cshperspect.a006049.
63. Amarante-Mendes GP, Adjemian S, Branco LM, Zanetti LC, Weinlich R, Bortoluci KR. Pattern recognition receptors and the host cell death molecular machinery. *Front Immunol.* 2018;9:2379. doi:10.3389/fimmu.2018.02379.
64. Tang D, Kang R, Coyne CB, Zeh HJ, Lotze MT. PAMPs and DAMPs: signal 0s that spur autophagy and immunity. *Immunol Rev.* 2012;249:158–75.
65. Vardhana SA, Wolchok JD. The many faces of the anti-COVID immune response. *J Exp Med.* 2020;217(6):e20200678. doi:10.1084/jem.20200678.
66. Wherry EJ. T cell exhaustion. *Nat Immunol.* 2011;12(6):492–99. doi:10.1038/ni.2035.
67. Saeidi A, Zandi K, Cheok YY, Saeidi H, Wong WF, Lee CYQ, Cheong HC, Yong YK, Larsson M, Shankar EM. T-cell exhaustion in chronic infections: reversing the state of exhaustion and reinvigorating optimal protective immune responses. *Front Immunol.* 2018;9:2569. doi:10.3389/fimmu.2018.02569.
68. Callow KA, Parry HF, Sergeant M, Tyrrell DA. The time course of the immune response to experimental coronavirus infection of man. *Epidemiol Infect.* 1990;105(2):435–46. doi:10.1017/S0950268800048019.
69. Wu L-P, Wang N-C, Chang Y-H, Tian X-Y, Na D-Y, Zhang L-Y, Zheng L, Lan T, Wang L-F, Liang G-D, et al. Duration of antibody responses after severe acute respiratory syndrome. *Emerg Infect Dis.* 2007;13(10):1562–64. doi:10.3201/eid1310.070576.
70. Li CK-F, Wu H, Yan H, Ma S, Wang L, Zhang M, Tang X, Temperton NJ, Weiss RA, Brenchley JM, et al. T cell responses to whole SARS coronavirus in humans. *J Immunol.* 2008;181(8):5490–500. doi:10.4049/jimmunol.181.8.5490.
71. Park WB, Perera RAMP, Choe PG, Lau EHY, Choi SJ, Chun JY, Oh HS, Song K-H, Bang JH, Kim ES, et al. Kinetics of serologic responses to MERS coronavirus infection in humans, South Korea. *Emerg Infect Dis.* 2015;21(12):2186–89. doi:10.3201/eid2112.151421.
72. Ko J-H, Müller MA, Seok H, Park GE, Lee JY, Cho SY, Ha YE, Baek JY, Kim SH, Kang J-M, et al. Serologic responses of 42 MERS-coronavirus-infected patients according to the disease severity. *Diagn Microbiol Infect Dis.* 2017;89(2):106–11. doi:10.1016/j.diagmicrobio.2017.07.006.
73. Huang AT, Garcia-Carreras B, Hitchings MDT, Yang B, Katselnick L, Rattigan SM, Borgert B, Moreno C, Solomon BD, Rodriguez-Barrquer I, et al. A systematic review of antibody mediated immunity to coronaviruses: antibody kinetics, correlates of protection, and association of antibody responses with severity of disease. *medRxiv.* 2020;20065771. doi:10.1101/2020.04.14.20065771.
74. Cheng MP, Papenburg J, Desjardins M, Kanjilal S, Quach C, Libman M, Dittrich S, Yansouni CP. Diagnostic testing for severe acute respiratory syndrome-related coronavirus-2. *Ann Intern Med.* 2020;M20-1301. doi:10.7326/M20-1301.
75. Mohammadi A, Esmaeilzadeh E, Li Y, Bosch RJ, Li JZ. SARS-CoV-2 detection in different respiratory sites: a systematic review and meta-analysis. *EBioMedicine.* 2020;59:102903. doi:10.1016/j.ebiom.2020.102903.
76. Tang Y-W, Schmitz JE, Persing DH, Stratton CW, McAdam AJ. Laboratory diagnosis of COVID-19: current issues and challenges. *J Clin Microbiol.* 2020;58(6):e00512–20. doi:10.1128/JCM.00512-20.
77. Barr JN, Fearn R. How RNA viruses maintain their genome integrity. *J Gen Virol.* 2010;91(6):1373–87. doi:10.1099/vir.0.020818-0.
78. Khailany RA, Safdar M, Ozaslan M. Genomic characterization of a novel SARS-CoV-2. *Gene Rep.* 2020;19:100682. doi:10.1016/j.genrep.2020.100682.
79. Sanjuán R, Domingo-Calap P. Mechanisms of viral mutation. *Cell Mol Life Sci.* 2016;73(23):4433–48. doi:10.1007/s00018-016-2299-6.
80. Kumar S, Nyodu R, Maurya VK, Saxena SK. Host immune response and immunobiology of human SARS-CoV-2 infection. *Coronavirus Dis 2019 (COVID-19).* 2020;43–53.
81. Ortega-Prieto AM, Dorner M. Immune evasion strategies during chronic hepatitis B and C virus infection. *Vaccines (Basel).* 2017;5:24.

82. Guha D, Ayyavoo V. Innate immune evasion strategies by human immunodeficiency virus type 1. *Ism Aids*. 2013;2013:954806. doi:10.1155/2013/954806.
83. Guerra-Assunção JA, Houben RMGJ, Crampin AC, Mzembe T, Mallard K, Coll F, Khan P, Banda L, Chiwaya A, Pereira RPA, et al. Recurrence due to relapse or reinfection with *Mycobacterium tuberculosis*: a whole-genome sequencing approach in a large, population-based cohort with a high HIV infection prevalence and active follow-up. *J Infect Dis*. 2015;211:1154–63. doi:10.1093/infdis/jiu574.
84. Brown JR, Roy S, Tutill H, Williams R, Breuer J. Super-infections and relapses occur in chronic norovirus infections. *J Clin Virol*. 2017;96:44–48. doi:10.1016/j.jcv.2017.09.009.
85. Bao L, Deng W, Gao H, Xiao C, Liu J, Xue J, Lv Q, Liu J, Yu P, Xu Y, et al. Reinfection could not occur in SARS-CoV-2 infected rhesus macaques. *BioRxiv*. 2020. doi:10.1101/2020.03.13.990226.
86. Chandrashekar A, Liu J, Martinot AJ, McMahan K, Mercado NB, Peter L, Tostanoski LH, Yu J, Maliga Z, Nekorchuk M, et al. SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science*. 2020;369(6505):812–817. doi:10.1126/science.abc4776.
87. Saif LJ. Bovine respiratory coronavirus. *Vet Clin North Am Food Anim Pract*. 2010;26(2):349–64. doi:10.1016/j.cvfa.2010.04.005.
88. Monto AS, Lim SK. The tecumseh study of respiratory illness. VI. Frequency of and relationship between outbreaks of coronavirus infection. *J Infect Dis*. 1974;129(3):271–76. doi:10.1093/infdis/129.3.271.
89. Gaspar HB, Goldblatt D. Immunodeficiency syndromes and recurrent infection. *Br J Hosp Med*. 1997;58:565–68.
90. Plotkin SA. Vaccines: past, present and future. *Nat Med*. 2005;11(S4):S5–11. doi:10.1038/nm1209.
91. WHO. Draft landscape of COVID-19 candidate vaccines. Geneva: World Health Organization; 2020.
92. Enjuanes L, Zuniga S, Castano-Rodriguez C, Gutierrez-Alvarez J, Canton J, Sola I. Molecular basis of coronavirus virulence and vaccine development. *Adv Virus Res*. 2016;96:245–86.
93. Blanco-Melo D, Nilsson-Payant BE, Liu WC, Uhl S, Hoagland D, Moller R, Jordan TX, Oishi K, Panis M, Sachs D, et al. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell*. 2020;181(5):1036–45e9. doi:10.1016/j.cell.2020.04.026.
94. Janeway CA Jr., Medzhitov R. Innate immune recognition. *Annu Rev Immunol*. 2002;20(1):197–216. doi:10.1146/annurev.immunol.20.083001.084359.
95. Crampton SP, Voynova E, Bolland S. Innate pathways to B-cell activation and tolerance. *Ann N Y Acad Sci*. 2010;1183(1):58–68. doi:10.1111/j.1749-6632.2009.05123.x.
96. Cyster JG, Allen CDC. B cell responses: cell interaction dynamics and decisions. *Cell*. 2019;177(3):524–40. doi:10.1016/j.cell.2019.03.016.
97. Khodadadi L, Cheng Q, Radbruch A, Hiepe F. The maintenance of memory plasma cells. *Front Immunol*. 2019;10:721.
98. Munz C, Steinman RM, Fujii S. Dendritic cell maturation by innate lymphocytes: coordinated stimulation of innate and adaptive immunity. *J Exp Med*. 2005;202(2):203–07. doi:10.1084/jem.20050810.
99. Shin KS, Jeon I, Kim BS, Kim IK, Park YJ, Koh CH, Song B, Lee J-M, Lim J, Bae E-A, et al. Monocyte-derived dendritic cells dictate the memory differentiation of CD8(+) T cells during acute infection. *Front Immunol*. 2019;10:1887. doi:10.3389/fimmu.2019.01887.
100. Iwasaki A, Medzhitov R. Regulation of adaptive immunity by the innate immune system. *Science*. 2010;327(5963):291–95. doi:10.1126/science.1183021.
101. Gellin BG, Greenberg RN, Hart RH, Bertino, Jr. JS Jr., Stein DH, Deloria MA, Clements-Mann ML. Immunogenicity of two doses of yeast recombinant hepatitis B vaccine in healthy older adults. *J Infect Dis*. 1997;175(6):1494–97. doi:10.1086/516485.
102. Nisini R, Poerio N, Mariotti S, De Santis F, Fraziano M. The multirole of liposomes in therapy and prevention of infectious diseases. *Front Immunol*. 2018;9:155.
103. van Doremalen N, Lambe T, Spencer A, Belij-Rammerstorfer S, Purushotham JN, Port JR, Avanzato V, Bushmaker T, Flaxman A, Ulaszewska M, et al. ChAdOx1 nCoV-19 vaccination prevents SARS-CoV-2 pneumonia in rhesus macaques. *bioRxiv*. 2020 May 13. 093195. doi:10.1101/2020.05.13.093195.
104. Aldhamen YA, Seregin SS, Amalfitano A. Immune recognition of gene transfer vectors: focus on adenovirus as a paradigm. *Front Immunol*. 2011;2:40. doi:10.3389/fimmu.2011.00040.
105. Steinmetz NF, Lim S, Sainsbury F. Protein cages and virus-like particles: from fundamental insight to biomimetic therapeutics. *Biomater Sci*. 2020;8(10):2771–77. doi:10.1039/D0BM00159G.
106. Zha L, Zhao H, Mohsen MO, Hong L, Zhou Y, Yao C, Guo L, Li Z, Chen H, Liu X, et al. Development of a COVID-19 vaccine based on the receptor binding domain displayed on virus-like particles. *bioRxiv*. 2020 May 06. 079830. doi:10.1101/2020.05.06.079830.
107. Bachmann MF, Rohrer UH, Kundig TM, Burki K, Hengartner H, Zinkernagel RM. The influence of antigen organization on B cell responsiveness. *Science*. 1993;262(5138):1448–51. doi:10.1126/science.8248784.
108. Raghunandan R. Virus-like particles: innate immune stimulators. *Expert Rev Vaccines*. 2011;10(4):409–11. doi:10.1586/erv.11.37.
109. Fu C, Jiang A. Dendritic cells and CD8 T cell immunity in tumor microenvironment. *Front Immunol*. 2018;9:3059. doi:10.3389/fimmu.2018.03059.
110. Keller SA, Bauer M, Manolova V, Muntwiler S, Saudan P, Bachmann MF. Cutting edge: limited specialization of dendritic cell subsets for MHC class II-associated presentation of viral particles. *J Immunol*. 2010;184(1):26–29. doi:10.4049/jimmunol.0901540.
111. Rynda-Apple A, Patterson DP, Douglas T. Virus-like particles as antigenic nanomaterials for inducing protective immune responses in the lung. *Nanomedicine*. 2014;9(12):1857–68. doi:10.2217/nmm.14.107.
112. Wiley JA, Richert LE, Swain SD, Harmsen A, Barnard DL, Randall TD, Jutila M, Douglas T, Broomell C, Young M, et al. Inducible bronchus-associated lymphoid tissue elicited by a protein cage nanoparticle enhances protection in mice against diverse respiratory viruses. *PLoS One*. 2009;4(9):e1742. doi:10.1371/journal.pone.0007142.
113. Rynda-Apple A, Dobrinen E, McAlpine M, Read A, Harmsen A, Richert LE, Calverley M, Pallister K, Voyich J, Wiley JA, et al. Virus-like particle-induced protection against MRSA pneumonia is dependent on IL-13 and enhancement of phagocyte function. *Am J Pathol*. 2012;181(1):196–210. doi:10.1016/j.ajpath.2012.03.018.
114. Patel C, Brotherton JML, Pillsbury A, Jayasinghe S, Donovan B, Macartney K, Marshall H. The impact of 10 years of human papillomavirus (HPV) vaccination in Australia: what additional disease burden will a nonavalent vaccine prevent? *Eurosurveillance*. 2018;23(41):1700737. doi:10.2807/1560-7917.ES.2018.23.41.1700737.
115. Schwarz TF, Huang LM, Valencia A, Panzer F, Chiu CH, Decreux A, Poncet S, Karkada N, Folschweiler N, Lin L, et al. A ten-year study of immunogenicity and safety of the AS04-HPV-16/18 vaccine in adolescent girls aged 10–14 years. *Hum Vaccin Immunother*. 2019;15(7–8):1970–79. doi:10.1080/21645515.2019.1625644.
116. Slifka MK, Amanna IJ. Role of multivalency and antigenic threshold in generating protective antibody responses. *Front Immunol*. 2019;10:956.
117. Zur Hausen H. Papillomaviruses and cancer: from basic studies to clinical application. *Nat Rev Cancer*. 2002;2(5):342–50. doi:10.1038/nrc798.
118. Hung CF, Ma B, Monie A, Tsen SW, Wu TC. Therapeutic human papillomavirus vaccines: current clinical trials and future directions. *Expert Opin Biol Ther*. 2008;8:421–39.
119. Dashti NH, Abidin RS, Sainsbury F. Programmable in vitro coencapsulation of guest proteins for intracellular delivery by virus-like particles. *ACS Nano*. 2018;12(5):4615–23. doi:10.1021/acsnano.8b01059.
120. Myoung J, Lee SA, Lee HR. Beyond viral interferon regulatory factors: immune evasion strategies. *J Microbiol Biotechnol*. 2019;29:1873–81.

121. Chen S, Yang C, Zhang W, Mahalingam S, Wang M, Cheng A. Flaviviridae virus nonstructural proteins 5 and 5A mediate viral immune evasion and are promising targets in drug development. *Pharmacol Ther.* 2018;190:1–14. doi:10.1016/j.pharmthera.2018.05.004.
122. Park A, Iwasaki A. Type I and Type III interferons - induction, signaling, evasion, and application to combat COVID-19. *Cell Host Microbe.* 2020;27(6):870–78. doi:10.1016/j.chom.2020.05.008.
123. Menachery VD, Yount BL, Debbink K, Agnihothram S, Gralinski LE, Plante JA, Graham RL, Scobey T, Ge X-Y, Donaldson EF, et al. A SARS-like cluster of circulating bat coronaviruses shows potential for human emergence. *Nat Med.* 2015;21:1508–13. doi:10.1038/nm.3985.
124. Siu KL, Kok KH, Ng MH, Poon VK, Yuen KY, Zheng BJ, Jin D-Y. Severe acute respiratory syndrome coronavirus M protein inhibits type I interferon production by impeding the formation of TRAF3-TANK/TBK1/IKKepsilon complex. *J Biol Chem.* 2009;284(24):16202–09. doi:10.1074/jbc.M109.008227.
125. Yang Y, Zhang L, Geng H, Deng Y, Huang B, Guo Y, Zhao Z, Tan W. The structural and accessory proteins M, ORF 4a, ORF 4b, and ORF 5 of Middle East respiratory syndrome coronavirus (MERS-CoV) are potent interferon antagonists. *Protein Cell.* 2013;4(12):951–61. doi:10.1007/s13238-013-3096-8.
126. Kopecky-Bromberg SA, Martinez-Sobrido L, Frieman M, Baric RA, Palese P. Severe acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and nucleocapsid proteins function as interferon antagonists. *J Virol.* 2007;81(2):548–57. doi:10.1128/JVI.01782-06.
127. Lokugamage KG, Hage A, Schindewolf C, Rajsbaum R, Menachery VD. SARS-CoV-2 is sensitive to type I interferon pretreatment. *BioRxiv.* 2020 March 07.982264. doi:10.1101/2020.03.07.982264.
128. Thoms M, Buschauer R, Ameisemeier M, Koepke L, Denk T, Hirschenberger M, Kratzat H, Hayn M, Mackens-Kiani T, Cheng J, et al. Structural basis for translational shutdown and immune evasion by the Nsp1 protein of SARS-CoV-2. *Science.* 2020;369(6508):1249–55. doi:10.1126/science.abc8665.
129. Konno Y, Kimura I, Uriu K, Fukushima M, Irie T, Koyanagi Y, Nakagawa S, Sato K. SARS-CoV-2 ORF3b is a potent interferon antagonist whose activity is further increased by a naturally occurring elongation variant. *bioRxiv.* 2020 May 11.088179. doi:10.1101/2020.05.11.088179.
130. Antia A, Ahmed H, Handel A, Carlson NE, Amanna IJ, Antia R, Slika M. Heterogeneity and longevity of antibody memory to viruses and vaccines. *PLoS Biol.* 2018;16(8):e2006601. doi:10.1371/journal.pbio.2006601.
131. Mayer A, Zhang Y, Perelson AS, Wingreen NS. Regulation of T cell expansion by antigen presentation dynamics. *Proc Natl Acad Sci U S A.* 2019;116(13):5914–19. doi:10.1073/pnas.1812800116.
132. Zinkernagel RM. Immunological memory not equal protective immunity. *Cell Mol Life Sci: CMLS.* 2012;69(10):1635–40. doi:10.1007/s00018-012-0972-y.
133. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu N-H, Nitsche A, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell.* 2020;181(2):271–80e8. doi:10.1016/j.cell.2020.02.052.
134. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol.* 2020;5(4):562–69. doi:10.1038/s41564-020-0688-y.
135. Pinto D, Park YJ, Beltramello M, Walls AC, Tortorici MA, Bianchi S, Jaconi S, Culap K, Zatta F, De Marco A, et al. Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. *Nature.* 2020;583(7815):290–95. doi:10.1038/s41586-020-2349-y.
136. Wang X, Xu W, Hu G, Xia S, Sun Z, Liu Z, Xie Y, Zhang R, Jiang S, Lu L, et al. Retraction Note to: SARS-CoV-2 infects T lymphocytes through its spike protein-mediated membrane fusion. *Cell Mol Immunol.* 2020;17(8):894–894. doi:10.1038/s41423-020-0498-4.
137. Zhou B, She J, Wang Y, Ma X. The duration of viral shedding of discharged patients with severe COVID-19. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa451.
138. Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Moderbacher CR, Rawlings SA, Sutherland A, Premkumar L, Jardi RS, et al. Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell.* 2020;181(7):1489–1501.e15.
139. Braun J, Loyal L, Frentsch M, Wendisch D, Georg P, Kurth F, Hippenstiel S, Dingeldey M, Kruse B, Fauchere F, et al. Presence of SARS-CoV-2 reactive T cells in COVID-19 patients and healthy donors. *medRxiv.* 2020 Apr 17. 20061440. doi:10.1101/2020.04.17.20061440.
140. Alexander J, Bilsel P, Del Guercio MF, Marinkovic-Petrovic A, Southwood S, Stewart S, Ishioka G, Kotturi MF, Botten J, Sidney J, et al. Identification of broad binding class I HLA supertype epitopes to provide universal coverage of influenza A virus. *Hum Immunol.* 2010;71(5):468–74. doi:10.1016/j.humimm.2010.02.014.
141. Masters PS. The molecular biology of coronaviruses. *Adv Virus Res.* 2006;66:193–292.
142. Zhao Z, Li H, Wu X, Zhong Y, Zhang K, Zhang YP, Boerwinkle E, Fu YX. Moderate mutation rate in the SARS coronavirus genome and its implications. *BMC Evol Biol.* 2004; 28(4):21. doi:10.1186/1471-2148-4-21.
143. Menachery VD, Yount BL Jr, Josset L, Gralinski LE, Scobey T, Agnihothram S, Katze MG, Baric RS. Attenuation and restoration of severe acute respiratory syndrome coronavirus mutant lacking 2'-O-methyltransferase activity. *J Virol.* 2014;88(8):4251–64. doi:10.1128/JVI.03571-13.
144. Elbahesh H, Saletti G, Gerlach T, Rimmelzwaan GF. Broadly protective influenza vaccines: design and production platforms. *Curr Opin Virol.* 2019;34:1–9. doi:10.1016/j.coviro.2018.11.005.
145. Vemula SV, Sayedahmed EE, Sambhara S, Mittal SK. Vaccine approaches conferring cross-protection against influenza viruses. *Expert Rev Vaccines.* 2017;16(11):1141–54. doi:10.1080/14760584.2017.1379396.
146. Milian E, Kamen AA. Current and emerging cell culture manufacturing technologies for influenza vaccines. *Biomed Res Int.* 2015;2015:504831. doi:10.1155/2015/504831.
147. Soema PC, Kompier R, Amorij JP, Kersten GF. Current and next generation influenza vaccines: formulation and production strategies. *Eur J Pharm Biopharm.* 2015;94:251–63. doi:10.1016/j.ejpb.2015.05.023.
148. Brown LE, Kelso A. Prospects for an influenza vaccine that induces cross-protective cytotoxic T lymphocytes. *Immunol Cell Biol.* 2009;87(4):300–08. doi:10.1038/icb.2009.16.
149. Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Moderbacher CR, Rawlings SA, Sutherland A, Premkumar L, Jardi RS, et al. Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell.* 2020;181(7):1489–501.e15.
150. Gilbert SC. T-cell-inducing vaccines - what's the future. *Immunology.* 2012;135(1):19–26. doi:10.1111/j.1365-2567.2011.03517.x.
151. Braun J, Loyal L, Fnotesrentsch M, Wendisch D, Georg P, Kurth F, Hippenstiel S, Dingeldey M, Kruse B, Fauchere F, et al. SARS-CoV-2-reactive T cells in healthy donors and patients with COVID-19. *Nature.* 2020. doi:10.1038/s41586-020-2598-9.
152. Yuan J, Kou S, Liang Y, Zeng J, Pan Y, Liu L. PCR assays turned positive in 25 discharged COVID-19 patients. *Clin Infect Dis.* 2020. doi:10.1093/cid/ciaa398.
153. Nakayama T. Causal relationship between immunological responses and adverse reactions following vaccination. *Vaccine.* 2019;37(2):366–71. doi:10.1016/j.vaccine.2018.11.045.
154. Chen J. Pathogenicity and transmissibility of 2019-nCoV-A quick overview and comparison with other emerging viruses. *Microbes Infect.* 2020;22(2):69–71. doi:10.1016/j.micinf.2020.01.004.
155. Metcalf CJE, Ferrari M, Graham AL, Grenfell BT. Understanding herd immunity. *Trends Immunol.* 2015;36(12):753–55. doi:10.1016/j.it.2015.10.004.
156. Rodpothong P, Auewarakul P. Viral evolution and transmission effectiveness. *World J Virol.* 2012;1(5):131–34. doi:10.5501/wjv.v1.i5.131.

ORIGINALITY REPORT

% **13**
SIMILARITY INDEX

% **9**
INTERNET SOURCES

% **10**
PUBLICATIONS

% **4**
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to University of the Pacific
Student Paper % **1**

2 www.koelnmesse.jp
Internet Source <% **1**

3 dokumen.pub
Internet Source <% **1**

4 Kuldeep Dhama, Senthilkumar Natesan, Mohd. Iqbal Yattoo, Shailesh Kumar Patel et al. "Plant-based vaccines and antibodies to combat COVID-19: current status and prospects", Human Vaccines & Immunotherapeutics, 2020
Publication <% **1**

5 tampub.uta.fi
Internet Source <% **1**

6 Mohd. Iqbal Yattoo, Zeenat Hamid, Izhar Rather, Qurat Ul Ain Nazir et al. "Immunotherapies and immunomodulatory approaches in clinical trials - a mini review", Human Vaccines & Immunotherapeutics, 2021
Publication <% **1**

7	ijbs.com Internet Source	<% 1
8	hdl.handle.net Internet Source	<% 1
9	discovery.ucl.ac.uk Internet Source	<% 1
10	Xuesong Gao, Zheng Yuan, Di Yang, Hongjie Li, Yijin Zhang, Ping Gao, Xiaomin Liu, Wenshan Zhao, Te Xiao, Xuefei Duan. "A Family Cluster of Severe Acute Respiratory Syndrome Coronavirus 2 Infections", Research Square, 2020 Publication	<% 1
11	pubcovid19.pt Internet Source	<% 1
12	Submitted to Jordan University of Science & Technology Student Paper	<% 1
13	Alexander A. Lehmann, Greg A. Kirchenbaum, Ting Zhang, Pedro A. Reche, Paul V. Lehmann. "Deconvoluting the T cell response to SARS-CoV-2: specificity versus chance- and cognate cross-reactivity", Cold Spring Harbor Laboratory, 2020 Publication	<% 1

14

patents.justia.com

Internet Source

<% 1

15

bmcpublichealth.biomedcentral.com

Internet Source

<% 1

16

dels.nas.edu

Internet Source

<% 1

17

Rayhane Nchioua, Dorota Kmiec, Janis Mueller, Carina Conzelmann et al. "The Zinc Finger Antiviral Protein restricts SARS-CoV-2", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

18

M Sokolowska, Z Lukasik, I Agache, CA Akdis et al. "Immunology of COVID-19: mechanisms, clinical outcome, diagnostics and perspectives – a report of the European Academy of Allergy and Clinical Immunology (EAACI)", Allergy, 2020

Publication

<% 1

19

Ali A. Rabaan, Shamsah H. Al-Ahmed, Ranjit Sah, Jaffar. A. Al-Tawfiq et al. "Recent advances in vaccine and immunotherapy for COVID-19", Human Vaccines & Immunotherapeutics, 2020

Publication

<% 1

20

apps.who.int

Internet Source

<% 1

21 yuan yuan, Jun He, Lei Gong, Weiwei Li et al. "Molecular epidemiology of SARS-CoV-2 clusters caused by asymptomatic cases in Anhui Province, China", Research Square, 2020
Publication <% 1

22 Annsea Park, Akiko Iwasaki. "Type I and Type III Interferons – Induction, Signaling, Evasion, and Application to Combat COVID-19", Cell Host & Microbe, 2020
Publication <% 1

23 Submitted to East Tennessee State University
Student Paper <% 1

24 jprsolutions.info
Internet Source <% 1

25 Submitted to University of Arizona
Student Paper <% 1

26 www.immunology.ox.ac.uk
Internet Source <% 1

27 Submitted to American University of Beirut
Student Paper <% 1

28 Limin Wang, Julian Candia, Lichun Ma, Yongmei Zhao et al. "Serological Responses to Human Virome Define Clinical Outcomes of Italian Patients Infected with SARS-CoV-2", Cold Spring Harbor Laboratory, 2020
Publication <% 1

-
- 29 Submitted to Al-Nahrain University <% 1
Student Paper
-
- 30 Submitted to Syiah Kuala University <% 1
Student Paper
-
- 31 mbio.asm.org <% 1
Internet Source
-
- 32 outbreakdaily.wp.tulane.edu <% 1
Internet Source
-
- 33 repub.eur.nl <% 1
Internet Source
-
- 34 Peng-jiao An, Yi Zhun Zhu, Li-ping Yang. <% 1
"Biochemical indicators of coronavirus disease
2019 exacerbation and the clinical implications",
Pharmacological Research, 2020
Publication
-
- 35 Francesco Antonio Ciarleglio, Marta Rigoni,
Liliana Mereu, Tommaso Cai et al. "Impact of
Covid-19 Disease and National Lockdown on
Outcomes of Emergency Surgery. Results From
A Retrospective Comparative Cohort Study In A
Tertiary Referral Teaching Hospital", Research
Square, 2021 <% 1
Publication
-
- 36 www.dovepress.com <% 1
Internet Source
-

37	www.medsci.org Internet Source	<% 1
38	actaneurocomms.biomedcentral.com Internet Source	<% 1
39	Rynda-Apple, Agnieszka, Dustin P Patterson, and Trevor Douglas. "Virus-like particles as antigenic nanomaterials for inducing protective immune responses in the lung", <i>Nanomedicine</i> , 2014. Publication	<% 1
40	www.bioscience.org Internet Source	<% 1
41	molecular-cancer.biomedcentral.com Internet Source	<% 1
42	www.mycoastnow.com Internet Source	<% 1
43	annalsofintensivecare.springeropen.com Internet Source	<% 1
44	E.O. Gubernatorova, E.A. Gorshkova, A.I. Polinova, M.D. Drutskaya. "IL-6: relevance for immunopathology of SARS-CoV-2", <i>Cytokine & Growth Factor Reviews</i> , 2020 Publication	<% 1
45	John S. Tregoning, Emily S. Brown, Hannah M. Cheeseman, Katie E. Flight et al. "Vaccines for	<% 1

COVID-19", Clinical & Experimental Immunology, 2020

Publication

46

eprints.gla.ac.uk

Internet Source

<% 1

47

Alyssa Silva-Cayetano, William S. Foster, Silvia Innocentin, Sandra Belij-Rammerstorfer et al. "A booster dose enhances immunogenicity of the COVID-19 vaccine candidate ChAdOx1 nCoV-19 in aged mice", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

48

David Baker, Sandra Amor, Angray S. Kang, Klaus Schmierer, Gavin Giovannoni. "The underpinning biology relating to multiple sclerosis disease modifying treatments during the COVID-19 pandemic.", Multiple Sclerosis and Related Disorders, 2020

Publication

<% 1

49

fjfsdata01prod.blob.core.windows.net

Internet Source

<% 1

50

www.jbc.org

Internet Source

<% 1

51

Patricia Figueiredo-Campos, Birte Blankenhaus, Catarina Mota, Andreia Gomes et al. "Seroprevalence of anti-SARS-CoV-2 antibodies

<% 1

in COVID-19 patients and healthy volunteers",
Cold Spring Harbor Laboratory, 2020

Publication

52

jurnal.ugm.ac.id

Internet Source

<% 1

53

worldwidescience.org

Internet Source

<% 1

54

www.ebmedicine.net

Internet Source

<% 1

55

Wangxue Chen. "Promise and challenges in the development of COVID-19 vaccines", Human Vaccines & Immunotherapeutics, 2020

Publication

<% 1

56

Prashant Dogra, Javier Ruiz-Ramírez, Kavya Sinha, Joseph D. Butner et al. "Innate immunity plays a key role in controlling viral load in COVID-19: mechanistic insights from a whole-body infection dynamics model", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

57

Matan Yechezkel, Amit Weiss, Idan Rejwan, Edan Shahmoon, Shachaf Ben-Gal, Dan Yamin. "Human mobility and poverty as key factors in strategies against COVID-19", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

58

jfmipc.com

Internet Source

<% 1

59

Carolina M Voloch, Ronaldo da Silva F, Luiz G P de Almeida, Otavio J. Brustolini et al. "Intra-host evolution during SARS-CoV-2 persistent infection", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

60

www.escardio.org

Internet Source

<% 1

61

Colin Pawlowski, AJ Venkatakrishnan, Eshwan Ramudu, Christian Kirkup et al. "Pre-existing conditions are associated with COVID patients' hospitalization, despite confirmed clearance of SARS-CoV-2 virus", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

62

Zhiyong Han, Fortunato Battaglia, Stanley R Terlecky. "Discharged COVID-19 patients testing positive again for SARS-CoV-2 RNA: A minireview of published studies from China", Journal of Medical Virology, 2020

Publication

<% 1

63

Ania Wajnberg, Mayce Mansour, Emily Leven, Nicole M Bouvier et al. "Humoral immune response and prolonged PCR positivity in a cohort of 1343 SARS-CoV 2 patients in the New

<% 1

York City region", Cold Spring Harbor Laboratory, 2020

Publication

64

Laura A Bienvenu, Jonathan Noonan, Xiaowei Wang, Karlheinz Peter. "Higher mortality of COVID-19 in males: sex differences in immune response and cardiovascular comorbidities", Cardiovascular Research, 2020

Publication

<% 1

65

Raife Dilek Turan, Cihan Tastan, Derya Dilek Kancagi, Bulut Yurtsever et al. "Gamma-irradiated SARS-CoV-2 vaccine candidate, OZG-38.61.3, confers protection from SARS-CoV-2 challenge in human ACEII-transgenic mice", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

66

Sebastian L. Johnston, Paul F McKay, Tatiana Keadze, Kai Hu et al. "Evaluation of the Abbott Architect, Roche Elecsys and Virtus S1 SARS-CoV-2 antibody tests in community-managed COVID-19 cases", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

67

m.amedeo.com

Internet Source

<% 1

68

daten-quadrat.de

Internet Source

<% 1

69

Zoe Swank, Grégoire Michielin, Hon Ming Yip, Patrick Cohen et al. "A high-throughput microfluidic nano-immunoassay for detecting anti-SARS-CoV-2 antibodies in serum or ultra-low volume dried blood samples", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

70

Chan, Jasper F. W., Susanna K. P. Lau, Kelvin K. W. To, Vincent C. C. Cheng, Patrick C. Y. Woo, and Kwok-Yung Yuen. "Middle East Respiratory Syndrome Coronavirus: Another Zoonotic Betacoronavirus Causing SARS-Like Disease", Clinical Microbiology Reviews, 2015.

Publication

<% 1

71

Jason Rosado, Charlotte Cockram, Sarah Helene Merklung, Caroline Demeret et al. "Serological signatures of SARS-CoV-2 infection: Implications for antibody-based diagnostics", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

72

Haitao Yang, Yuzhu Lan, Xiujuan Yao, Sheng Lin, Baosong Xie. "Evaluation on the Diagnostic Efficiency of Different Methods in Detecting COVID-19.", Research Square, 2020

Publication

<% 1

73

Estela Gimenez, Eliseo Albert, Ignacio Torres,

<% 1

Maria Jose Remigia et al. "SARS-CoV-2-reactive interferon- γ -producing CD8+ T cells in patients hospitalized with Coronavirus viral disease-2019", Cold Spring Harbor Laboratory, 2020

Publication

74

Naru Zhang, Shibo Jiang, Lanying Du. "Current advancements and potential strategies in the development of MERS-CoV vaccines", Expert Review of Vaccines, 2014

Publication

75

Bin Zhou, Tran Thi Nhu Thao, Donata Hoffmann, Adriano Taddeo et al. "SARS-CoV-2 spike D614G variant confers enhanced replication and transmissibility", Cold Spring Harbor Laboratory, 2020

Publication

76

Tao Liu, Sanyun Wu, Huangheng Tao, Guang Zeng, Fuling Zhou, Xinghuan Wang. "Prevalence of IgG Antibodies to SARS-CoV-2 in Wuhan – Implications for the Longevity of Antibodies Against SARS-CoV-2", Research Square, 2020

Publication

77

Bing Liu, Junyan Han, Xiaohuan Cheng, Long Yu, Li Zhang, Wei Wang, Lan Ni, Chaojie Wei, Yafei Huang, Zhenshun Cheng. "Persistent

<% 1

<% 1

<% 1

<% 1

SARS-CoV-2 presence is accompanied with defects in adaptive immune system in non-severe COVID-19 patients", Cold Spring Harbor Laboratory, 2020

Publication

78

www.tremedica.org

Internet Source

<% 1

79

Elisa Kortela, Vesa Kirjavainen, Maarit J. Ahava, Suvi T. Jokiranta et al. "Real-life clinical sensitivity of SARS-CoV-2 RT-PCR test in symptomatic patients", Cold Spring Harbor Laboratory, 2020

Publication

<% 1

80

Harapan Harapan, Naoya Itoh, Amanda Yufika, Wira Winardi et al. "Coronavirus disease 2019 (COVID-19): A literature review", Journal of Infection and Public Health, 2020

Publication

<% 1

81

Harapan Harapan, Mirza Ryan, Benediktus Yohan, Rufika Shari Abidin et al. "Covid-19 and dengue: Double punches for dengue-endemic countries in Asia", Reviews in Medical Virology, 2020

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE ON

BIBLIOGRAPHY

EXCLUDE MATCHES < 5

WORDS