

# Effect of Vitamin B Complex and Phosphorus on Stress Resistance and Phosphorus Uptake in Barramund

*by Aslamiyah --*

---

**Submission date:** 29-Apr-2023 02:02PM (UTC+0700)

**Submission ID:** 2079051670

**File name:** Ahmad\_Yani\_2022\_Israeli.pdf (451.71K)

**Word count:** 5212

**Character count:** 26995



The IJA is a peer-reviewed open-access, electronic journal, freely available without charge to users  
Produced by the AquacultureHub non-profit Foundation  
Sale of IJA papers is strictly forbidden



2

## Effect of Vitamin B Complex and Phosphorus on Stress Resistance and Phosphorus Uptake in Barramundi

Ahmad Yani<sup>1\*</sup>, Muhammad Yusri Karim<sup>2</sup>, Zainuddin Zainuddin<sup>2</sup>,  
Siti Aslamiyah<sup>2</sup>

<sup>1</sup> Doctoral Study Program in Fisheries Science, Faculty of Marine Science and Fisheries,  
Hasanuddin University, Makassar, Indonesia

<sup>2</sup> Department of Fisheries, Faculty of Marine Science and Fisheries, Hasanuddin  
University, Makassar, 90245, Indonesia

(Received Aug 18, 2022; Accepted Dec 16, 2022; Published Dec 27, 2022)

Keywords: aquaculture media enrichment, domestication, larviculture, *Lates calcarifer*

### Abstract

Barramundi, the Asian sea bass (*Lates calcarifer* Bloch 1790), is a valuable aquaculture commodity, but seed survival is generally low. This study used a completely randomized design (two factors; 16 treatments; 3 replicates) to examine the effect of adding vitamin B complex and phosphorus to the rearing media on the survival, stress resistance, and phosphorus content of barramundi larvae. Barramundi larvae aged two days were stocked at a density of 25 fish L<sup>-1</sup> in 40 L green plastic basins (30 L water) and reared for 30 days with natural feed (Chlorella and rotifers). The factor I was vitamin B complex with 4 levels: 0, 75, 150, and 225 mg L<sup>-1</sup>. Factor II was phosphorus with four levels: 0, 0.5, 1.0, and 1.5 mg L<sup>-1</sup>. Commercially available B vitamin complex and phosphorus mineral supplements were dissolved in water and added to the larval-rearing media. Analysis of variance showed that the combination of vitamin B complex and phosphorus treatments had a significant effect ( $P < 0.01$ ) on survival, stress resistance, and phosphorus uptake. Survival and phosphorus content in barramundi larvae were highest with a combination of 150 mg L<sup>-1</sup> vitamin B complex and 1.0 mg L<sup>-1</sup> phosphorus and lowest in control (0 mg L<sup>-1</sup> of both additives).

\* Corresponding author. Tel.: +62-813-24015293, e-mail: ahmadyani\_fpapua\_papua@yahoo.com

## Introduction

Barramundi (*Lates calcarifer* Bloch 1790), also known as the Asian sea bass, is a high-value euryhaline fish with good potential as an aquaculture commodity because of its relatively fast growth and ease of adaptation to the culture environment, in particular concerning salinity (Wijayanto et al., 2020). One of the determining factors for the success of barramundi culture is the availability of seeds (Thépot et al., 2016). In Indonesia, most barramundi seeds are still wild-caught, and availability is seasonal. The development of seed production in hatcheries is needed to support the sustainable development of the barramundi culture (Wijayanto et al., 2020). Several hatcheries have succeeded in the captive breeding of barramundi, but the survival of larvae to fingerling size is often still very low (Thépot et al., 2016). The larvae and fingerlings up to around 7 cm in length the larvae/fingerlings are easily stressed, leading to high mortality rates, especially at high stocking densities, which can lead to reduced dissolved oxygen concentrations making the fish vulnerable to stress (Noval et al., 2018). Several studies have reported low survival rates from 0 % to around 30 % (Hassan et al., 2021; Noval et al., 2019; Srichanun et al., 2014; Thépot et al., 2016; Wijayanto et al., 2020). High larval mortality rates can be due to problems with environmental factors and poor quality or inappropriate feed (Srichanun et al., 2014; Thépot et al., 2016). Therefore, it is necessary to provide the larvae with suitable high quality feed and a rearing environment that meets their needs (Noval et al., 2019; Srichanun et al., 2014; Thépot et al., 2016; Wijayanto et al., 2020).

One way to increase barramundi larvae's survival and stress resistance is by improving nutrition. Two potential supplements are the vitamin B complex and the mineral phosphorus. The benefits of B vitamins in humans include reducing fatigue, supporting the normal functioning of the immune system, and nervous systems, and red blood cell formation (Shabbir et al., 2020). Similarly, in fish, the B vitamins, especially vitamin B6, play a role in forming red blood cells and synthesizing insulin and hemoglobin. In contrast, vitamin B12 regulates the brain and nervous system (Vilain and Baran, 2016). Research on the use of B vitamins in fish has shown positive effects. For example, supplementation with B vitamins resulted in 100 % survival of Atlantic salmon (*Salmo salar* Linnaeus 1758) (Hemre et al., 2012) with similar results reported in other taxa, including juvenile crabs and fishes (Hansen et al., 2015; Wei et al. 2014).

In addition to vitamin B complex, phosphorus supplementation could also be expected to reduce stress levels in barramundi larvae. Increasing the availability of phosphorus and, thereby, phosphorus uptake can significantly increase the growth performance (e.g., final body weight, specific growth rate, and feeding rate) of juvenile yellow catfish *Pelteobagrus fulvidraco* Richardson 1846 (Tang et al., 2012; Wang et al., 2016), rainbow trout *Oncorhynchus mykiss* Walbaum 1792 (Hernández and Roman, 2016; Morales et al., 2018), Prussian carp *Carassius auratus gibelio* var. Bloch 1782 (Jiang et al. 2012), and Nile tilapia *Oreochromis niloticus* Linnaeus 1758 (Dogan et al. 2015). Furthermore, the provision of phosphorus in a mineral form is expected to increase phosphorus uptake and, thus, the phosphorus content of larvae.

The combination of appropriate levels of supplementation with vitamin B complex and phosphorus should result in improved survival, stress resistance, and phosphorus uptake in barramundi larvae. However, there needs to be more data on phosphorus and B vitamin complex supplementation in barramundi larvae, including the appropriate dosage or quantity of each supplement. Therefore, research is needed to determine the appropriate levels of each supplement for the optimal combination of vitamin B complex and phosphorus when rearing barramundi larvae. This study examined the effect of combining different levels of vitamin B complex and phosphorus supplementation on the survival, stress resistance, and phosphorus content of barramundi (*L. calcarifer*) larvae.

32  
Materials and Methods

## Experimental material

The study was carried out at the Takalar Brackish Water Aquaculture Centre. The 48 rearing containers used were 40 L green plastic basins with a length of 30 cm, a width of 50 cm, and a height of 30 cm filled with 30 L of water and equipped with aeration equipment. Before use, the basins were sterilized by soaking in a 10 ppm chlorine solution for 24 hours; the chlorine was then neutralized with a 5 ppm solution of sodium thiosulfate ( $\text{Na}_2\text{S}_2\text{O}_3$ ). The culture media was sea water obtained from the coastal waters in Galesong, Takalar Regency. The water was pumped using a submersible pump, and the pipe's end was covered with a filter bag (20x40 cm). The water was then filtered through a sand filter and stored in a water reservoir.

The experimental animals used were barramundi (*Lates calcarifer*) larvae aged two days after the egg yolk had been exhausted. The larvae were obtained from the spawning barramundi broodstock at the Takalar Brackish Water Aquaculture Centre (BPBAP). Larvae were stocked at a density of 25 fish  $\text{L}^{-1}$  and reared for 30 days. Salinity was maintained within the 30-31 ppt range with a temperature range of 29-31°C. The experimental fish were fed with natural feed from *Chlorella* and rotifers. Both natural feeds were obtained from mass culture systems at the Takalar Brackish Water Aquaculture Centre (BPBAP).

## Experimental Feed

The vitamin B complex was a readily available commercial formulation produced by PT. Indosco, Surabaya. The vitamin B complex composition was: Vitamin B1 (10,000 mg); B2 (20,000 mg); B6 (10,000 mg); B12 (10,000 mcg); Ca-d pantothenate (20,000 mg); nicotinic Acid (50,000 mg). The phosphorus mineral supplement used in this research was commercially available  $\text{Na}_2\text{HPO}_4$ . To get the desired phosphorus content in the compound  $\text{Na}_2\text{HPO}_4$  (2:1:1:4), the mass ratio (46:1:31:64) was considered.

The vitamin B complex and phosphorus supplements were administered through the culture medium (water) on the first day after stocking in doses appropriate for each treatment. The same vitamin B complex and phosphorus doses were administered every 2 days after changing the water until the end of the study. The vitamin B complex and phosphorus, according to each treatment, were mixed with water in a 1 L dipper before being added to the rearing containers. This according to the research of Nurfadilah (2021), that implemented the same study to evaluate the effect of vitamin B complex on the survival rate, growth, and stress resistance of Nile tilapia (*O. niloticus*).

The study used a completely randomized design (CRD) with two factors. The factor reference refers to Nurfadilah (2021), which uses the same treatment in the research. The factor I was phosphorus, with four levels: 0, 0.5, 1.0, and 1.5 mg  $\text{L}^{-1}$ . Factor II was a soluble vitamin B complex with four levels: 0, 75, 150, and 225 mg  $\text{L}^{-1}$ . With 16 treatment combinations and three replicates, there were 48 experimental units (Table 1).

**Table 1** Phosphorus (P) and vitamin B complex (B) treatment combinations

Treatment		P	B	Treatment		P	B
Letter	Code	( $\text{mg L}^{-1}$ )	( $\text{mg L}^{-1}$ )	Letter	Code	(mg)	(mg)
A	F1B1	0	0	I	F3B1	1.0	0
B	F1B2	0	75	J	F3B2	1.0	75
C	F1B3	0	150	K	F3B3	1.0	150
D	F1B4	0	225	L	F3B4	1.0	225
E	F2B1	0.5	0	M	F4B1	1.5	0
F	F2B2	0.5	75	N	F4B2	1.5	75
G	F2B3	0.5	150	O	F4B3	1.5	150
H	F2B4	0.5	225	P	F4B4	1.5	225

### Procedures and variables

The variables observed in this study were larval survival, stress resistance, and the amount of phosphorus absorbed and deposited by the larvae. The survival rate was calculated as the percentage of live barramundi larvae at the end of the experiment. The stress resistance of the barramundi larvae was evaluated using the Cumulative Stress Index (CSI). A stress resistance test was carried out to evaluate the physiological condition of barramundi larvae under the various phosphorous and vitamin B complex treatments. The test followed the methods in Roy et al. (2002) to measure the resistance of barramundi larvae to osmotic shock. At the end of the study, ten barramundi larvae were randomly taken from each experimental unit and placed in a glass beaker filled with 1 L of low-salinity water (0 - 5 ppt). The number of stressed barramundi larvae was observed at 5-minute intervals for one hour. Stress assessment was based on behavioral responses such as abnormal movements of the barramundi larvae or death. Abnormal movements indicative of stress included unusual swimming up and down behavior and spinning around until death occurred. The Cumulative Stress Index (CSI) was calculated using the following formula adapted from Shabbir et al. (2020):

$$[CSI = D5 + D10 + D15 + \dots + D60]$$

Where:

DX = number of larvae exhibiting stress after X minutes when exposed to the osmotic shock test.

The phosphorus content of the barramundi larvae was calculated using the procedures described in the Association of Analytical Chemists (2012) in adapting and summarizing from the original publication reports on two experiments conducted to characterize better the nutritional value of a high-quality fish meal against standard fish byproduct meal in barramundi and the potential for phosphorus supplementation (Schreck and Tort 2016). Supporting data on water quality were collected throughout the research period. Temperature and dissolved oxygen (DO) were measured using a DO meter (YSI 58, Yow Springs Instrument Co. Inc., USA), and pH was measured using a pH meter (Meterlab PHM 201, Radiometer Analytical, S.A., France). Temperature, pH, and DO were measured twice a day, in the morning (08:00) and afternoon (16:00). Ammonia levels were measured three times at the experiment's start, middle, and end. Water samples were collected in bottles from each experimental unit and taken to the Water Quality Laboratory in the Fisheries Department of the Faculty of Marine Science and Fisheries at Hasanuddin University, where ammonia levels were measured using a spectrophotometer.

### Statistical analysis

The data obtained were tabulated in Microsoft Excel 2010 and reported as mean values and standard deviation (SD). All statistical analyses were carried out in SPSS version 23.0. Analysis of variance (ANOVA) was used to evaluate treatment differences at the 95 % confidence level ( $\alpha = 0.05$ ). When significant differences were found, the W-Tukey posthoc test was applied. The water quality parameters were analyzed descriptively concerning the barramundi larvae survival rates.

## Results

### Survival rate, stress resistance, and phosphorus uptake

The survival rate, cumulative stress index (CSI), and phosphorus content of the barramundi larvae under the 16 vitamin B complex and phosphorus treatments are shown in **Table 2**. The ANOVA indicated that the vitamin B complex and phosphorus combination had a highly significant effect ( $P < 0.01$ ) on barramundi larval survival rate, CSI, and phosphorus content. The data in Table 1 show that survival was highest under treatment K (150 mg L<sup>-1</sup> vitamin B complex and 1.0 mg L<sup>-1</sup> phosphorus), reaching values around 60 %, more than 10 times higher than the lowest survival rate of around 5 %

under the control treatment with no added vitamin B or phosphorus during the study, water quality was in fairly good condition, except for low dissolved oxygen in the morning. Temperatures ranged between 27 – 32°C, salinity 28 - 35 ppt, pH 6.9 – 8.9, dissolved oxygen 2.1 – 5.6 ppm, and ammonia 0.002 ppm.

**Table 2** The survival rate, cumulative stress index (CSI), and phosphorus content of barramundi (*Lates calcarifer*) larvae under vitamin B complex and phosphorus treatments.

Vitamin B Complex and Phosphorus Treatment (mg L <sup>-1</sup> )	Survival Rate (%)	Cumulative Stress Index (CSI)	Phosphorus Content (%)
A 0 and 0	5.06 <sup>f</sup> ± 2.095	101.333 <sup>a</sup> ± 0.577	0.365 <sup>a</sup> ± 0.111
B 75 and 0	24.62 <sup>ghi</sup> ± 1.776	93.333 <sup>abcde</sup> ± 1.52	0.377 <sup>a</sup> ± 0.034
C 150 and 0	28.93 <sup>ij</sup> ± 1.638	87.333 <sup>cdefgh</sup> ± 3.512	0.705 <sup>b</sup> ± 0.034
D 225 and 0	15.06 <sup>cde</sup> ± 0.833	91 <sup>abcdef</sup> ± 6.245	0.361 <sup>a</sup> ± 0.031
E 0 and 0.5	17.91 <sup>efg</sup> ± 1.646	95.667 <sup>abcd</sup> ± 2.082	0.548 <sup>c</sup> ± 0.027
F 75 and 0.5	30 <sup>de</sup> ± 2.800	83.333 <sup>efgh</sup> ± 2.517	0.656 <sup>b</sup> ± 0.010
G 150 and 0.5	32.35 <sup>bc</sup> ± 0.758	78.333 <sup>h</sup> ± 3.055	0.770 <sup>b</sup> ± 0.029
H 225 and 0.5	19.95 <sup>cde</sup> ± 2.694	89 <sup>bcd</sup> ± 2.000	0.384 <sup>a</sup> ± 0.039
I 0 and 1.0	10.93 <sup>def</sup> ± 1.638	96.333 <sup>abcd</sup> ± 3.055	0.213 <sup>f</sup> ± 0.077
J 75 and 1.0	38.26 <sup>cde</sup> ± 1.010	81.333 <sup>gh</sup> ± 5.100	0.382 <sup>a</sup> ± 0.077
K 150 and 1.0	59.82 <sup>a</sup> ± 3.041	78.667 <sup>gh</sup> ± 3.512	0.781 <sup>b</sup> ± 0.016
L 225 and 1.0	21.37 <sup>b</sup> ± 0.658	97 <sup>abc</sup> ± 2.646	0.513 <sup>c</sup> ± 0.064
M 0 and 1.5	34.57 <sup>hi</sup> ± 1.150	99 <sup>ab</sup> ± 2.646	0.420 <sup>a</sup> ± 0.077
N 75 and 1.5	30.8 <sup>gh</sup> ± 0.611	88 <sup>cdefgh</sup> ± 3.606	0.461 <sup>d</sup> ± 0.214
O 150 and 1.5	45.06 <sup>gh</sup> ± 8.331	78 <sup>h</sup> ± 5.196	0.724 <sup>b</sup> ± 0.027
P 225 and 1.5	13.86 <sup>hi</sup> ± 2.203	86.333 <sup>defgh</sup> ± 3.786	0.444 <sup>d</sup> ± 0.043

Note: in each column different superscript letters indicate significantly different mean values at the 95 % confidence level (P < 0.05)

The cumulative stress index (CSI) data in **Table 2** show that the stress levels are lowest and therefore stress resistance highest under treatments G, K and O, which were not significantly different from each other. All three treatments had 150 mg L<sup>-1</sup> B vitamin B complex with phosphorus levels of 0.5, 1.0 and 1.5 mg L<sup>-1</sup>, respectively. The highest CSI and therefore the lowest stress resistance occurred under the control treatment A with no additives. The increased stress resistance associated with vitamin B complex was evident at 75 mg L<sup>-1</sup>, highest at 150 mg L<sup>-1</sup> and much lower at the highest rate of 225 mg L<sup>-1</sup>, while all three doses of phosphorus had a similar positive effect.

The treatments had an effect on phosphorus intake, as evidenced by the significant differences (P < 0.05) in the phosphorus content of the barramundi larvae at the end of the study (**Table 2**). Barramundi larvae showed an increase in phosphorus content with increasing doses of phosphorus and vitamin B complex up to 1.5 mg and 150 mg L<sup>-1</sup>, respectively. The highest phosphorus content in barramundi larvae was 0.78 %, obtained under treatment K with the combination of 150 mg L<sup>-1</sup> vitamin B complex and 1.0 mg L<sup>-1</sup> phosphorus. The lowest phosphorus content was 0.21 % under treatment A, with no additives.

#### Water quality

The physical and chemical water quality parameters measured during the experiment (**Table 3**) did not differ significantly between the 16 vitamin B complex and phosphorus enrichment treatments. Throughout the study, the parameters remained within the ranges considered suitable for barramundi larval rearing.

#### Ammonia

Ammonia causes stress and damages gills and other tissues, even in small amounts. Fish exposed to low levels of ammonia over time are more susceptible to bacterial infections, have poor growth, and will not tolerate routine handling as well as they otherwise would. Ammonia is a killer in higher concentrations, and many unexplained production losses have likely been caused by ammonia. But in this research, the

average range of ammonia is in good condition to barramundi larvae between 0.002 – 0.005 ppm.

**Table 3** Water quality parameters were measured in the barramundi larvae rearing units during the 30-day experimental period.

Parameter	Observed Range	Recommended Range*
Temperature (°C)	27 - 32	28.6 - 32.2
pH	6.9 - 8.9	7 - 9
Salinity (ppt)	28 - 35	
Dissolved oxygen (ppm)	2.1 - 5.6	> 4.0
Ammonia (ppm)	0.002 - 0.005	< 0.1

### Discussion

The results indicate that providing barramundi larvae with a combination of phosphorus and vitamin B complex in appropriate dosages can improve the phosphorus content, and that a higher phosphorus content was associated with an improvement in the performance and quality of barramundi larvae, as evidenced by the results of the stress test. The very low survival rate of barramundi larvae in the control treatment A, with all supplementation treatments showing some increase in survival, most likely indicates a deficiency of these nutrients in the larvae.

Stressors can be external, such as environmental conditions, or internal such as injury, disease, or nutritional status. The ability to withstand and recover from stress and stress-induced health problems depends significantly on the condition of an organism, in particular the energy balance and metabolic processes (Rahmi and Ramses, 2017; Tang et al., 2012).

The B vitamins and phosphorus play essential roles in fish metabolic processes, while deficiencies will cause abnormal symptoms (Thépot et al., 2016; Tort, 2011; Vilain and Baran, 2016; Waagbø, 2010; Wang et al., 2021; Wang et al., 2016; Wei et al., 2014); these roles likely contributed to the observed the positive effect of supplementation. Vitamin B deficiency symptoms can include poor appetite, low feed efficiency, and severe digestive problems weight loss, while phosphorus deficiency can result in a loss of or low appetite; these effects can cause weakness and elevated mortality (Thépot et al., 2016; Tort, 2011; Vilain and Baran, 2016). Furthermore, the vitamin B complex also plays a role in antibody production, so the absorption of vitamin B complex by the larvae could improve immunity and resistance to stress as well as disease (Thépot et al., 2016; Tort, 2011).

The vitamin B complex functions as a coenzyme that helps the fish to obtain energy from food (Wijayanto et al., 2020). Thiamine (B1), in particular, aids in releasing energy from food and promotes appetite. Riboflavin (B2) also aids in releasing energy from food and is important for growth, cell development, and the proper functioning of cells within the body. Niacin (B3) plays a role in energy production and other crucial cell functions. Pyridoxine (B6) is crucial for protein metabolism, the production of red blood cells, and the transmission of nerve signals. The amino acid folate, often known as folic acid, aids in the digestion of proteins, encourages the production of red blood cells, and lowers the incidence of spinal abnormalities; Biotin aids in the release of energy from carbohydrates in food in the metabolism of lipids, proteins, carbohydrates; and Pantothenic acid (B5) is involved in energy and hormone production and the metabolism of lipids, proteins and carbohydrate from food. Cyanocobalamin (B12) plays a role in developing genetic material, producing normal red blood cells, and maintaining the nervous system (Thépot et al., 2016; Vilain and Baran, 2016; Tort, 2011). In this study, the ability of the larvae to survive with vitamin B complex supplementation, alone and in combination with phosphorus, is thought to be due to positive effects on metabolic processes, in particular, an increased energy availability.

In particular, phosphorus plays essential roles in muscles and energy metabolism, carbohydrate metabolism, nervous tissue metabolism, regulating amino acids, normalizing blood chemistry, the transport of fatty acids and other lipids, the formation of protein, nucleic acids, and cell membranes; maintaining the balance of hormone production and functions; bone mineralization to form hard tissues, in particular the skeleton and other structural components in the fish body (Waagbø, 2010; Wang et al., 2021; Wang et al., 2016; Wei et al., 2014). However, fish cannot absorb phosphorus from water because the concentration of phosphorus dissolved in water is generally very low (Wang et al., 2016). Phosphorus deficiency affects not only hard tissue, and in humans, is responsible for malformations causing rickets but also affects various intermediate metabolism aspects, thereby affecting weight gain and feed conversion (Tort, 2011; Waagbø, 2010). Signs of phosphorus deficiency have been found in fish, especially bone abnormalities but also loss of appetite; the abnormalities can include curved spines, softened bones, reduced head/skull size, and other skeletal deformities, and in extreme cases, the deleterious effects on bone formation can lead to the death of the fish affected (Tort, 2011; Wei et al., 2014). In some fish species, phosphorus deficiency is characterized by slow growth, poor feed efficiency, and stunted bone mineralization (Waagbø, 2010; Wang et al., 2021; Wang et al., 2016; Wei et al., 2014). Conversely, excess phosphorus intake in the diet or through the gills is also deleterious to fish health (Vilain and Baran, 2016). In this study, phosphorus also appears to be a key factor in improving larval quality; furthermore, phosphorus assists in activating B vitamins through the attachment of phosphate groups (Thépot et al., 2016), the positive interaction between the two additives could be expected.

The high survival rate of barramundi larvae under treatment K indicates that the combination of 150 mg L<sup>-1</sup> vitamin B complex and 1.0 mg L<sup>-1</sup> phosphorus is optimal or very close to the optimum value of both additives for promoting barramundi larval survival. For each of the two additives, this combination represents the middle value in treatments tested, indicating that higher or lower values of each additive are less effective, alone or in combination. In these conditions, Singh (2016) in Nurfadilah (2021) stated that those treatments play a role in improving the function of the adrenal glands that manage stress. If the stress can be overcome, the bodywork system in fish larvae will return to normal, and larvae can exist to live and grow (Rehman et al., 2017).

In conclusion, adding a combination of vitamin B complex and phosphorus to the rearing medium (water) significantly affected the survival, stress resistance, and phosphorus uptake of barramundi (*Lates calcarifer*) larvae. The combination of 150 mg L<sup>-1</sup> vitamin B complex and 1 mg L<sup>-1</sup> phosphorus (treatment K) gave the best results overall, with the highest survival rate (around 60 %) and high-stress resistance and phosphorus uptake. The control treatment (no vitamin B complex or phosphorus intake) performed poorly on all three parameters, with a low survival rate (≈ 5 %), poor stress resistance, and low phosphorus content. All other treatments performed better than the control but less well than treatment K on one or more of the three variables measured, indicating that both higher and lower levels of the vitamin B complex (75 and 300 mg L<sup>-1</sup>) and phosphorus (0.5 and 1.5 mg L<sup>-1</sup>) enrichment were less effective. It can be concluded that the combination of 150 mg L<sup>-1</sup> vitamin B complex and 1 mg L<sup>-1</sup> phosphorus is effective and probably close to optimal for the rearing of barramundi larvae.

### Acknowledgments

The authors acknowledge support during the research from the Dean and staff of the Faculty of Fisheries and Marine Science, Universitas Hasanuddin, and the staff of the Takalar Brackish Water Aquaculture Centre, South Sulawesi, Indonesia.

## References

- Dogan, Z., Atli, G. and Canli, M. 2015. Effects of Lead on ATPases in Tissues of Freshwater Fish (*Oreochromis niloticus*) in Differing Calcium Levels. Turkish Journal of Fisheries and Aquatic Sciences 15:223–233. <https://www.trjfas.org/abstract.php?id=757>.
- Hansen, A. C., Waagbø, R. and Hemre, G. I. 2015. New B vitamin recommendations in fish when fed plant-based diets. Aquaculture Nutrition 21:507–527. <https://doi.org/10.1111/anu.12342>.
- Hassan, H. U., Mohammad Ali, Q., Ahmad, N., Masood, Z., Yeamin Hossain, M., Gabol, K., Khan, W., Hussain, M., Ali, A., Attaullah, M. and Kamal, M. 2021. Assessment of growth characteristics, the survival rate and body composition of Asian sea bass *Lates calcarifer* (Bloch, 1790) under different feeding rates in closed aquaculture system. Saudi Journal of Biological Sciences 28:1324–1330. <https://doi.org/10.1016/j.sjbs.2020.11.056>.
- Hemre, G. I., Lock, E. J., Olsvik, P. A., Hamre, K., Espe, M., Torstensen, B. E., Silva, J., Hansen, A. C., Waagbø, R., Johansen, J. S., Sanden, M. and Sissener, N. H. 2016. Atlantic salmon (*Salmo salar*) require increased dietary levels of B-vitamins when fed diets with high inclusion of plant based ingredients. PeerJ 2016:1–28. <https://doi.org/10.7717/peerj.2493>.
- Hernández, A. J. and Roman, D. 2016. Phosphorus and nitrogen utilization efficiency in rainbow trout (*Oncorhynchus mykiss*) fed diets with lupin (*Lupinus albus*) or soybean (*Glycine max*) meals as partial replacements to fish meal. Czech Journal of Animal Science 61:67–74. doi: 10.17221/8729-CJAS.
- Jiang, H., Yang, H., Kong, X., Wang, S., Liu, D. and Shi, S. 2012. Response of Acid and Alkaline Phosphatase Activities to Copper Exposure and Recovery in Freshwater Fish *Carassius auratus gibelio* var. Life Science Journal 9:233–245. <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.1049.7239&rep=rep1&type=pdf>.
- Morales, G. A., Azcuy, R. L., Casaretto, M. E., Márquez, L., Hernández, A. J., Gómez, F., Koppe, W. and Mereu, A. 2018. Effect of different inorganic phosphorus sources on growth performance, digestibility, retention efficiency and discharge of nutrients in rainbow trout (*Oncorhynchus mykiss*). Aquaculture 495:568–574. <https://doi.org/10.1016/j.aquaculture.2018.06.036>.
- Nurfadilah, Jamaluddin, R. and Yusri, M. K., The Effect of Vitamin B Complex on Survival, Growth and Stress Resistance of Nila Tilapia (*Oreochromis niloticus*) Larvae. International Journal of Scientific and Research Publications, Volume 11, Issue 11. Indonesia. <http://dx.doi.org/10.29322/IJSRP.11.11.2021.p11929>
- Roy, P. K., Witten, P. E., Hall, B. K. and Lall, S. P. 2002. Effects of dietary phosphorus on bone growth and mineralisation of vertebrae in haddock (*Melanogrammus aeglefinus* L.). Fish Physiology and Biochemistry 27:35–48. <https://doi.org/10.1023/B:FISH.0000021778.99011.ce>.
- Schreck, C. B., and Tort, L. 2016. The Concept of Stress in Fish. Fish Physiology 35:1–34. <https://doi.org/10.1016/B978-0-12-802728-8.00001-1>.
- Shabbir, H., Shabbir, I., Aslam, M., Sarwar, M. F., Sarwar, M. H. and Sarwar, M. 2020. Fundamental Aspects of Vitamin B complex in Human Nourishment and Fitness. American Journal of Food Science and Health 6:109–118.
- Srichanun, M., Tantikitti, C., Kortner, T. M., Krogdahl, Å. and Chotikachinda, R. 2014. Effects of different protein hydrolysate products and levels on growth, survival rate and digestive capacity in Asian seabass (*Lates calcarifer* Bloch) larvae. Aquaculture 428–429:195–202. <https://doi.org/10.1016/j.aquaculture.2014.03.004>.
- Tang, Q., Wang, C., Xie, C., Jin, J. and Huang, Y. 2012. Dietary available phosphorus affected growth performance, body composition, and hepatic antioxidant property of juvenile yellow catfish *Pelteobagrus fulvidraco*. The Scientific World Journal 2012. <https://doi.org/10.1100/2012/987570>.
- Thépot, V., Mangott, A. and Pirozzi, I. 2016. Rotifers enriched with a mixed algal diet promote survival, growth and development of barramundi larvae, *Lates calcarifer* (Bloch). Aquaculture Reports 3:147–158. <https://doi.org/10.1016/j.aqrep.2016.02.003>.
- Tort, L. 2011. Stress and immune modulation in fish. Developmental and Comparative Immunology 35:1366–1375.
- Vilain, C. and Baran, E. 2016. Nutritional and Health Value of Fish: the Case of Cambodia. Page World Journal of Zoology. <https://hdl.handle.net/20.500.12348/103>.
- Waagbø, R. 2010. Water-soluble vitamins in fish ontogeny. Aquaculture Research 41:733–744. <https://doi.org/10.1111/j.1365-2109.2009.02223.x>.
- Wang, C., Tang, Q., Luo, W. and Zhao, Y. 2016. Modeling phosphorus metabolism in fish species: an example for juvenile yellow catfish, *Pelteobagrus fulvidraco*. Aquaculture International 24:281–294. <https://doi.org/10.1007/s10499-015-9925-4>.

- Wang, P., Li, X., Xu, Z., Ji, D., He, M., Dang, J. Y. and Jun Leng, X. 2021. The digestible phosphorus requirement in practical diet for largemouth bass (*Micropterus salmoides*) based on growth and feed utilization. *Aquaculture and Fisheries* 7:632–638.
- Wei, J., Yu, N., Tian, W., Zhang, F., Wu, Q., Li, E., Zhang, M., Du, Z., Qin, J. and Chen, L. 2014. Dietary vitamin B12 requirement and its effect on non-specific immunity and disease resistance in juvenile Chinese mitten crab *Eriocheir sinensis*. *Aquaculture* 434:179–183. <https://doi.org/10.1016/j.aquaculture.2014.08.010>.
- Wijayanto, D., Bambang, A. N., Nugroho, R. A., Kurohman, F. and Nursanto, D. B. 2020. The effect of different low salinities on growth, feed conversion ratio, survival rate and profit of asian seabass cultivation. *AACL Bioflux* 13:3706–3712. <http://www.bioflux.com.ro/docs/2020.3706-3712.pdf>.

# Effect of Vitamin B Complex and Phosphorus on Stress Resistance and Phosphorus Uptake in Barramund

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://blr.scholasticahq.com">blr.scholasticahq.com</a> Internet Source	2%
2	<a href="http://buscador.una.edu.ni">buscador.una.edu.ni</a> Internet Source	2%
3	Submitted to Kang Chiao International School Student Paper	1%
4	<a href="http://www.suweco.sk">www.suweco.sk</a> Internet Source	1%
5	C.J. Simon, M.J. Salini, S. Irvin, D. Blyth, N. Bourne, R. Smullen. "The effect of poultry protein concentrate and phosphorus supplementation on growth, digestibility and nutrient retention efficiency in barramundi <i>Lates calcarifer</i> ", <i>Aquaculture</i> , 2019 Publication	1%
6	<a href="http://besta-indonesia.com">besta-indonesia.com</a> Internet Source	1%
7	R Hidayat, M Zainuddin, S Safruddin, A Mallawa, S A Farhum. " Skipjack Tuna ( ) catch	1%

in relation to the Thermal and Chlorophyll-a Fronts during May – July in the Makassar Strait ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

---

8	<a href="https://researchonline.jcu.edu.au">researchonline.jcu.edu.au</a> Internet Source	1 %
9	Florencia Nery Sompie, Jein Rinny Leke, Jacquélien Laihad, Linda Tangkau, Malcky Telleng. "The effect dietary peanut flour ( <i>Arachis hypogaea</i> L.) on the quality of chicken eggs", AIP Publishing, 2023 Publication	1 %
10	<a href="http://www.aiep.pl">www.aiep.pl</a> Internet Source	<1 %
11	Park, I.S.. "Anesthetic effects of lidocaine-hydrochloride on water parameters in simulated transport experiment of juvenile winter flounder, <i>Pleuronectes americanus</i> ", Aquaculture, 20090901 Publication	<1 %
12	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	<1 %
13	<a href="http://www.slideshare.net">www.slideshare.net</a> Internet Source	<1 %
14	Hung Duc Pham, Le-Minh Hoang, Khuong V. Dinh, Muhammad A.B. Siddik, Do-Huu Hoang,	<1 %

Ngo-Van Manh. "Effects of enrichment Artemia with organic selenium and essential fatty acids on growth performance and fatty acid composition of barramundi (*Lates calcarifer*) larvae", *Regional Studies in Marine Science*, 2022

Publication

15

[link.springer.com](https://link.springer.com)

Internet Source

<1 %

16

Submitted to Berkshire College of Agriculture

Student Paper

<1 %

17

Submitted to Curtin University of Technology

Student Paper

<1 %

18

[static-curis.ku.dk](https://static-curis.ku.dk)

Internet Source

<1 %

19

[garuda.kemdikbud.go.id](https://garuda.kemdikbud.go.id)

Internet Source

<1 %

20

[repositorio-aberto.up.pt](https://repositorio-aberto.up.pt)

Internet Source

<1 %

21

[rsdjournal.org](https://rsdjournal.org)

Internet Source

<1 %

22

Submitted to Asian Institute of Technology

Student Paper

<1 %

23

[repository.ung.ac.id](https://repository.ung.ac.id)

Internet Source

<1 %

24	<a href="http://www.thaiscience.info">www.thaiscience.info</a> Internet Source	<1 %
25	Younis Mohd Khan, Mukhtar A. Khan. " Effects of dietary cyanocobalamin on growth performance, non - specific immune response, antioxidant capacity, haematological parameters, body composition and liver cyanocobalamin concentration of fingerling major carp, (Hamilton) ", Aquaculture Nutrition, 2020 Publication	<1 %
26	Yu - Chen Ding, Sih - Shien Lee, Sheng - Kai Peng, Wen - Kai Yang, Tsung - Han Lee. " Salinity - dependent changes in branchial morphometry and Na , K - ATPase responses of euryhaline Asian sea bass, ", Journal of Experimental Zoology Part A: Ecological and Integrative Physiology, 2023 Publication	<1 %
27	<a href="http://akuakultur.fpik.undip.ac.id">akuakultur.fpik.undip.ac.id</a> Internet Source	<1 %
28	<a href="http://digitalarchive.worldfishcenter.org">digitalarchive.worldfishcenter.org</a> Internet Source	<1 %
29	<a href="http://ebvgroup4.wikispaces.com">ebvgroup4.wikispaces.com</a> Internet Source	<1 %
30	<a href="http://repositorio.ufmg.br">repositorio.ufmg.br</a> Internet Source	<1 %

- 
- 31 [www.frontiersin.org](http://www.frontiersin.org) Internet Source <1 %
- 
- 32 [www.scielo.br](http://www.scielo.br) Internet Source <1 %
- 
- 33 [www.thejaps.org.pk](http://www.thejaps.org.pk) Internet Source <1 %
- 
- 34 [www.walshmedicalmedia.com](http://www.walshmedicalmedia.com) Internet Source <1 %
- 
- 35 Ali Mohd Yattoo, Md. Niamat Ali, Zarka Zaheen, Zahoor Ahmad Baba et al. "Assessment of pesticide toxicity on earthworms using multiple biomarkers: a review", Environmental Chemistry Letters, 2022  
Publication <1 %
- 
- 36 Gro-Ingunn Hemre, Erik-Jan Lock, Pål Asgeir Olsvik, Kristin Hamre et al. " Atlantic salmon ( ) require increased dietary levels of B-vitamins when fed diets with high inclusion of plant based ingredients ", PeerJ, 2016  
Publication <1 %
- 
- 37 Karima El - Naggat, Radi Mohamed, Mohamed I. El - katcha, Safaa E. Abdo, Mosaad A. Soltan. "Plant Ingredient diet supplemented with lecithin as fish meal and fish oil alternative affects growth <1 %

performance, serum biochemical, lipid metabolism and growth - related gene expression in Nile tilapia", Aquaculture Research, 2021

Publication

---

38	<a href="https://collections.plymouth.ac.uk">collections.plymouth.ac.uk</a> Internet Source	<1 %
39	<a href="https://dspace.plymouth.ac.uk">dspace.plymouth.ac.uk</a> Internet Source	<1 %
40	<a href="https://ecsdev.org">ecsdev.org</a> Internet Source	<1 %
41	<a href="https://imbbs.hcmr.gr">imbbs.hcmr.gr</a> Internet Source	<1 %
42	<a href="https://pubag.nal.usda.gov">pubag.nal.usda.gov</a> Internet Source	<1 %
43	<a href="https://repositorio.ual.es:8080">repositorio.ual.es:8080</a> Internet Source	<1 %
44	<a href="https://silo.tips">silo.tips</a> Internet Source	<1 %
45	<a href="https://vdoc.pub">vdoc.pub</a> Internet Source	<1 %
46	<a href="https://www.science.gov">www.science.gov</a> Internet Source	<1 %
47	Christian E.W. Steinberg. "Aquatic Animal Nutrition", Springer Science and Business	<1 %

---

48

Ibrahim Adeshina, Benjamin U. Akpoilih, Blessing F. Udom, Olarinke V. Adeniyi, Mohsen Abdel-Tawwab. "Interactive effects of dietary phosphorus and microbial phytase on growth performance, intestinal morphometry, and welfare of Nile tilapia (*Oreochromis niloticus*) fed on low-fishmeal diets", *Aquaculture*, 2023

Publication

---

<1 %

49

Mónica Costa, Benjamin Costas, Marina Machado, Carla Teixeira et al. "Anchovy and giant squid hydrolysates can enhance growth and the immune response of European seabass (*Dicentrarchus labrax*) fed plant-protein-based diets", *Aquaculture*, 2020

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      < 5 words

Exclude bibliography      On