

SYNTHESIS,
CHARACTERIZATION AND
CYTOTOXIC ACTIVITY OF
Zn(II) CYSTEINE
DITHIOCARBAMATE IN
BREAST CANCER (MCF-7)
by

FILE	RYZAL_ZN_II_1.PDF (313.06K)	WORD COUNT	2497
TIME SUBMITTED	31-JUL-2019 04:08PM (UTC+0700)	CHARACTER COUNT	13259
SUBMISSION ID	1156462757		



Research Article

15

SYNTHESIS, CHARACTERIZATION AND CYTOTOXIC ACTIVITY OF Zn(II) CYSTEINE DITHIOCARBAMATE IN BREAST CANCER (MCF-7)

Rizal Irfandi ¹, Prihantono ², Indah Raya ^{1*}

¹Department of Chemistry, Faculty of Mathematics, and Natural Science, Hasanuddin University, Perintis Kemerdekaan Street Km. 10 Tamalanrea, Makassar, 90245, Indonesia

²Department of Surgery, Faculty of Medical, Hasanuddin University, Perintis Kemerdekaan Street Km. 10 Tamalanrea, Makassar, 90245, Indonesia

¹Department of Chemistry, Faculty of Mathematics, and Natural Science, Hasanuddin University, Perintis Kemerdekaan Street Km. 10 Tamalanrea, Makassar, 90245, Indonesia

*Corresponding Author Email: indahraya05@gmail.com

Article Received on: 20/03/19 Approved for publication: 19/04/19

DOI: 10.7897/2230-8407.1004126

ABSTRACT

To synthesize and characterize complex compound of Zn(II) with cysteine dithiocarbamate ligands and studies its cytotoxic activity in breast cancer (MCF-7). The complex of Zn(II) cysteine dithiocarbamate was prepared by "in situ method" and characterized by using Ultraviolet-Visible (UV-Vis) and Infra-Red (IR) spectroscopy, melting point and conductivity. The results of UV-Vis maximum spectrums of Zn(II)cysteine dithiocarbamate at 282 nm and 374 nm indicated that electronic transition $\pi \rightarrow \pi^*$ dan $n \rightarrow \pi^*$ of CS₂ and N=C=S. While the presence of IR spectra at the wavelength in the region of 397-534 cm⁻¹ indicated that has been coordination occurred between Zn(II) with Sulphur (S), Nitrogen (N) and Oxygen(O) atoms from cysteine dithiocarbamate ligands. Complex characterization using UV-Vis and IR showed that complex Zn(II) cysteine dithiocarbamate compounds were successfully synthesized by the in-situ method. The results of the cytotoxicity assay of Zn(II) cysteine dithiocarbamate complex compounds in MCF-7 cells obtained IC₅₀ values of Zn(II) complex = 98.60 µg/mL and cisplatin (IC₅₀ = 200 µg/mL). These result can be seen that IC₅₀ value of Zn(II) complex is lower than cisplatin so that Zn(II) complex has very good inhibitory activity against MCF-7 cancer cells compared to cisplatin and it can even cause apoptosis in cancer cells.

Keywords: Complexes; Cysteine; Dithiocarbamate; Apoptosis; Breast cancer; Chemotherapy.

12

INTRODUCTION

Breast cancer (MCF-7) is one of the highest lethal cancers in 7 countries, namely Indonesia, Japan, Malaysia, Philippines, Singapore, Sri Lanka and Taiwan ¹. Treatment of breast cancer is carried out in a multidisciplinary manner such as surgery, chemotherapy and radiation ².

At present chemotherapy is a very important component in the paradigm of handling breast cancer ³. Some drugs that are often used for chemotherapy are platinum derivative compounds, such as cisplatin, carboplatin, and oxaliplatin. Cisplatin is still the main choice in the treatment of cancer. However, cisplatin has very dangerous side effects, especially showing high toxicity in the body ^{4,5,6,7,8}. So that it needs effective and less toxic metal-based anticancer drugs ⁵. The use of essential metals is interesting for researchers to develop metal-based anticancer drugs.

The use of appropriate ligands can significantly increase the activity of complex compounds in inhibiting cancer cells ⁹. This study used a dithiocarbamate ligands. Dithiocarbamate compounds is a material that is very potential in the fields of agriculture, industry, and health. In the field of health, dithiocarbamate compounds can be character-based metal poisoning drugs that act as good chelating agents ^{10,11}. In addition, these compounds can also be used as radio chemotherapy target

agents in tumors ^{12,13}. Dithiocarbamate compounds can be complexed mostly with metal ions from the transition element, because dithiocarbamate compounds have a very special structure where there are S groups that can donate electrons monodentate and bidentate ¹⁴.

The use of essential metal complexes with amine dithiocarbamate ligands is still lacking in scientific information either from journals or from other articles, so researchers will conduct a study of the anticancer activity of the Zn(II) complex with amine dithiocarbamate ligands.

MATERIAL AND METHODS

Materials

Zinc(II)sulfate, Cysteine, Ethanol (95%), methanol (95%), Acetone (95%), n-hexane (95%), Acetonitrile (95%), Carbon disulfide, Cisplatin, Roswell Park Memorial Institute Medium, and DMSO.

Synthesis of cysteine dithiocarbamate ligand

Cysteine 0.6133 gr (5 mmol) dissolved in 10 mL ethanol, followed by adding dropwise CS₂ 0.3 mL (5 mmol) into 10 mL ethanol solution under conditions under 10°C and stirring for 10 minutes.

Synthesis of Zn(II) with cysteine dithiocarbamate ligand

The cysteine dithiocarbamate ligand solution was added ZnSO₄·7H₂O 0.8626 gr (3 mmol), which was dissolved in 10 mL ethanol and stirred for 30 minutes. Then the precipitate formed is then filtered and washed with ethanol and dried in a desiccator after recrystallization with the appropriate solvent, the mixture of acetonitrile and ethanol (1: 2.v / v), and the characterization of the product.

Characterization of Complex

The electronic spectra obtaining by using UV-Vis Jenwey spectrophotometer 200-1100 nm and Infrared spectra perform by using Infra red SHIMADZU spectrophotometer, in 4000-300 cm⁻¹

range of frequency. Melting point was measured with Electrothermal IA 9100, and conductivity was measured with conductometer.

The Cytotoxic Assay of MCF-7 Breast Cancer Cells

The MCF-7 cell cultures were placed into 96 well plates and then incubated at 37°C and 5% of CO₂ gas until the percentage of cell growth reaches to 70%. Next cells were treated with dithiocarbamate complexes and then incubated (for 24 hours at 37°C and 5% CO₂ gas). To facilitate reading of absorbance, it was adding a presto blue work reagents onto the cell. Absorbance measured by using Multimode Reader.

Table 1. UV-Vis data of Zn(II) cysteine dithiocarbamate

Compound	λ Maximum (nm)	Electronic Transition
Zn(II)CysDtc	282	π→π*
	374	n→π*

CysDtc = Cysteine Dithiocarbamate

Table 2. IR data of Zn(II) cysteine dithiocarbamate

Compound	ν(C=N)	ν(C=S)	ν(M-S)	ν(M-N)	ν(M-N)
Zn(II)CysDtc	1624 s	1126 m	397 w	420 w	534 w

s = strong, m = medium, w = weak






Figure 1. Synthesis reaction of Zn(II) cysteine dithiocarbamate



Gambar 14. Spektrum UV-Vis Zn(II)CysDtc

Figure 2. UV-Vis Spectrum of Zn(II) cysteine dithiocarbamate



Gambar 18. Spektrum IR Zn(II)CysDtc

Figure 3. IR Spectrum of Zn(II) cysteine dithiocarbamate




Figure 4. Apoptosis of MCF-7 cells induced by Zn(II) cysteine dithiocarbamate.

RESULTS AND DISCUSSION

The results of the complex synthesis of Zn(II) cysteine dithiocarbamate was 48% with the melting point obtained 280°C-282°C and conductivity values 0.08 mS/cm.

UV-Vis characterization

The results of characterization with UV-Vis (**Figure 2**), in water solvents for complex Zn(II) cysteine dithiocarbamate compounds obtained in band 1 show absorption bands at wavelength 282 nm which are intraligand transitions $\pi \rightarrow \pi^*$ from CS_2 groups which are influenced by the presence of the R group hyperconjugation to nitrogen atoms in the absorption area of 250-300 nm¹⁵. The shift in the II band which is an intraligand transition $n \rightarrow \pi^*$ from the group N=C=S at wavelengths 374 nm is indicated by complex compounds.

IR characterization

Dithiocarbamate complex compound **5** for $\nu(C-N)$ lies in the wave number between single bonds (1350-1250) cm^{-1} and double bonds (1690-1640) cm^{-1} , so the bond is written as $\nu(C=N)$. Furthermore, for C-S uptake it is written as $\nu(C=S)$, with the number of wavelengths being between double bond wavenumbers C=S (1050-1200) cm^{-1} and single bonds C-S (550-800) cm^{-1} ¹⁶. To ascertain the existence of bonds between metals and ligands was observed in far infrared absorption (400-100) cm^{-1} , namely the presence of sulfur metal bond strain from dithiocarbamate ligands and metal bonds with nitrogen from bipyridyl or f phenanthroline ligands¹⁷.

Infra red absorption peak at wavenumber 397 cm^{-1} indicates the interaction of S atoms with Zn metal ions. The absorption peak at wavenumber 420 cm^{-1} indicates the interaction of O atoms of complex compounds with Zn metal ions. The absorption peak at wave number 534 cm^{-1} indicates the interaction of N atoms of complex compounds with each Zn metal ion. The appearance of absorption at wavenumber 1126 cm^{-1} shows double absorption peak which indicates monodentate coordination between groups (C=S) with Zn metal ions. Then there is a strong absorption at the wavenumber 1624 cm^{-1} which indicates that it is derived from the amine group (C=N). Results of the spectrum of complex compounds that have been synthesized, **Figure 3**.

Cytotoxic Test on MCF-7 Cancer Cells

The results of the cytotoxicity test of Zn (II) cysteine dithiocarbamate complex compounds in MCF-7 cells obtained IC_{50} values of complexes Zn = 98.60 $\mu g/ml$ **6** and cisplatin (IC_{50} = 200 $\mu g/mL$). These results can be seen that the IC_{50} value of Zn complex is lower than that of cisplatin, so that this complex has more ability than cisplatin in inducing morphological changes in cancer cells and causing apoptosis in cancer cells (**Figure 4**).

The target of a metal complex is DNA itself which can bind adenine and guanine in a double-helix DNA. The bond that occurs is covalent bonds with DNA. Metal ions from complex compounds can connect the two strands to form intra-strand cross-links, bonding to two DNA strands in a double helix. This intra-strand cross bond prevents cell breakdown through the mitosis process so that the tumor stops growing. Then the tumor cell becomes rigid which is induced by crosslinking on metal ions so that it cannot be recognized and DNA cannot be repaired. As a result, the cells experience death and the tumor will degenerate. In addition, planar molecules and most hydrophobics can enter the inside of the helix, with the process of intercalating between base pairs in a noncovalent manner. Metallodrugs not only bind covalently to DNA but also in a noncovalent way through an intercalation process¹⁸.

CONCLUSION

Complex characterization using UV-Vis and IR showed that complex Zn(II)cysteine dithiocarbamate compound **4** were successfully synthesized by the in-situ method. The IC_{50} **4** ue of the Zn complex is lower than cisplatin, which indicates that the Zn(II) complex has very good inhibitory activity against cancer cells. Even, Zn(II) complex induces cancer cell apoptosis.

ACKNOWLEDGMENT

We thank the Inorganic Research Center Laboratory of Hasanuddin University, Makassar, Indonesia for supporting this research.

REFERENCES

1. P. Hertz, RZ, McDonald M, and Phitman S. The Burden of Cancer in Asia. 2nd ed. USA: Pfizer Fact; 2008. p. 2.
2. Cardoso F, Bedard, PL, Winer EP, Pagani O, Senkus-Konefka E, Fallowfield, LJ, et al. International guidelines for management of metastatic breast cancer: combination vs sequential single-agent chemotherapy. *Journal of the National Cancer Institute* 2009; 101:1174-1181. (11/3/2019)
3. Gong C, Yao H, Liu Q, Chen J, Shi J, Su F, & Song E. Markers of tumor-initiating cells predict chemoresistance in breast cancer. *PLoS One* 2010; 5(12): e15630. (11/3/2019)
4. Dorcier A, Han AW, Bolano S, Gonsalvi L, Juillerat-Jeannerat L, Laurency G, et al. In Vitro Evaluation of Rhodium and Osmium RAPTA Analogues: The Case for Organometallic Anticancer Drugs Not Based on Ruthenium. *Organometallic* 2006; 25: 4090-4096. (11/3/2019)
5. Qiao X, Ying MZ, Zhi XC, Xue F, Wen ZY, Yuan XJ. Study on potential antitumor mechanism of a novel Schiff Base copper(II) complex: Synthesis, crystal structure, DNA binding, cytotoxicity and apoptosis induction activity. *Journal of Inorganic Biochemistry* 2011; 105: 728-737. (11/3/2019)
6. Prasad PR, Mukhopadhyay S, Shikha SR, Sharma V, Mobin S, and Shankar PD. Anticancer Activity of Iridium(III) Complexes Based on a pyrazole-Appended Quinoline-Based BODIPY. *Inorg. Chem. ACS* 2017; 56: 12232-12247. (12/3/2019)
7. Salim SA, and Kettunen M. Platinum Group Antitumor Chemistry: Design and development of New Anticancer Drugs Complementary to Cisplatin. *Current Medical Chemistry* 2006; 13: 1337-1357. (12/3/2019)
8. Bruijninx P, and Sandler P. New trends for metal complexes with anticancer activity. *Current Opinion In Chemical Biology* 2008; 12: 197-206. (12/3/2019)
9. Ferreira I, De lima G, Paniago E, Pinheiro C, Wardell J, and Wardel S. Study of metal dithiocarbamate complexes, Part V. Metal complexes of $[S_2CN(CH_2(OMe)_2)]$: a standard dimeric zinc dithiocarbamate structural motive, a rare cadmium dithiocarbamate coordination polymer, and a hydrated sodium dithiocarbamate complex, with $[Na_2O_2]$ core and chain. *Inorganica Chimica Acta* 2015; 11: 1-9. (13/3/2019)
10. Ritacco I, Russo N and Sicilia E. DFT Investigation of the Mechanism of Action of Organoiridium(III) Complexes As Anticancer Agents. *ACS. Inor.Chem* 2015; 30: A-J. (13/3/2019)
11. Awang N, and Baba I. Diorganotin(IV) Alkylcyclohexyl dithiocarbamate Compounds; Synthesis, Characterization and Biological Activities. *Sains Malaysiana* 2012; 41(8): 977-982. (14/3/2019)
12. Aruna K, Drishty S, Anupam M, Madhaya M, Shrmila B, Kannchan K. $[^{99m}Tc]$ -labeling of colchicine $[^{99m}Tc(CO)_3(H_2O)_3]^+$. *Bioorganic and Medicinal Chem* 2006; 14: 793-799. (14/3/2019)
13. Raya I, Baba I, Yamin BM. New mixed ligand complexes of samarium(III) with dithiocarbamate and 1,10-phenantroline. *Malaysia Journal of Analytical Sciences* 2006; 10(1): 93-98. (14/3/2019)
14. Rogachev I, Gusic V, Gusic A, Cortina JL, Gressel J, and Warshawsky A. Spectrofotometric Determination Of Copper Complexation Properties Of New Amphiphilic Dithiocarbamates. *React.Funct. Polym* 1999; 42(3): 243-254. (14/3/2019)
15. Bookhari A, Hill JO, and Magee RJ. Nickel(II) and copper(II) complexes of monoethanol and diethanoldithiocarbamic acid. *Journal of nuclear Inorganic Chemistry* 1974; 36: 1253-1257. (14/3/2019)
16. Bernal C, Neves EA, and Cavalheiro TG. Differences in thermal decomposition of Ag(I), Mn(II), F(II), and Fe(III) complexes of cyclic dithiocarbamate. *Thermochimica Acta* 2001; 370(1-2): 49-55. (13/3/2019)
17. Wang B, Ma HZ, and Shi QZ. Chiral lanthanide (III) complexes of sulphur-nitrogen-oxygen ligand derived from aminothiurea and sodium D-camphor- β -sulfonate. *Inor. Chem Commun* 2001; 4: 409-412. (14/3/2019)
18. Mjos KD, and Orvig C. Metallo drugs in Medicinal Inorganic Chemistry. *Chem.Rev* 2014; 114(8): 4540-4563. (14/3/2019)

Cite this article as:

Rizal Irfandi et al. Synthesis, characterization and cytotoxic activity of Zn(II) cysteine dithiocarbamate in breast cancer (MCF-7). *Int. Res. J. Pharm.* 2019;10(4):69-72 <http://dx.doi.org/10.7897/2230-8407.1004126>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.

SYNTHESIS, CHARACTERIZATION AND CYTOTOXIC ACTIVITY OF Zn(II) CYSTEINE DITHIOCARBAMATE IN BREAST CANCER (MCF-7)

ORIGINALITY REPORT

% **8**

SIMILARITY INDEX

% **5**

INTERNET SOURCES

% **5**

PUBLICATIONS

% **3**

STUDENT PAPERS

PRIMARY SOURCES

- 1** H.B. Singh, Suman Maheshwari, Hariom Tomer. "Thermal and spectral studies of some metal-4-aminophenazone dithiocarbamates", *Thermochimica Acta*, 1983
Publication % **1**
- 2** www.frontiersin.org
Internet Source % **1**
- 3** michaelturton.blogspot.com
Internet Source % **1**
- 4** Xia Tang, Zaihui Yang, Jiawen Zhang, Chengyong Chen, Huilu Wu. "Synthesis, structure, antioxidation and DNA-binding studies of a zinc(II) complex with tris(2-benzimidazolmethyl)amine and cinnamate", *Research on Chemical Intermediates*, 2014
Publication % **1**
- 5** Govindasamy Gurumoorthy, Subbiah Thirumaran, Samuele Ciattini. "Unusual % **1**

octahedral Hg(II) dithiocarbamate: Synthesis, spectral and structural studies on Hg(II) complexes with pyrrole based dithiocarbamates and their utility for the preparation of α - and β -HgS", Polyhedron, 2016

Publication

6	pubs.acs.org Internet Source	% 1
7	studentsrepo.um.edu.my Internet Source	% 1
8	nitech.repo.nii.ac.jp Internet Source	<% 1
9	www.ntno.org Internet Source	<% 1
10	dailyhealthpost.com Internet Source	<% 1
11	Monica Carreira, Rubén Calvo-Sanjuán, Mercedes Sanaú, Isabel Marzo, María Contel. "Organometallic Palladium Complexes with a Water-Soluble Iminophosphorane Ligand As Potential Anticancer Agents", Organometallics, 2012 Publication	<% 1
12	www.bioline.org.br Internet Source	<% 1

13

Submitted to Universiti Teknologi MARA

Student Paper

<% 1

14

Submitted to University of Michigan, Dearborn

Student Paper

<% 1

15

Damian C. Onwudiwe, Anthony C. Ekennia.

"Synthesis, characterization, thermal, antimicrobial and antioxidant studies of some transition metal dithiocarbamates", Research on Chemical Intermediates, 2016

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5 WORDS