

THE_CARDIOPROTECTIVE_EFFECT_OF_EXTRA_VIRGIN_OLIVE.pdf

f

by

Submission date: 26-May-2023 07:13PM (UTC+0700)

Submission ID: 2102419721

File name: THE_CARDIOPROTECTIVE_EFFECT_OF_EXTRA_VIRGIN_OLIVE.pdf (537.36K)

Word count: 5753

Character count: 30868

THE CARDIOPROTECTIVE EFFECT OF EXTRA VIRGIN OLIVE OIL AND VIRGIN COCONUT OIL ON ISOPROTERENOL-INDUCED MYOCARDIAL INFARCTION IN RATS

SYAADATUN NADIAH¹, YULIA YUSRINI DJABIR^{2*}, M. ARYADI ARSYAD³, NUR RAHMI²

¹Postgraduate Program, Magister of Pharmacy Program, Faculty of Pharmacy, Hasanuddin University, Makassar, 90245, Indonesia

²Laboratory of Clinical Pharmacy, Faculty of Pharmacy, Hasanuddin University, Makassar, 90245, Indonesia

³Department of Physiology, Faculty of Medicine, Hasanuddin University, Makassar, 90245, Indonesia

*corresponding author: yulia.yusrini@unhas.ac.id

Manuscript received: January 2022

Abstract

Routine consumption of extra virgin olive oil (EVOO) and virgin coconut oil (VCO) may improve cardiovascular function. This study examined the cardioprotective effect of the combination of EVOO and VCO on isoproterenol-induced myocardial infarction in rats. Rats (n = 30) were divided into five groups: a healthy group, an isoproterenol group without treatment, and three treatment groups that were given an EVOO-VCO combination at 1:1, 1:2 and 2:1 (2 mL/200 g) orally for 14 days before injecting isoproterenol (100 mg/kg) for two consecutive days. After 24 hours, blood samples were taken to analyse cardiac biomarkers, and the heart was examined for histopathological changes. Isoproterenol injection caused a substantial increase in the creatine kinase-myocardial band, lactate dehydrogenase and aspartate aminotransferase levels. Pretreatment with an EVOO-VCO combination in all ratios significantly reduced these biomarker levels (p < 0.05). Myocardial damage was significantly reduced, especially with the EVOO-VCO combination of 2:1. In conclusion, the EVOO-VCO combination significantly reduced cardiac injury biomarker levels and improved myocardial structure in isoproterenol-induced myocardial infarction in rats. The EVOO-VCO combination of 2:1 was superior in preserving myocardial structure compared to the other ratios.

Rezumat

Consumul uleiului de măsline extravirgin (EVOO) și a uleiului de cocos virgin (VCO) poate îmbunătăți funcția cardiovasculară. Acest studiu a examinat efectul cardioprotector al combinației de EVOO și VCO asupra infarctului miocardic indus de izoproterenol la șobolani. Șobolanii (n = 30) au fost împărțiți în cinci grupuri: un grup sănătos, un grup de izoproterenol fără tratament și trei grupuri de tratament cărora li s-a administrat o combinație de EVOO-VCO în proporție de 1:1, 1:2 și 2:1 (2 mL/200 g) pe cale orală timp de 14 zile înainte de injectarea de izoproterenol (100 mg/kg) timp de două zile consecutive. După 24 de ore, au fost prelevate probe de sânge pentru a analiza biomarkerii cardiaci, iar la nivelul inimii au fost examinate modificările histopatologice. Administrarea de izoproterenol a provocat o creștere substanțială a nivelurilor de creatin-kinază miocardică, lactat dehidrogenază și aspartat aminotransferază. Pretratarea cu o combinație EVOO-VCO, în toate proporțiile, a redus semnificativ aceste niveluri ale biomarkerilor (p < 0,05). Leziunile miocardice au fost diminuate, în special la administrarea combinației EVOO-VCO 2:1. În concluzie, asocierea EVOO-VCO a redus semnificativ nivelurile biomarkerilor cardiaci și a îmbunătățit structura miocardului în modelul experimental ales. Combinația EVOO-VCO de 2:1 a fost superioară în ceea ce privește conservarea structurii miocardice în comparație cu celelalte proporții.

Keywords: extra virgin olive oil, virgin coconut oil, cardioprotective, isoproterenol, myocardial infarction

Introduction

Isoproterenol (ISO) is a non-selective β -adrenergic (β -1 and β -2) agonist drug [1]. The injection of ISO in high doses has been used in experimental myocardial infarction (MI) in rat models [2, 3]. ISO can trigger an increase in calcium levels in the myocardium, resulting in overstimulation, increased contractile force, oxygen demand and excessive depletion of ATP, leading to myocardial injury [2, 4]. In addition, ISO also cause an increase in oxidative stress, such as reactive oxygen species (ROS) and lipid peroxidase (LPO) [2, 5]. In most coronary artery diseases, a prolonged lack of

oxygen in the myocardium leads to necrosis, which initiates the formation of MI [6].

Lifestyle changes and routine consumption of healthy foods such as vegetables, fruits and antioxidant-rich plants can help improve cardiovascular function [7]. Extra virgin olive oil (EVOO) and virgin coconut oil (VCO) have high fatty acids (FAs) and antioxidants [8]. VCO contains capric acid, caproic acid and lauric acid, which have potential as anti-thrombogenic, anti-arthritis, antihyperlipidemic, cardioprotective, anti-osteoporosis, anti-inflammatory, antimicrobial, hepatoprotective and neuroprotective agents [8]. EVOO

contains FAs that are rich in oleic acid in the form of triacylglycerols and vitamin E [8]. It has potential antimicrobial, antioxidant and anti-inflammatory effects [9]. EVOO at a dose of 10 g/day can reduce the risk of cardiovascular disease by up to 10% [10]. A clinical trial reported that consuming 10 - 50 mL per day of EVOO could significantly reduce blood pressure [11]. Additionally, a preclinical study reported that a single administration of VCO had a cardioprotective effect against cardiac remodelling and blood pressure elevation [12]. The protective effects of EVOO and VCO in combination were shown to be superior against oxidative stress in multiple organs in doxorubicin-treated rats [13]. The study showed that the combination could prevent the development of myocardial infarction after doxorubicin injection [14]. Accordingly, the present study aimed to examine the protective effect of an EVOO and VCO combination in a ratio of 1:1, 1:2 and 2:1 in ISO-treated rats.

Materials and Methods

Chemicals and drugs

EVOO and VCO were purchased from a local pharmacy in Makassar, Indonesia. The identification of the chemical compounds in the oils was by gas chromatography-mass spectrophotometry (GC-MS). Ethanol 70%, diethyl ether, 20% formalin and ISO (USP Cat. from Sigma Chemical Co.) were purchased from official chemical distributors in Jakarta, Indonesia. Creatine kinase-myocardial band (CK-MB), lactate dehydrogenase (LDH), alanine aminotransferase (ALT) and aspartate aminotransferase (AST) kit reagents (Human, Germany) were obtained from the official distributor for Human World Diagnostic in Indonesia.

Chemical component analysis

EVOO and VCO components were analysed using a Trace 1310 gas chromatograph with a TSQ 8000 Evo mass spectrometer (Thermo Scientific; Mundelein, IL, USA). The column size used was 20 mm x 0.18 mm (TG-5MS), with helium as a carrier. The initial temperature of the oven was 50°C, increasing to 330°C at the end.

Preparation of animals

Male albino rats (*Rattus norvegicus*) were used as the animal model. Thirty rats weighing 180 - 300 g were cared for in plastic cages and given standard feed and drinking water ad libitum. The research was conducted in accordance with Directive 2010/63/EU of the European Parliament and of the Council on the protection of animals used for scientific purposes, and the experimental protocol was approved by the research ethics committee of the Faculty of Medicine at Hasanuddin University, Indonesia, (number 671/UN4.6.4.5.31/PP36/22).

Preparation of isoproterenol and EVOO-VCO combination

The dose of isoproterenol used was 100 mg/kg body weight (20 mg/200 g body weight of rats) dissolved in 1 mL of 0.9% NaCl. The combination of EVOO and VCO was made in ratios of 1:1, 1:2 and 2:1, with an oral administration volume of 10 mL/kg rat body weight.

Experimental protocol

The rats were divided into five groups of six: (1) a healthy control group, (2) an ISO group (without any pre-treatment) and (3 - 5) the treatment groups, which received pre-treatment with EVOO and VCO in ratios of 1:1, 1:2 and 2:1, respectively. The treatments were given for 14 days, followed by ISO 100 mg/kg subcutaneous (s.c.) injection for 2 days in a row. After 24 hours following the last ISO injection, blood samples were drawn to analyse the levels of serum biomarkers for MI: CK-MB, LDH, AST and ALT. The blood samples were centrifuged at 3,000 rpm for 10 minutes to obtain serum. The obtained serum was prepared according to the instructions from the reagent kit to analyse the levels of CK-MB, LDH, AST and ALT using a spectrophotometry instrument (Humalyzer 3500). Surgical procedures were performed to remove the heart. Once retrieved, the hearts were rinsed in 0.9% NaCl, drained and weighed. The ratio of the organ weight to the body weight of rats was calculated to obtain the relative organ weight.

Histopathological examination

The rats' hearts were fixed in 10% formalin. After 48 hours, the tissue was vertically cut, processed in a tissue processor for 12 hours, and prepared into paraffin blocks. The tissue was sliced 4 - 5 µm thick using a microtome and floated in a water bath. The sections were placed on glass slides and dried. After 2 hours, staining was performed using haematoxylin and eosin. Histopathological examination was carried out using a light microscope (Olympus®), and photomicrographs were taken with a magnification of 200x. The histopathology score was used to obtain semi-quantitative data for the intensity of myocardial injury. The scores were determined by an anatomical pathologist who was blinded to the treatments given to the rats. A score of 0 indicated no damage in the observed tissue, a score of 1 indicated mild injury (< 25%), a score of 2 indicated moderate damage (26 - 50%), a score of 3 indicated severe damage (51 - 75%) and a score of 4 indicated massive damage (> 75%) [15].

Statistical analysis

The normality of the data was determined using Shapiro-Wilk analysis. The biomarker levels were compared using one-way ANOVA followed by Tukey's HSD test. The histopathological data were compared using the Kruskal-Wallis and Mann-Whitney U tests to determine significant differences between groups. The data are presented as mean ± standard deviation

(SD). The level of statistical significance was set at $p < 0.05$.

Results and Discussion

Chemical components

The GC-MS analysis of the EVOO used in this study revealed 55 peaks of chemical compounds (Table I). FAs, squalene and α -tocopherol were also found. The VCO had 49 peaks (Table II). Like EVOO, VCO is high in FAs. Based on the GC-MS analysis, the FA

content of the EVOO used in this study consisted of 46 g-chain fatty acids (LCFAs) (Table III). Of these, monounsaturated fatty acids (MUFAs) and polyunsaturated fatty acids (PUFAs) were higher in EVOO than in VCO. In VCO, no SCFA content was found, but the MCFA content was much higher than that of EVOO. The LCFA content was lower in VCO than in EVOO. Different from EVOO, the VCO had a high SFA content compared to MUFAs and PUFAs.

84 **Table I**
The chemical compounds of the extra virgin olive oil used based on GC-MS analysis

No.	RT.	RM.	Component	Synonym	Rel. Area (%)
1.	3.772 - 4.068	C ₃ H ₈ O ₃	Gliserin	Gliserin	0.06190
2.	3.955	C ₈ H ₁₄ O ₂	Pentanoic acid, 2-propenyl ester	Velaric acid	0.01697
3	4.241	C ₅ H ₈ O ₂ S	4-Oxopentanethioic acid	4-Oxopentanethioic acid	0.00108
4.	4.891	C ₉ H ₁₈ O ₂	Octanoic acid, methyl ester	Caprylic acid, methyl ester	0.00999
5.	6.173	C ₁₁ H ₂₂ O ₂	Decanoic acid, methyl ester	Capric acid methyl ester	0.00979
6.	7.452	C ₁₅ H ₂₄	α -farnesene	Trans,trans- α -Farnesene	0.00258
7.	7.54 - 7.615	C ₁₃ H ₂₆ O ₂	Dodecanoic acid, methyl ester	Lauric acid methyl ester	0.08249
8.	8.802 - 8.86	C ₁₅ H ₂₈ O ₂	Methyl myristoleate	Cis-9-tetradecanoic acid methyl ester	0.01155
9.	8.914 - 9.026	C ₁₅ H ₃₀ O ₂	Methyl tetradecanoate	Tetradecanoic acid, methyl ester	0.09688
10.	9.424 - 9.591	C ₁₉ H ₃₆ O ₂	9-Octadecanoic acid (Z)-, methyl ester	Oleic acid, methyl ester, cis-	0.07185
11.	9.747	C ₁₆ H ₃₂ O ₂	Pentadecanoic acid, methyl ester	Lauric acid	0.01121
12.	10.159 - 10.213	C ₁₇ H ₃₂ O ₂	9-Hexadecanoic acid, methyl ester, (Z)-	Palmitoleic acid methyl ester	1.05613
13.	10.295 - 10.608	C ₁₇ H ₃₄ O ₂	Hexadecanoic acid, methyl ester	Palmitic acid, methyl ester	28.1459
14.	10.832 - 11.064	C ₁₈ H ₃₄ O ₂	Cis-10-Heptadecenoic acid, methyl ester	Cis-10-Heptadecenoic acid, methyl ester	0.72335
15.	11.247 - 11.287	C ₁₈ H ₃₆ O ₂	Heptadecanoic acid, methyl ester	Margaric acid, methyl ester	0.40585
16.	11.206 - 11.353	C ₁₈ H ₃₆ O ₂	Hexadecanoic acid, 14-methyl-, methyl ester	14-methylhexadecanoic acid methyl ester	0.04777
17.	11.414	C ₁₉ H ₃₄ O ₂	9,12-Octadecadienoic acid (Z,Z)-, methyl ester	Linoleic acid methyl ester	0.14758
18.	11.462 - 11.523	C ₁₉ H ₃₆ O ₂	11-Octadecenoic acid, methyl ester	trans-Vaccenic acid methyl ester	5.08255
19.	11.56 - 12.067	C ₁₉ H ₃₆ O ₂	9-Octadecenoic acid, methyl ester, (E)-	Methyl trans-oleate	13.79887
20.	11.679 - 12.839	C ₁₉ H ₃₆ O ₂	9-Octadecenoic acid (Z)-, methyl ester	Oleic acid, Methyl ester, Cis-	37.19101
21.	12.308	C ₁₉ H ₃₈ O ₂	Methyl stearate	Stearic acid, methyl ester	3.14399
22.	12.455	C ₂₀ H ₃₈ O ₂	Cis-10-Nonadecanoic acid, methyl ester	Cis-10-Nonadecanoic acid, methyl ester	0.29706
23.	12.54	C ₂₀ H ₄₀ O ₂	Nonadecanoic acid, methyl ester	Methyl nonadecanoate	0.08822
24.	12.921	C ₁₉ H ₃₆ O ₃	Oxiranic acid, 3-octyl-, methyl ester, cis-	cis-9,10-Ethoxystearic Acid, methyl ester	0.04413
25.	12.995	C ₂₁ H ₄₀ O ₂	Cis-11-Eicosenoic acid, methyl ester	Eicosenoic acid methyl ester, 11-(Z)-	1.70871
26.	13.118 - 13.149	C ₂₁ H ₄₂ O ₂	Eicosanoic acid, methyl ester	Methyl cis-11-eicosenoate	2.26083

No.	RT.	RM.	Component	Synonym	Rel. Area (%)
27.	13.213	C ₂₅ H ₄₂ O ₂	Cyclopropanebutanoic acid, 2-[[2-[(2-pentylcyclopropyl) methyl] cyclopropyl] methyl] cyclopropyl methyl]-, methyl ester	Butanoic acid	0.0109
28.	13.298	C ₂₀ H ₃₈ O ₂	Cyclopropanoic acid, 2-octyl-, methyl ester	Methyl dihydrosterculate	0.049981
29.	13.39	C ₂₀ H ₃₆ O ₂	Linoleic acid ethyl ester	Linoleic acid ethyl ester	0.09103
30.	13.444	C ₂₀ H ₃₆ O ₂	Methyl 2-octylcyclopropene-1-octanoate	Sterculic acid methyl ester	0.0725
31.	13.567	C ₁₉ H ₃₆ O ₂	10-Octadecenoic acid, methyl ester	10-Octadecenoic acid, methyl ester	0.00591
32.	13.676	C ₁₉ H ₃₆ O ₂	16-Octadecenoic acid, methyl ester	16-Octadecenoic acid, methyl ester	0.00484
	14.873				
	15.305				
33.	13.73	C ₂₂ H ₄₄ O ₂	Heneicosanoic acid, methyl ester	Methyl heneicosanoate	0.09611
34.	14.084	C ₂₁ H ₄₀ O ₄	9-Octadecenoic acid (Z)-, 2-hydroxy-1-(hydroxymethyl) ethyl ester	2-Glyceryl monooleate	0.10305
35.	14.135	C ₂₃ H ₄₀ O ₃	11-(3,4-Dimethyl-5-pentyl-2-furyl)-dodecanoic acid, methyl ester	11-(3,4-Dimethyl-5-pentyl-2-furyl)-dodecanoic acid, methyl ester	0.00577
36.	14.169	C ₁₉ H ₃₆ O ₃	Methyl 5-oxo-octadecanoate	5-Oxostearic acid methyl ester	0.02064
37.	14.257	C ₂₃ H ₄₄ O ₂	Docosenoic acid, methyl ester, (Z)-	Erucic acid methyl ester	0.01851
38.	14.308	C ₁₉ H ₃₈ O ₄	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester	2-Palmitoylglycerol	0.01476
39.	14.434	C ₂₃ H ₄₆ O ₂	Docosanoic acid, methyl ester	Behenic acid methyl ester	0.73223
40.	14.614	C ₁₉ H ₃₆ O ₂	14-Octadecenoic acid, methyl ester	14-Octadecenoic acid, methyl ester	0.00862
41.	15.101	C ₂₄ H ₄₈ O ₂	Tricosanoic acid, methyl ester	Methyl tricosanoate	0.15563
42.	15.795	C ₂₅ H ₅₀ O ₂	Tetracosanoic acid, methyl ester	Lignoceric acid methyl ester	0.35066
43.	16.254	C ₂₇ H ₅₂ O ₄ Si ₂	9,12,15-Octadecatrienoic, 2,3-bis-[(trimethylsilyl)oxy] propyl ester, [Z,Z,Z]-	Monolinolenin TMS	0.00233
44.	16.51 - 16.57	C ₃₀ H ₅₀	Squalene	Squalene	3.46622
45.	16.91	C ₃₅ H ₇₀	Pentatriacontene	17-Pentatriacontene	0.01202
46.	17.121	C ₂₇ H ₅₄ O ₂	Hexacosanoic acid, methyl ester	Cerotic acid methyl ester	0.04682
47.	17.189 - 17.689	C ₂₆ H ₄₄ O ₅	Ethyl iso-allocholate	Cholic acid ethyl ester	0.01695
48.	17.322 - 21.066	C ₂₇ H ₅₆ O ₄ Si ₂	1-Monolinoleylglycerol trimethylsilyl ether	Monoolein TMS	0.03142
49.	18.689	C ₂₉ H ₅₀ O ₂	(+)- α -Tocopherol	(+)- α -Tocopherol	0.05603
50.	19.658	C ₂₈ H ₄₈ O	Campesterol	Campesterol	0.00841
51.	20.593	C ₂₉ H ₅₀ O	Sitosterol	Angelicin (Steroid)	0.19757
52.	20.764	C ₃₀ H ₅₀ O	Cholest-5-en-3-ol, 24-propylidene-, (3 β)-	(E)-24-propylidenecholesterol	0.01523
53.	21.002	C ₃₀ H ₅₀ O ₂	Betulin	Betulin	0.00586
54.	21.417	C ₃₂ H ₅₂ O ₃	9,19-Cyclolanost-24-en-3-ol, acetat, (3 β)-	9,19-Cyclolanost-24-en-3-ol, acetat, (3 β)-	0.00854
55.	22.24	C ₃₀ H ₄₈ O ₅	9,19-Cyclolanost-24-en-3-ol, methylene-, (3 β)-	Cimigol	0.06169
Total					100

82

Table II

The chemical compound of the virgin coconut oil used is based on GC-MS analysis

No.	RT.	RM.	Component	Synonym	Rel. Area (%)
1.	3.456 - 3.806	C ₇ H ₁₄ O ₂	Hexanoic acid, methyl ester	Caproic acid methyl ester	1.54681
2.	3.942 - 4.262	C ₆ H ₁₂ O ₂	Hexanoic acid	Caproic acid	0.25573
3.	4.911 - 5.02	C ₉ H ₁₈ O ₂	Octanoic acid, methyl ester	Caprylic acid methyl ester	3.55331
4.	5.211 - 5.84	C ₈ H ₁₆ O ₂	Octanoic acid	Caprylic acid	5.55028
5.	6.221 - 6.292	C ₁₁ H ₂₂ O ₂	Decanoic acid, methyl ester	Capric acid methyl ester	4.31038

No.	RT.	RM.	Component	Synonim	Rel. Area (%)
6.	6.615 - 7.326	C ₁₀ H ₂₀ O ₂	Decanoic acid	Capric acid	4.16892
7.	7.605 - 8.016	C ₁₃ H ₂₆ O ₂	Dodecanoic acid, methyl ester	Lauric acid methyl ester	12.06083
8.	8.054 - 8.918	C ₁₂ H ₂₄ O ₂	Dodecanoic acid	Lauric acid	18.47322
9.	8.993 - 9.319	C ₁₅ H ₃₀ O ₂	Methyl tetradecanoate	Miristic acid methyl ester	9.49689
10.	9.435 - 10.101	C ₁₄ H ₂₈ O ₂	Tetradecanoic acid	Miristic acid	8.79295
11.	10.224	C ₁₈ H ₃₄ O ₂	Heptadecanoic acid	Elaidic acid	0.02006
12.	10.346 - 10.618	C ₁₇ H ₃₄ O ₂	Hexadecanoic acid, methyl ester	Palmitic acid methyl ester	7.48533
13.	10.642 - 10.982	C ₁₆ H ₃₂ O ₂	n-Hexadecanoic acid	Palmitic acid	3.14541
14.	11.02	C ₁₈ H ₃₆ O ₂	Octadecanoic acid	Stearic acid	0.531309
	12.105				
	12.717				
15.	11.482	C ₁₉ H ₃₆ O ₂	9,12-Octadecadienoic acid (Z,Z)-, methyl ester	Linoleic acid methyl ester	0.02808
16.	11.547 - 11.564	C ₁₉ H ₃₆ O ₂	9-Octadecanoic acid (Z)-, methyl ester	Oleic acid methyl ester	3.58107
	11.72 - 11.741				
17.	11.618	C ₁₉ H ₃₆ O ₂	9-Octadecanoic acid, methyl ester, (E)	Elaidic acid	3.11044
	11.669				
18.	11.632	C ₁₉ H ₃₆ O ₂	11-Octadecanoic acid, methyl ester	Trans veccanic acid methyl ester	1.30384
19.	11.771	C ₁₅ H ₃₀ O ₄	Dodecanoic acid, 2,3-dihydroxypropyl ester	Laurin 1-mono	1.12757
20.	11.822 - 11.907	C ₁₈ H ₃₆ O ₂	Methyl stearate	Stearic acid methyl ester	4.34627
21.	11.921	C ₁₈ H ₃₂ O ₂	9,12-Octadecadienoic acid (Z,Z)-	Linoleic acid	0.72384
22.	11.975	C ₁₈ H ₃₄ O ₂	9-Octadecanoic acid, (E)-	Trans-oleic acid	0.42989
23.	11.996 - 12.383	C ₁₈ H ₃₄ O ₂	Oleic acid	Oleic acid	1.01164
	13.155				
	13.431				
24.	12.785	C ₁₉ H ₃₆ O ₃	Methyl 5 oxo-octadecanoate	Methyl 5 oxo-octadecanoate	0.00843
25.	12.843	C ₁₉ H ₃₆ O ₃	9-Octadecenoic acid, 12-hydroxy-, methyl ester, [R-(Z)]-	Ricinoleic acid methyl ester	0.01373
26.	12.887	C ₂₁ H ₄₀ O ₂	Cis-11-Eicosenoic acid, methyl ester	Cis-11-Eicosenoic acid, methyl ester	0.132
27.	12.945 - 13.019	C ₁₇ H ₃₄ O ₄	Tetradecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester	Miristic acid beta monoglyceride	0.69048
28.	13.37	C ₁₃ H ₂₆ O ₆	Xylitol, 1-O-octanoyl	Octanoil-o-xylitol	0.04892
29.	13.06	C ₂₁ H ₄₂ O ₂	Heptacosanoic acid, methyl ester	Arracidic acid methyl ester	0.23068
30.	14.067	C ₂₂ H ₄₆ O ₃ Si	Hexadecanoic acid, 3-(trimethylsilyloxy) propyl ester	3-[trimethylsilyloxy) propyl palmitat	0.00585
31.	14.247	C ₁₉ H ₃₈ O ₄	Octadecanoic acid, 9,10-dihydroxy-, methyl ester	Methyl 9,10 dihydroxystearate	0.01784
32.	14.325	C ₁₉ H ₃₈ O ₄	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester	Palmitic acid, beta monoglyceride	0.2536
33.	14.407	C ₁₉ H ₃₈ O ₄	Docosanoic acid, methyl ester	Behenic acid methyl ester	0.05816
34.	14.601	C ₁₃ H ₂₆ O ₄	Decanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester	Decanoic acid 2 monoglyceride	0.13775
	14.73				
35.	14.665	C ₁₅ H ₃₀ O ₄	Docosanoic acid, 2,3-dihydroxypropyl ester	Laurin acid alpha monoglyceride	0.18336
	14.764				
36.	14.863	C ₃₅ H ₆₈ O ₅	Hexadecanoic, 1-(hydroxymethyl)-1,2-ethenediyl ester	Dipalmitin	0.00583
	15.448				
37.	15.087	C ₂₄ H ₄₈ O ₂	Methyl 21-methyldocosanoat	Methyl 21-methyldocosanoat	0.00514
38.	15.298	C ₁₄ H ₂₆ F ₂ O ₂	2,2-Difluoroheptacosanoic acid	2,2-Difluoroheptacosanoic acid	0.00425
39.	15.591	C ₂₁ H ₄₀ O ₄	9-Octadecenoic acid (Z)-, 2,3-dihydroxypropyl ester	Olein 2-mono	0.13879
40.	15.74	C ₂₁ H ₄₂ O ₄	Octadecanoic acid, 2,3-dihydroxypropyl ester	Stearin 1-mono	0.02981
41.	15.778	C ₂₅ H ₅₀ O ₂	Tetracosanoic acid, methyl ester	Lignoceric acid methyl ester	0.05691

No.	RT.	RM.	Component	Synonym	Rel. Area (%)
42.	15.968 - 20.781	C ₂₇ H ₅₂ O ₅	Dodecanoic acid, 1-(hydroxymethyl)-1,2-ethanediyl ester	Ethyl laurat	2.77783
43.	16.485	C ₃₉ H ₈₀	Squalene	Squalene	0.01603
44.	17.118	C ₁₈ H ₃₆ O ₂	Hexadecanoic acid, 14-methyl-, methyl ester	Montanic acid methyl ester	0.00466
45.	18.57	C ₂₉ H ₅₄ O ₆	Dinonanoin monocaprylin	Dinonanoin monocaprylin	0.00421
46.	19.682	C ₂₈ H ₄₈ O	Campesterol	Campesterol	0.00237
47.	19.975	C ₂₉ H ₄₈ O	Stigmasterol	Stigmasterol	0.0031
48.	20.339	C ₂₇ H ₅₀ O ₆	Capric acid triglyceride	Caprilic acid trigliseride	0.06873
49.	21.478	C ₂₃ H ₅₂ O ₃	9,19-Cyclolanost-24-en-3-ol, (3β)-	9,19-Cyclolanost-24-en-3-ol, acetate, (3β)-	0.00417
Total					100

Table III

The types of fatty acid content of EVOO and VCO

Types of Fatty Acid				
EVOO	SCFA	0.11%	SFA	35.99%
	MCFA	0.10%	MUFA	60.23%
	LCFA	99.77%	PUFA	3.71%
VCO	SCFA	-	SFA	89.49%
	MCFA	49.92%	MUFA	9.74%
	LCFA	50.07%	PUFA	0.77%

8

SCFA: short-chain fatty acids; MCFA: medium chain fatty acids; LCFA: long-chain fatty acids; SFA: saturated fatty acids; MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids

Relative organ weights

The relative organ weight in the healthy control group was 4.15 ± 0.19 mg/g. The relative heart weight significantly increased in the ISO group, with an average of 6.03 ± 0.51 mg/g (p < 0.0001). Pretreatment with a combination of EVOO and VCO in all ratios maintained the relative organ weights of the rats; hence, the weights were similar to those in the healthy control group (Figure 1).

Biomarker analysis

Four cardiac serum biomarkers were used to confirm the presence of MI: CK-MB, LDH, AST and ALT. In the healthy control group, the mean CK-MB, LDH, AST and ALT levels were 29.59 ± 8.39 U/L, 108.50 ± 26.53 U/L, 105.20 ± 5.86 U/L and 59.55 ± 9.78 U/L, respectively. The levels of these biomarkers, except for ALT, were significantly increased with ISO injection without pretreatment (p < 0.05). In contrast, the CK-MB and LDH levels of rats treated with an EVOO-VCO combination at 1:1, 1:2 and 2:1 were not significantly elevated after ISO injection; the values were not significantly different from that of the healthy control. Furthermore, the EVOO-VCO combination in a 1:2 and 2:1 ratio, but not 1:1, significantly inhibited the increase in AST levels (p < 0.05) (Figure 2).

Histopathological analysis

Representative histopathological features of the heart tissue can be seen in Figure 3, which was scored based on the intensity of injury in Table IV. The healthy control group showed no sign of myocardial tissue damage. In contrast, rats in the ISO group without EVOO-VCO pretreatment had evident

myocardial injury characterized by diffused necrotic areas, haemorrhage and inflammation. In the treatment groups, the administration of EVOO-VCO in any ratio before the injection of ISO reduced the damage to myocardial tissue, but only the ratio of 2:1 led to superior improvement of the myocardial tissue. With EVOO-VCO (2:1), the occurrence of necrosis was only scattered in small areas, with no excessive bleeding, and the infiltration of inflammatory cells was not prominent.




Figure 1.

Comparison of the relative weights of rats' heart among treatment groups

(**) indicates p < 0.001 (***) indicates p < 0.001, (****) indicates p < 0.0001




Figure 2.

Comparison of biomarker levels among rat treatment groups
 a) CK-MB, b) LDH, c) AST and d) ALT; (*) indicate $p < 0.05$, (**) indicate $p < 0.01$, (****) indicates $p < 0.0001$ and (ns) indicates non-significant from the ISO group

Table IV

The score of myocardial damage based on the presence of necrosis, inflammation and haemorrhage

Group	Necrosis	Inflammation	Haemorrhage
Healthy group	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00
ISO group	2.33 ± 0.58	2.33 ± 0.58	1.00 ± 0.00
EVOO:VCO (1:1) + ISO	1.67 ± 0.58	1.33 ± 0.58	1.00 ± 0.00
EVOO:VCO (1:2) + ISO	1.33 ± 0.58	1.33 ± 0.58	0.67 ± 0.58
EVOO:VCO (2:1) + ISO	1.00 ± 0.00	1.00 ± 1.00	0.00 ± 0.00

Score 0: no damage; score 1: a mild injury (< 25%); score 2: moderate damage (26% - 50%); score 3: severe damage (51% - 75%); score 4: massive damage (> 75% - 100%)




Figure 3.

7 Comparison of histopathological features of cardiac tissue

a-b: healthy group; c-d: ISO group 100 mg/kg BW; e-f: group EVOO:VCO ratio 1:1; g-h: group EVOO:VCO 1:2; i-j: group EVOO:VCO 2:1. Red dotted line (necrosis); yellow arrows (inflammation); and green arrows (haemorrhage)

This study was conducted to determine the cardio-protective effect of various combinations (either 1:1, 1:2, or 2:1) of EVOO and VCO in MI rats induced by ISO. Different ratios of EVOO–VCO combinations may result in different cardioprotective effects. Previously, Utari *et al.* (2022) showed that an EVOO–VCO combination in a 1:1 ratio was superior in inhibiting myocardial injury in DOX-induced rats compared to either EVOO or VCO alone. The EVOO–VCO combination was believed to have a synergistic effect in inhibiting cellular damage and death [13]. EVOO and VCO are plant-based oils with very different [68] mical compounds and FA profiles. FAs can be classified based on the length of the carbon chains (SCFAs, MCFAs and LCFAs) [16], or the presence of double bonds in their carbon chains (SFAs, MUFAs and PUFAs) [17]. In this study, EVOO predominantly contained LCFAs, whereas the FA content of VCO primarily consisted of MCFAs and LCFAs. A high LCFA content in EVOO can function as an energy source and maintain heart contractile function [18]. The use of LCFA as the energy source of cardiac metabolism can result in better adenosine triphosphate (ATP) production when compared to glucose. A previous study demonstrated that the incidence of heart failure is more common when energy sources are derived from glucose [19, 20]. The rich MCFA content in VCO, apart from acting as an energy source, [3] n molecularly increase and activate GPCR84 and play a role in the immune system and inflammatory pathways, which results in the inhibition of cAMP [21–23]. Therefore, the initial administration of the EVOO–VCO combination can inhibit the inflammation that occurs in ISO-induced MI rats. In addition, MCFAs such as caprylic acid may act as anti-inflammatory agents, inhibiting the toll-like receptor-4 (TRL4) activation and inhibiting the NF- κ B pathway [24]. By combining EVOO and VCO, the FA content can [66] optimally prevent heart damage and reduce mortality patients with cardiovascular disease [25]. According to a meta-analysis study by Schwingshackl and Hoffmann in 2014, MUFAs may be beneficial to reduce cardiovascular events and death [26]. ISO can cause MI through several mechanisms [2, 27, 28]. The administration of ISO can cause an increase in Ca^{2+} levels in the myocardium, which results in an overstimulation of myocytes, with increased contractility and oxygen demand, causing ATP depletion [5]. The occurrence of excessive intracellular calcium affects the activation of phospholipase and protease enzymes that inhibit Na^+/K^+ ATPase, which causes an increase in intracellular Na^+ and Ca^{2+} , prompting cellular dysfunction and cardiotoxicity [29]. The alteration of myocardial metabolism also triggers an inflammatory reaction. Moreover, infarct-size reducing properties can be related to anti-inflammatory activity of a drug [30]. The neutrophil accumulation in the myocardium

reaches its peak after 24 hours [31]. Monocytes and macrophages predominate in cellular infiltration. In this ph[36] an increase occurs in the production of several cytokines such as IL-1, IL-18, IL-6 and TNF [32]. Inflammatory cells release proteolytic enzymes and ROS, which damage the myocytes. MI is characterized by irreversible damage to cardiomyocytes and results in necrosis [18]. In this study, MI was noticeable after two injections of ISO at the dose of 100 mg/kg, characterized by moderate to severe necrosis in a large area of myocardium. The histological structures of cardiomyocytes were significantly different in the hearts of the rats receiving EVOO–VCO treatment at a ratio of 2:1. Although mild injury was still present, the areas of necrosis and inflammation were significantly reduced. The occurrence of myocardial injury triggers the release of cardiac biomarkers to the systemic circulation. Hence, CK-MB, LDH and AST were significantly elevated after ISO injection in the ISO group [27]. Based on the results of this study, the FA contained in the EVOO–VCO combination could inhibit cellular damage, presumably by providing an energy source that can be effectively metabolized to preserve myocardial energy when ISO induces overstimulation of beta-receptors. Another explanation may come from the anti-inflammatory effect of EVOO and VCO [20, 21], which helps reduce the pro-inflammatory response [24]. Interestingly, the 2:1 ratio elicited better protection against MI. This may indicate that the LCFA and MUFA content of EVOO was required in a higher proportion than the MCFAs contained in VCO. EVOO is also known to contain more antioxidants compared to VCO, which may also explain this result.

Conclusions

The combination of EVOO and VCO [2] ratios of 1:1, 1:2 and 2:1 inhibited the increase in the levels [44] CK-MB, LDH and AST, indicating a cardio-protective effect against ISO-induced MI. In addition, the combination of EVOO and VCO, especially in a ratio of 2:1, provided superior protection against myocardial necrosis and inflammation. Further studies [86] needed to measure the prooxidant and antioxidant levels in rats treated with ISO and EVOO–VCO to confirm [18] whether this cardioprotective effect is related to its antioxidant activity.

Conflict of interest

The authors declare no conflict of interest.

References

1. Neumar RW, Otto CW, Link MS, Kronick SL, Shuster M, Callaway CW, Kudenchuk PJ, Ornato JP, McNally B, Silvers SM, Passman RS, White RD, Hess EP, Tang W, Davis D, Sinz E, Morrison LJ, Part 8: adult advanced cardiovascular life support:

- 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*, 2010; 122(18 Suppl): S729-S767.
2. Ferg M, Khanna D, Exploration of pharmacological interventions to prevent isoproterenol-induced myocardial infarction in experimental models. *Ther Adv Cardiovasc Dis.*, 2014; 8(5): 155-169.
 3. Shaik AH, Rasool SN, Kareem MA, Krushna GS, Akhtar PM, Devi KL, Maslinic acid protects against isoproterenol-induced cardiotoxicity in albino Wistar rats. *J Med Food*, 2012; 15(8): 741-746.
 4. Szymanski MW, Singh DP, Isoproterenol. [Updated 2022 Sep 27]. In: StatPearls, Treasure Island (FL): StatPearls Publishing, 2022; 1-15.
 5. Zhang GX, Kimura S, Nishiyama A, Shokoji T, Rahman M, Yao L, Nagai Y, Fujisawa Y, Miyatake A, Abe Y, Cardiac oxidative stress in acute and chronic isoproterenol-infused rats. *Cardiovasc Res.*, 2005; 65(1): 230-238.
 6. Ojha S, Nandave M, Arora S, Arya DS, Effect of isoproterenol on tissue defense enzymes, hemodynamic and left ventricular contractile function in rats. *Indian Biochem.*, 2010; 25(4): 357-361.
 7. Gilliss JG, Clark AM, Ernst E, Hufford CD, Johnston GAR, Rimoldi JM, Weimar BJ, Natural and synthetic substances related to human health (IUPAC Technical Report). *Pure Appl Chem.*, 2002; 74(10): 1957-1985.
 8. Kamri AM, Ningsih R, Wiyani L, Wiyani Latri, Rahman S, Sistematis review: The effect of virgin coconut oil as a source of antioxidant. *IJRAR.*, 2021; 8(4): 213-219.
 9. Yubero-Serrano EM, Lopez-Moreno J, Gomez-Delgado F, Lopez-Miranda J, Extra virgin olive oil: More than a healthy fat. *Eur J Clin Nutr.*, 2019; 72: 8-17.
 10. Guasch-Ferré M, Hu FB, Martínez-González MA, Fitó M, Bulló M, Estruch R, Ros E, Corella D, Recondo J, Gomez-Gracia E, Fiol M, Lapetra J, Serra-Majem L, Muñoz MA, Pintó X, Lamuela-Raventós RM, Basora J, Buil-Cosiales P, Sorlí JV, Ruiz-Gutiérrez V, Martínez JA, Salas-Savado J, Olive oil intake and risk of cardiovascular disease and mortality in the PREDIMED Study. *BMC Med.*, 2014; 12(78): 1-11.
 11. Zamora-Zamora F, Martínez-Galiano JM, Gaforio JJ, Delgado-Rodríguez M, Effects of olive oil on blood pressure: A systematic review and meta-analysis. *Grasas y Aceites*, 2018; 69(4): 1-9.
 12. Kamisah Y, Periyah V, Lee KT, Noor-Izwan N, Nurul-Hamizah A, Nurul-Iman BS, Subermaniam K, Jaarin K, Azam A, Faizah O, Qodriyah HMS, Cardioprotective effect of virgin coconut oil in heated palm oil diet-induced hypertensive rats. *Pharm Biol.*, 2015; 53(9): 1243-1249.
 13. Utari AU, Djabir YY, Palinggi BP, A combination of virgin coconut oil and extra virgin olive oil elicits superior protection against doxorubicin cardiotoxicity in rats. *Turk J Pharm Sci.*, 2022; 19(2): 138-144.
 14. Rus I, Tertiş M, Paşcalău V, Pavel C, Melean B, Suciuc M, Moldovan C, Topală T, Popa C, Săndulescu R, Cristea C, Simple and fast analytical method for the evaluation of the encapsulation and release profile of doxorubicin from drug delivery systems. *Farmacia*, 2011; 69(4): 670-681.
 15. Gibson-Corley KN, Olivier AK, Meyerholz DK, Principles for valid histopathologic scoring in research. *Pathol.*, 2013; 50(6): 1007-1015.
 16. Yin CL, Nijholt KT, Bakker BM, Westenbrink KH, Short-Chain fatty acids in the Metabolism of heart failure – rethinking the fat stigma. *Front Cardiovasc Med.*, 2022; 9: 1-9.
 17. Savchenko L, Ivanauskas L, Jarukas L, Georgiyants V, Determination and comparison of fatty acids composition of apricot and peach oils. *Farmacia*, 2021; 69(5): 941-947.
 18. Labarthe F, Gélinas R, Rosiers CD, Medium-chain fatty acids as metabolic therapy in cardiac disease. *Cardiovasc Drugs Ther.*, 2008; 22: 97-106.
 19. Angelini A, Saha PK, Jain A, Jung SY, Mynatt RL, Pi X, Xie L, PHDs/CPT1B/VDAC1 axis regulates long-chain fatty acid oxidation in cardiomyocytes. *Cell Rep.*, 2021; 37(1): 1-14.
 20. Lopaschuk GD, Ussher JR, Folmes CDL, Jaswal JS, Stanley WC, Myocardial fatty acid metabolism in health and disease. *Physiol Rev.*, 2010; 90(1): 209-258.
 21. Chopashree PG, Shetty SS, Kumari SN, Effect of medium chain fatty acid in human health and disease. *J Funct Foods*, 2021; 87: 1-11.
 22. Toit ED, Browne L, Irving-Rodgers H, Massa HM, Fozzard N, Jennings MP, Peak IR, Effect of GPR84 deletion on obesity and diabetes development in mice fed long chain or medium chain fatty acid rich diets. *Eur J Nutr.*, 2018; 57: 1737-1746.
 23. Wang J, Wu X, Simonavicius N, Tian H, Ling L, Medium-chain fatty acids as ligands for orphan G protein-coupled receptor GPR84. *JBC.*, 2006; 281(84): 34457-34464.
 24. Zhang X, Xue C, Xu Q, Zhang Y, Li H, Li F, Liu Y, Guo C, Caprylic acid suppresses inflammation via TLR4/NF-κB signaling and improves atherosclerosis in ApoE-deficient mice. *Nutr Metab (Lond.)*, 2019; 16(40): 1-16.
 25. Balta I, Stef L, Pet I, Iancu T, Stef D, Corcionivoschi N, Essential Fatty Acids as Biomedicines in Cardiac Health. *Biomedicines*, 2021; 9(10): 1-24.
 26. Schwingshackl L, Hoffmann G, Monounsaturated fatty acids, olive oil and health status: a systematic review and meta-analysis of cohort studies. *Lipids Health Dis.*, 2014; 13: 1-15.
 27. Afroz R, Tanvir EM, Karim N, Hossain MS, Alam N, Gan SH, Khalil MI, Sundarban Honey Confers Protection against Isoproterenol-Induced Myocardial Infarction in Wistar Rats. *Biomed Res Int.*, 2016; 2016: 1-10.
 28. Mohamed ME, Abduldaium MS, Younis NS, Cardioprotective effect of linalool against isoproterenol-induced myocardial infarction. *Life*, 2021; 11(2): 1-17.
 29. Djabir Y, Dobson GP, Hemodynamic rescue and ECG stability during chest compressions using adenosine and lidocaine after 8-minute asphyxial hypoxia in the rat. *Am J Emerg Med.*, 2013; 31(11): 1539-1545.
 30. Ramos K, Combs AB, Acosta D, Role of calcium in isoproterenol cytotoxicity to cultured myocardial cells. *Biochem Pharmacol.*, 1984; 33(12): 1989-1992.

31. Kain V, Halade GV, Role of neutrophils in ischemic heart failure. *Pharmacol Ther.*, 2020; 205: 1-22.
32. Turillazzi E, Pomara C, Bello S, Neri M, Riezzo I, Fineschi V, Meaning of different forms of structural myocardial injury, immune response and timing of infarct necrosis and cardiac repair. *Curr Vasc Pharmacol.*, 2015; 13(1): 6-19.

ORIGINALITY REPORT

24%

SIMILARITY INDEX

18%

INTERNET SOURCES

19%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	9lib.net Internet Source	1%
2	cms.turkjps.org Internet Source	1%
3	www.frontiersin.org Internet Source	1%
4	Gupta, Mahesh, Pallavi Sharma, Arindam Ghosh Mazumder, Vikram Patial, and Damanpreet Singh. "Dwindling of cardio damaging effect of isoproterenol by Punica granatum L. peel extract involve activation of nitric oxide-mediated Nrf2/ARE signaling pathway and apoptosis inhibition", Nitric Oxide, 2015. Publication	1%
5	eprints.undip.ac.id Internet Source	1%
6	smujo.id Internet Source	1%

7	<p>Manjunatha S., Althaf Hussain Shaik, Maruthi Prasad E., Suliman Yousef Al Omar, Altaf Mohammad, Lakshmi Devi Kodidhela.</p> <p>"Combined cardio-protective ability of syringic acid and resveratrol against isoproterenol induced cardio-toxicity in rats via attenuating NF-kB and TNF-α pathways", Scientific Reports, 2020</p> <p>Publication</p>	1 %
8	<p>ijponline.biomedcentral.com</p> <p>Internet Source</p>	1 %
9	<p>pubmed.ncbi.nlm.nih.gov</p> <p>Internet Source</p>	1 %
10	<p>Yannai, . "Name Index", Dictionary of Food Compounds with CD-ROM Second Edition, 2012.</p> <p>Publication</p>	1 %
11	<p>assets.researchsquare.com</p> <p>Internet Source</p>	<1 %
12	<p>Risa Kashiwagi-Takayama, Junji Kozawa, Yoshiya Hosokawa, Sarasa Kato et al.</p> <p>"Myocardial fat accumulation is associated with cardiac dysfunction in patients with type 2 diabetes, especially in elderly or female patients: a retrospective observational study", Cardiovascular Diabetology, 2023</p> <p>Publication</p>	<1 %

13	mobt3ath.com Internet Source	<1 %
14	repositorio.udec.cl Internet Source	<1 %
15	www.chemspider.com Internet Source	<1 %
16	www.phenomenex.com Internet Source	<1 %
17	Submitted to De Montfort University Student Paper	<1 %
18	www.jbiomeds.com Internet Source	<1 %
19	ijms.sums.ac.ir Internet Source	<1 %
20	Yusof Kamisah, Vengadesh Periyah, Kee Tat Lee, Norrashid Noor-Izwan et al. "Cardioprotective effect of virgin coconut oil in heated palm oil diet-induced hypertensive rats", <i>Pharmaceutical Biology</i> , 2015 Publication	<1 %
21	M. I. Cengiz. "Periodontal disease in patients with familial Mediterranean fever: from inflammation to amyloidosis", <i>Journal of Periodontal Research</i> , 06/2009 Publication	<1 %

22

Submitted to Universitas Hasanuddin

Student Paper

<1 %

23

Takashi Sano, Ryo Okabe, Maiko Iwahashi, Jun Imagi, Toshiro Sato, Toshiyuki Yamashita, Eiichiro Fukusaki, Takeshi Bamba. "Effect of Furan Fatty Acids and 3-Methyl-2,4-nonanedione on Light-Induced Off-Odor in Soybean Oil", Journal of Agricultural and Food Chemistry, 2017

Publication

<1 %

24

Nur Arfian, Danny A. P. Wahyudi, Ingesti B. Zulfatina, Arsitya N. Citta et al. "Chlorogenic Acid Attenuates Kidney Ischemic/Reperfusion Injury via Reducing Inflammation, Tubular Injury, and Myofibroblast Formation", BioMed Research International, 2019

Publication

<1 %

25

origin-www.fishersci.com

Internet Source

<1 %

26

www.jomes.org

Internet Source

<1 %

27

Zhipeng Liu, Rui Liu, Jing Chou, Jiaying Yu, Xiaowei Liu, Changhao Sun, Ying Li, Liyan Liu. "Targeted metabolomics analysis reveals the association between maternal folic acid supplementation and fatty acids and amino

<1 %

acids profiles in rat pups", Journal of
Chromatography B, 2018

Publication

28

Li, Hua, Yan-Hua Xie, Qian Yang, Si-Wang Wang, Bang-Le Zhang, Jian-Bo Wang, Wei Cao, Lin-Lin Bi, Ji-Yuan Sun, Shan Miao, Jing Hu, Xuan-Xuan Zhou, and Peng-Cheng Qiu. "Cardioprotective Effect of Paeonol and Danshensu Combination on Isoproterenol-Induced Myocardial Injury in Rats", PLoS ONE, 2012.

Publication

29

Submitted to University of Queensland

Student Paper

30

pubag.nal.usda.gov

Internet Source

31

elifesciences.org

Internet Source

32

isfj.ir

Internet Source

33

www.chemicalbook.com

Internet Source

34

Submitted to University of Bradford

Student Paper

35

repositorio.ufsc.br

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

36

www.internetchemie.info

Internet Source

<1 %

37

www.ocl-journal.org

Internet Source

<1 %

38

www.slideshare.net

Internet Source

<1 %

39

Yan-Ling Liu, Chun-Yan Kong, Peng Song, Heng Zhou, Xing-Sheng Zhao, Qi-Zhu Tang. "Maslinic acid protects against pressure overload-induced cardiac hypertrophy in mice", *Journal of Pharmacological Sciences*, 2018

Publication

<1 %

40

git.embl.de

Internet Source

<1 %

41

webbook.nist.gov

Internet Source

<1 %

42

www.molecularinfo.com

Internet Source

<1 %

43

"Final Report on the Safety Assessment of Ricinus Communis (Castor) Seed Oil, Hydrogenated Castor Oil, Glyceryl Ricinoleate, Glyceryl Ricinoleate SE, Ricinoleic Acid, Potassium Ricinoleate, Sodium Ricinoleate, Zinc Ricinoleate, Cetyl Ricinoleate, Ethyl Ricinoleate, Glycol Ricinoleate, Isopropyl

<1 %

Ricinoleate, Methyl Ricinoleate, and Octyldodecyl Ricinoleate", International Journal of Toxicology, 2007

Publication

44

Basant M. Al-Botaty, Abeer Elkhoely, Elsayed K. El-Sayed, Amany A.E. Ahmed. "Ethyl pyruvate attenuates isoproterenol-induced myocardial infarction in rats: Insight to TNF- α -mediated apoptotic and necroptotic signaling interplay", International Immunopharmacology, 2022

Publication

<1 %

45

Liu, X.. "Antioxidant activity and phenolics of an endophytic Xylaria sp. from Ginkgo biloba", Food Chemistry, 2007

Publication

<1 %

46

Pei Dai, Zhongxiang Tang, Mingming Qi, Dan Liu, Ousman Bajinka, Yurong Tan. "Dispersion and utilization of lipid droplets mediates respiratory syncytial virus - induced airway hyperresponsiveness", Pediatric Allergy and Immunology, 2021

Publication

<1 %

47

Romana Ahmed, E. M. Tanvir, Md. Sakib Hossen, Rizwana Afroz et al. "Antioxidant Properties and Cardioprotective Mechanism of Malaysian Propolis in Rats", Evidence-

<1 %

Based Complementary and Alternative Medicine, 2017

Publication

48

academicjournals.org

Internet Source

<1 %

49

publication.babcock.edu.ng

Internet Source

<1 %

50

www.acc.org

Internet Source

<1 %

51

www.freepatentsonline.com

Internet Source

<1 %

52

www.gjesm.net

Internet Source

<1 %

53

www.koreascience.or.kr

Internet Source

<1 %

54

www.wjgnet.com

Internet Source

<1 %

55

Alexandra Gyongyosi, Rita Zilinyi, Andras Czegledi, Agnes Tosaki, Arpad Tosaki, Istvan Lekli. "The Role of Autophagy and Death Pathways in Dose-dependent Isoproterenolinduced Cardiotoxicity", Current Pharmaceutical Design, 2019

Publication

<1 %

56

C. SAIZ-JIMENEZ. "Land use effects on the composition of organic matter in soil particle

<1 %

size separates. III. Analytical pyrolysis",
European Journal of Soil Science, 3/1996

Publication

57

Dalia Ismail, Asmaa ShamsEldeen, Laila Rashed, Ashraf Shama, Sara Ashour, Alshaymaa Aboulkhair. "Cardioprotective Potential of Zinc and Vitamin E Against Isoprenaline-Induced Myocardial Infarction in Albino Rats by Targeting Autophagy: A Histological and Biochemical Study", Egyptian Journal of Histology, 2020

Publication

<1 %

58

Holly Susan Haggerty, Hugh S. Lusted. "Histological Reaction to Polyimide Films in the Cochlea", Acta Oto-Laryngologica, 2009

Publication

<1 %

59

Kais Mnafgui, Raouf Hajji, Fatma Derbali, Anis Gammoudi et al. "Anti-inflammatory, Antithrombotic and Cardiac Remodeling Preventive Effects of Eugenol in Isoproterenol-Induced Myocardial Infarction in Wistar Rat", Cardiovascular Toxicology, 2015

Publication

<1 %

60

Sumayya Shahzad, Somaiya Mateen, P.M. Mubeena Mariyath, Syed Shariq Naeem, Kafil Akhtar, Waseem Rizvi, Shagufta Moin. "Protective effect of syringaldehyde on

<1 %

biomolecular oxidation, inflammation and histopathological alterations in isoproterenol induced cardiotoxicity in rats", Biomedicine & Pharmacotherapy, 2018

Publication

61

Yulia Yusrini Djabir, Aryadi Arsyad, Mufidah Murdifin, Rosany Tayeb et al. "Kleinhovia hospita extract alleviates experimental hepatic and renal toxicities induced by a combination of antituberculosis drugs", Journal of Herbmed Pharmacology, 2020

Publication

<1 %

62

academic.oup.com

Internet Source

<1 %

63

bestjournal.untad.ac.id

Internet Source

<1 %

64

doczz.net

Internet Source

<1 %

65

m.biomart.cn

Internet Source

<1 %

66

mspace.lib.umanitoba.ca

Internet Source

<1 %

67

old.21food.com

Internet Source

<1 %

68

patents.google.com

Internet Source

<1 %

69	sciendo-parsed-data-feed.s3.eu-central-1.amazonaws.com Internet Source	<1 %
70	storage.googleapis.com Internet Source	<1 %
71	ur.booksc.eu Internet Source	<1 %
72	wap.guidechem.com Internet Source	<1 %
73	worldwidescience.org Internet Source	<1 %
74	www.ehrenstorfer.com Internet Source	<1 %
75	www.potravinarstvo.com Internet Source	<1 %
76	www.researchsquare.com Internet Source	<1 %
77	www.tandfonline.com Internet Source	<1 %
78	Anna Gawron-Skarbek, Agnieszka Guligowska, Anna Prymont-Przymińska, Dariusz Nowak, Tomasz Kostka. "The Anti-Inflammatory and Antioxidant Impact of Dietary Fatty Acids in Cardiovascular Protection in Older Adults May	<1 %

Be Related to Vitamin C Intake", Antioxidants, 2023

Publication

79

S. A. Salami, A. Guinguina, J. O. Agboola, A. A. Omede, E. M. Agbonlahor, U. Tayyab. "Review: and effects of feed antioxidants in livestock: a review of the implications on authorization of antioxidant feed additives ", animal, 2016

Publication

80

Zhibo Fu, Hongyang Zhang, Zhijun Zeng, Fanghong Ning, Ziwei Xu, Chenyu Liu, Min Zhang, Ping Hu. "A pre - column derivatization high - performance liquid chromatography method for simultaneous determination of short - chain and medium - chain fatty acids in a fecal sample", Journal of Separation Science, 2022

Publication

81

Dong, Tao, Jun Wang, Chao Miao, Yubin Zheng, and Shulin Chen. "Two-step in situ biodiesel production from microalgae with high free fatty acid content", Bioresource Technology, 2013.

Publication

82

Kamisah, Yusof, Shu-Min Ang, Faizah Othman, Badlishah Sham Nurul-Iman, and Hj Mohd Saad Qodriyah. "Renoprotective effect of

<1 %

<1 %

<1 %

<1 %

virgin coconut oil in heated palm oil diet-induced hypertensive rats", Applied Physiology Nutrition and Metabolism, 2016.
Publication

83

Krivacsy, Z.. "Primary organic pollutants in New Zealand urban aerosol in winter during high PM¹⁰ episodes", Environmental Pollution, 200601

Publication

<1 %

84

Quintero-Flórez, Angélica, Laura Sinausia Nieva, Araceli Sánchez-Ortiz, Gabriel Beltran, and Javier S. Perona. "The Fatty Acid Composition of Virgin Olive Oil from Different Cultivars is Determinant for Foam Cell Formation by Macrophages", Journal of Agricultural and Food Chemistry

Publication

<1 %

85

Samah M. Ahmed, Shaimaa Ali Abdelrahman, Alaa Elsayed Salama. "Efficacy of gold nanoparticles against isoproterenol induced acute myocardial infarction in adult male albino rats", Ultrastructural Pathology, 2017

Publication

<1 %

86

Sudip Paul, Sagarika Das, E.M. Tanvir, Md. Sakib Hossen et al. "Protective effects of ethanolic peel and pulp extracts of Citrus macroptera fruit against isoproterenol-

<1 %

induced myocardial infarction in rats", Biomedicine & Pharmacotherapy, 2017

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On