

Renal Resistive Index in Hypertensive Patients: One Centre Study

by Hasyim Kasim

Submission date: 30-Apr-2023 03:51AM (UTC+0700)

Submission ID: 2079332055

File name: I_Resistive_Index_in_Hypertensive_Patients-_One_Centre_Study.pdf (616.37K)

Word count: 2973

Character count: 15269

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 2449-6170

e-ISSN: 2449-6162

Renal Resistive Index in Hypertensive Patients: One Centre Study

Authors: Hasyim Kasim, Khadijah Khairunnisa Hasyim, Andi Makbul Aman, Dimas Bayu, Nur Fitriani

DOI: 10.5603/AH.a2023.0003

30
Article type: Case report

Submitted: 2022-12-23

Accepted: 2023-01-23

Published online: 2023-01-24

2
This article has been peer reviewed and published immediately upon acceptance.
It is an open access article, which means that it can be downloaded, printed, and distributed freely,
provided the work is properly cited.
The final version may contain major or minor changes.

41

Renal resistive index in hypertensive patients: one centre study

10.5603/AH.a2023.0003

Hasyim Kasim, Khadijah Khairunnisa Hasyim, Andi Makbul Aman, Dimas Bayu, and Nur Fitriani

8

Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Tamalanrea, Makassar, Indonesia

8

Corresponding author: Hasyim Kasim, Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Tamalanrea, Makassar, Indonesia; e-mail: hasyimkasim.unhas@gmail.com

Abstract

31

Background: Hypertension (HTN) is a leading cause of kidney dysfunction. Renal resistive index (RRI) was an index to evaluate arterial compliance and/or resistance, reflecting the reduction of kidney function and microalbuminuria. We investigated the relationship of RRI in hypertensive patients to detect kidney dysfunction early detection.

Material and methods: This was a cross-sectional study at Wahidin Sudirohusodo hospital in June–November 2022. All hypertensive patient was evaluated for RRI. RRI was examined with intrarenal doppler ultrasound, and a cut-off ≥ 0.70 were used.

Results: This study included 61 subjects. Thirty-five subjects were female, and 26 subjects were male 90.2% of subjects were below 60 years. Estimated glomerular filtration rate (eGFR) level was 90.29 ± 25.19 in $RRI < 0.7$ and 64.91 ± 31.79 in $RRI \geq 0.7$. Our study found there was a significant difference between anti-hypertensive treatment and eGFR level with the RRI group (p -value < 0.05). There was no significant difference in sex, age, proteinuria, and HTN control status in both RRI groups.

Conclusion: The renal resistive index is a useful marker for early renal dysfunction in hypertensive patients despite normal eGFR.

Key words: hypertension; renal resistive index (RRI); eGFR level; proteinuria

Introduction

Renal resistive index (RRI) is an ultrasonographic Doppler measurement of flow velocities in intraparenchymal renal arteries [1]. It is a non-invasive and repeatable method

for assessing arterial compliance and/or resistance. RRI appears to have a significant role in assessing various secondary hypertension (HTN) patients. RRI is related to subclinical indicators of target organ damage and represents renal disease progression beyond albuminuria and creatinine clearance. Also, the RRI can evaluate cardiovascular and renal risk [2]. Several studies indicate that this index reflects systemic hemodynamic and depends on the aortic pulse pressure, which is affected by parameters like age, presence of HTN, or diabetes. In patients with widespread atherosclerosis or reduced vascular compliance, RRI may be increased even with normal kidney function [2]. An elevated RRI (≥ 0.70) is usually associated with impaired renal function, increased proteinuria, and poor prognosis [3]. Evaluation of RRI may also contribute to therapeutic decision-making. Given its straightforward assessment, RRI appears as a simple approach and “multifunctional” instrument that might aid in evaluating renal disease progression. The purpose of this review was to evaluate RRI in hypertensive patients.

Material and methods

Subjects

Sixty-one patients with HTN at Wahidin Sudirohusodo Hospital were chosen as a subject. All patients had signed a consent form and confirmed their voluntary participation in this research study. They were given an explanation regarding the purpose, benefits, and what was done in this study and agreed to participate in this research voluntarily. During the study, they had been given the right to ask questions or ask for clarification from researchers if there were still things that were not clear.

All essential HTN patients were evaluated for renal RI at the initial visit. HTN was defined as systolic blood pressure of 140 mm Hg and/or diastolic blood pressure of 90 mm Hg, measured three times in the sitting position using a brachial sphygmomanometer or therapy with antihypertensive medication. The exclusion criteria were chronic kidney disease with dialysis. The ethics committee of the Faculty of Medicine, Hasanuddin University, authorized the study with ethical number UH22090548. The study was conducted in accordance with the Declaration of Helsinki Ethical Principles and Good Clinical Practices.

Blood and urine collection

Using an automated analyser, haemoglobin, creatinine, sodium, potassium, and chloride were determined. The estimated glomerular filtration rate was computed utilizing the equation of Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI).

Duplex doppler ultrasonography

Doppler ultrasonography was used to investigate renal hemodynamics utilizing an **HI VISION Avius (Hitachi Aloka Medical, Tokyo, Japan.)** and a **3.5-MHz convex probe** fitted with a Doppler system. Doppler flow was measured in the interlobar arteries of both kidneys at three distinct places (superior, middle, and inferior) using colour flow mapping as a reference. Then, peak systolic velocity (PSV) and end-diastolic velocity (EDV) were calculated. The average resistive index (RI) was computed using the following formula:

$$RI = (PSV - EDV)/PSV [4].$$

Statistical analysis

Statistical analysis was implemented using the statistical package for social sciences (SPSS) software, version 25.0 for Windows. Data are expressed as mean \pm SD or median (interquartile range). Both data and normality were analysed using the Kolmogorov-Smirnov test. The chi-square test was used to evaluate significant differences between variable with normal data distribution. Statistical significance was defined as p-value < 0.05.

Results

This study included 61 patients (Tab. 1 and 2). Thirty-five subjects were female, and 26 subjects were male. 90.2% of subjects were below 60 years. Antihypertensive treatment usage was calcium channel blocker (50.8%), angiotensin receptor blocker (9.8%), angiotensin converting enzyme inhibitor (ACEI) (6.6%), a combination of calcium channel blocker (CCB) and angiotensin receptor blocker (ARB) (27.9%), a combination of CCB and ACEI (3.3%), and three combination drugs (1.6%). Based on the RRI group, 88.5% had the RRI value of < 0.7, and 11.5% had an RRI value of \geq 0.7. Based on proteinuria, 42.6% had proteinuria and 57.4% had no proteinuria. Based on hypertensive control, accounted 31.1% of subjects had controlled HTN, and 68.9% of subjects were uncontrolled. In Table 3, the average eGFR RRI < 0.7 is 90.29 ± 25.19 and the average RRI \geq 0.7 is 64.91 ± 31.79 . With a p-value of 0.030, there was a significant difference between eGFR and RRI. Age, systolic, diastolic, pulse rate, haemoglobin, creatinine, and potassium were not significantly different on the RRI (p-value > 0.05). In the RRI 0.7 group, there were 27 CCB subjects, 6 ARB subjects, 4 ACEI subjects, 14 CCB+ARB subjects, 2 CCB + ACEI subjects, and 1 person

receiving a combination of all three medicines. In the RRI > 0.7 group, 4 participants had CCB, 0 had ARB, 0 had ACEI, 3 had CCB + ARB, 0 had CCB + ACEi, and 0 had a combination of all 3 medications. There was a correlation between antihypertensive therapy and RRI ($p = 0.04$) (Tab. 4).

Discussion

This study showed a significant relationship between RRI values and eGFR even though the average eGFR value was more than 60 mL/min/1.73 m². Renal vasodilating capacity was reduced before the onset of established renal damage and in normal RRI values, meaning that functional rather than structural alterations might already be present, indicating a subclinical stage of renal damage. In predicting Chronic Kidney Disease (CKD) progression and poor outcomes in cases with mild to moderate renal impairment, RRI was superior to renal function assessment alone [5]. Four RRI has been linked to a quicker loss in renal function in individuals with proteinuria CKD or diabetics with microalbuminuria, even when GFR levels are normal [6]. In Table 3, there was a significant relationship between RRI and eGFR in 61 hypertensive patients. It seems that lower eGFR values were associated with higher RRI values. A similar relationship was reported by Gaurav et al. who reported that there was a significant negative correlation between RRI and eGFR [7].

This study did not show a significant association with proteinuria. It could be due to the fact that this study did not assess microalbuminuria or proteinuria in 24 hours. Hashimoto et al. studied 133 HTN individuals and found that each 0.1 increase in RRI was related to a 5.4-fold increase in the incidence of albuminuria [8]. In a investigation involving 66 patients with critical HTN, a strong correlation was seen between high RRI and future increases in urine albumin excretion [9].

The only variable that substantially predicted an >50% rise in the urine albumin to creatinine ratio over two years was RRI. The ideal RRI cut-off value that predicted this increase was 0.71 (sensitivity 52.4% and specificity 84.0%). This cut-off value was consistent with other studies indicating that RRI values >0.7 are more prevalent in patients with left ventricular hypertrophy or advanced carotid atherosclerosis and are associated with higher mortality in hypertensive patients with CKD and no clinically significant renal artery stenosis [10, 11].

The assessment of RRI may have therapeutic consequences as well. Our study found there was a significant difference ($p < 0.05$) between RRI and antihypertensive treatment. During chronic antihypertensive medication, there was evidence that changes in RRI parallel

changes in urine albumin excretion [11]. In addition, an increase in RRI indicated intrarenal stiffness and urges care in titrating renin-angiotensin system inhibitors to prevent renal function decline, particularly in CKD patients, diabetics, and the elderly. In particular, Renin angiotensin system inhibitors (RASi) such as valsartan and lisinopril can improve renal function in individuals with essential HTN, particularly those with microalbuminuria, by decreasing renal vascular resistance and so avoiding eventual renal failure [12, 12].

Details on the effect of treatment and RRI are given in Table 5. Overall, the majority of patients have RRI of less than 0.7. RRI values were affected by the type of therapy received by either the controlled or uncontrolled HTN. While monotherapy therapy is effective for controlled and uncontrolled HTN, combined therapy is less effective for uncontrolled therapy. Despite the treatment, none showed statistically significant ($p < 0.05$).

Conclusion

RRI is a useful marker for renal dysfunction in hypertensive patients.

Authorship

H.K.: conceptualization, data collection, writing, funding acquisition. K.K.H.: data collection, methodology. A.M.A.: methodology, writing. D.B.: data modelling. N.F.: writing — review and editing.

Conflict of interest

The authors declare no conflict of interest.

Funding

We want to thank the Hasanuddin University, Indonesia for their research grand number 1148/UN4.22/PT.01.06/2022.

Acknowledgments

We would like to thank the Department of Internal Medicine, Wahidin Sudirohusodo Hospital, Makassar, Indonesia for their support.

References

1. Viazzi F, Leoncini G, Derchi LE, et al. Ultrasound Doppler renal resistive index: a useful tool for the management of the hypertensive patient. *J Hypertens*. 2014; 32(1): 149-153, doi: [10.1097/HJH.0b013e328365b29c](https://doi.org/10.1097/HJH.0b013e328365b29c), indexed in Pubmed: [24172238](https://pubmed.ncbi.nlm.nih.gov/24172238/).
2. Andrikou I, Tsioufis C, Konstantinidis D, et al. Renal resistive index in hypertensive patients. *J Clin Hypertens (Greenwich)*. 2018; 20(12): 1739-1744, doi: [10.1111/jch.13410](https://doi.org/10.1111/jch.13410), indexed in Pubmed: [30362245](https://pubmed.ncbi.nlm.nih.gov/30362245/).
3. Giustiniano E, Pugliese L, Calabrò L, et al. Renal Resistive Index: Not Only for Kidney Clinics. *J Ultrasound Med*. 2022; 41(4): 1027-1030, doi: [10.1002/jum.15775](https://doi.org/10.1002/jum.15775), indexed in Pubmed: [34213804](https://pubmed.ncbi.nlm.nih.gov/34213804/).
4. Viazzi F, Leoncini G, Derchi LE, et al. Ultrasound Doppler renal resistive index: a useful tool for the management of the hypertensive patient. *J Hypertens*. 2014; 32(1): 149-153, doi: [10.1097/HJH.0b013e328365b29c](https://doi.org/10.1097/HJH.0b013e328365b29c), indexed in Pubmed: [24172238](https://pubmed.ncbi.nlm.nih.gov/24172238/).
5. Lubas A, Kade G, Niemczyk S. Renal resistive index as a marker of vascular damage in cardiovascular diseases. *Int Urol Nephrol*. 2014; 46(2): 395-402, doi: [10.1007/s11255-013-0528-6](https://doi.org/10.1007/s11255-013-0528-6), indexed in Pubmed: [23959401](https://pubmed.ncbi.nlm.nih.gov/23959401/).
6. Afsar B, Elsurur R. Increased renal resistive index in type 2 diabetes: Clinical relevance, mechanisms and future directions. *Diabetes Metab Syndr*. 2017; 11(4): 291-296, doi: [10.1016/j.dsx.2016.08.019](https://doi.org/10.1016/j.dsx.2016.08.019), indexed in Pubmed: [27594114](https://pubmed.ncbi.nlm.nih.gov/27594114/).
7. Gaurav K, Yalavarthy U, Chamberlain N, et al. Correlation between Renal Resistive Index and Estimated Glomerular Filtration Rate in Patients with Hypertension. *J Vasc Ultrasound*. 2018; 32(2): 82-84, doi: [10.1177/154431670803200204](https://doi.org/10.1177/154431670803200204).
8. Hashimoto J, Ito S. Central pulse pressure and aortic stiffness determine renal hemodynamics: pathophysiological implication for microalbuminuria in hypertension. *Hypertension*. 2011; 58(5): 839-846, doi: [10.1161/HYPERTENSIONAHA.111.177469](https://doi.org/10.1161/HYPERTENSIONAHA.111.177469), indexed in Pubmed: [21968753](https://pubmed.ncbi.nlm.nih.gov/21968753/).
9. Shimizu Y, Itoh T, Hougaku H, et al. Clinical usefulness of duplex ultrasonography for the assessment of renal arteriosclerosis in essential hypertensive patients. *Hypertens Res*. 2001; 24(1): 13-17, doi: [10.1291/hyres.24.13](https://doi.org/10.1291/hyres.24.13), indexed in Pubmed: [11213024](https://pubmed.ncbi.nlm.nih.gov/11213024/).
10. Toledo C, Thomas G, Schold JD, et al. Renal resistive index and mortality in chronic kidney disease. *Hypertension*. 2015; 66(2): 382-388, doi: [10.1161/HYPERTENSIONAHA.115.05536](https://doi.org/10.1161/HYPERTENSIONAHA.115.05536), indexed in Pubmed: [26077569](https://pubmed.ncbi.nlm.nih.gov/26077569/).
11. Miyoshi K, Okura T, Tanino A, et al. Usefulness of the renal resistive index to predict an increase in urinary albumin excretion in patients with essential hypertension. *J Hum Hypertens*. 2017; 31(1): 66-69, doi: [10.1038/jhh.2016.38](https://doi.org/10.1038/jhh.2016.38), indexed in Pubmed: [27278927](https://pubmed.ncbi.nlm.nih.gov/27278927/).
12. Trovato GM, Martines GF, Trovato FM, et al. Renal resistive index and parathyroid hormone relationship with renal function in nondiabetic patients. *Endocr Res*. 2012; 37(2): 47-58, doi: [10.3109/07435800.2011.625513](https://doi.org/10.3109/07435800.2011.625513), indexed in Pubmed: [22007967](https://pubmed.ncbi.nlm.nih.gov/22007967/).
13. Zhang F, Liu H, Liu Di, et al. Effects of RAAS Inhibitors in Patients with Kidney Disease. *Curr Hypertens Rep*. 2017; 19(9): 72, doi: [10.1007/s11906-017-0771-9](https://doi.org/10.1007/s11906-017-0771-9), indexed in Pubmed: [28791529](https://pubmed.ncbi.nlm.nih.gov/28791529/).

Table 1. Data descriptive

	Minimum	Maximum	Average	SD
Age [years]	32.0	63.0	50.7	8.6
Systolic blood pressure [mm Hg]	110.0	214.0	155.4	21.9
Diastolic blood pressure [mm Hg]	63.0	127.0	89.6	12.2
Heart rate [beat/min]	62.0	121.0	88.1	11.2
Haemoglobin [g/dL]	7.4	16.9	12.8	2.3
Creatinine [mg/dL]	0.4	3.4	0.9	0.5
GFR [mL/min/1.73 m ²]	14.8	138.0	87.4	26.9
RRI	0.5	0.8	0.6	0.1
Sodium [mEq/L]	127.0	150.0	139.3	4.2
Potassium [mEq/L]	2.8	5.3	3.9	0.6
Chloride [mEq/L]	96.0	113.0	104.8	4.0

SD — standard deviation; GFR — glomerular filtration rate; RRI — renal resistive index

Table 2. Data descriptive

Variable	n	%
Sex		
Female	35	57.4
Male	26	42.6
Age		
< 60 years	55	90.2
≥ 60 years	6	9.8
Antihypertensive treatment		
CCB	31	50.8
ARB	6	9.8
ACEI	4	6.6
CCB + ARB	17	27.9
CCB + ACEI	2	3.3
Combination of three	1	1.6
RRI		
< 0.7	54	88.5
≥ 0.7	7	11.5
Proteinuria		
Normal	35	57.4
Proteinuria	26	42.6
Hypertensive control		
Controlled	19	31.1
Uncontrolled	42	68.9

Note: Our study found there was a significant difference between antihypertensive treatment and estimated glomerular filtration rate (eGFR) level with the renal resistive index (RRI) group (p-value < 0.05). CCB — calcium channel blocker ARB — angiotensin receptor blocker; ACEI — angiotensin converting enzyme inhibitor

Table 3. Renal resistive index (RRI) profile in hypertensive patient

Variable	< 0.7 (n = 54)	≥ 0.7 (n = 7)	p-value*
Age	50.0 ± 8.8	55.5 ± 4.1	0.10
Systolic	156.1 ± 21.3	150.1 ± 27.2	0.50
Diastolic	90.5 ± 12.3	83.1 ± 9.9	0.14
Heart rate	88.8 ± 11.3	83.1 ± 9.5	0.21
Haemoglobin	12.7 ± 2.2	13.3 ± 3.1	0.51
Creatinine	0.9 ± 0.3	1.4 ± 0.9	0.10**
eGFR	90.3 ± 25.2	64.9 ± 31.8	0.03**
Potassium	3.9 ± 0.5	4.2 ± 0.5	0.19

*t-test at p < 0.05; **Mann Whitney test, the value was significant at p < 0.05. eGFR — estimated glomerular filtration rate

Table 4. Relationship of renal resistive index (RRI) and independent variable

Variable	RRI (n,%)		p-value*
	< 0,7 (n = 54)	≥ 0,7 (n = 7)	
Sex			
Female	32 (91.4)	3 (8.6)	0.41
Male	22 (84.0)	4 (15.4)	
Age			
< 60 years	49 (89.1)	6 (10.9)	0.67
≥ 60 years	5 (83.3)	1 (16.7)	
Anti-hypertensive treatment			
CCB	27 (87.1)	4 (12.9)	
ARB	6 (100.0)	0 (0)	0.04
ACEI	4 (100.0)	0 (0)	
CCB + ARB	14 (82.4)	3 (17.6)	
CCB + ACEI	2 (100.0)	0 (0)	
Combination of three	1 (100.0)	0 (0)	
Proteinuria			
Normal	29 (82.9)	6 (17.1)	0.10
Proteinuria	25 (96.2)	1 (3.8)	

Hypertensive control			
Controlled	16 (84.2)	3 (15.8)	0.47
Uncontrolled	38 (90.5)	4 (9.5)	
Value (average ± SD)			
Creatinine	0.88 ± 0.32	1.37 ± 0.93	0.10**
GFR	90.29 ± 25.19	64.91 ± 31.79	0.03**

*t-test at $p < 0.05$; **Mann Whitney test, the value was significant at $p < 0.05$;

CCB — calcium channel blocker ARB — angiotensin receptor blocker; ACEI — angiotensin converting enzyme inhibitor; SD — standard deviation; GFR — glomerular filtration rate

Table 5. Renal resistive index (RRI) values for controlled and uncontrolled hypertension patients

Hypertension			RRI		Total	p-value
			< 0.7	≥ 0.7		
Controlled hypertension	Monotherapy	N	12	3	15	0.52
		%	80.0%	20.0%	100.0%	
	Combination therapy	N	7	0	7	
		%	100.0%	0%	100.0%	
Total		N	19	3	22	
		%	86.4%	13.6%	100.0%	
Uncontrolled hypertension	Monotherapy	N	26	1	27	0.08
		%	96.3%	3.7%	100.0%	
	Combination therapy	N	9	3	12	
		%	75.0%	25.0%	100.0%	
Total		N	35	4	39	
		%	89.7%	10.3%	100.0%	

The value was significant at $p < 0.05$

Renal Resistive Index in Hypertensive Patients: One Centre Study

ORIGINALITY REPORT

35%

SIMILARITY INDEX

30%

INTERNET SOURCES

30%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1 pubmed.ncbi.nlm.nih.gov Internet Source 2%

2 repozytorium.ukw.edu.pl Internet Source 2%

3 www.ncbi.nlm.nih.gov Internet Source 2%

4 Daijiro Enomoto, Takafumi Okura, Tomoaki Nagao, Masanori Jotoku, Jun Irita, Ken-ichi Miyoshi, Jitsuo Higaki. "Relationship Between Renal Hemodynamics and Urinary Type IV Collagen in Patients with Essential Hypertension", Clinical and Experimental Hypertension, 2012
Publication 2%

5 journals.sagepub.com Internet Source 2%

6 www.dovepress.com Internet Source 2%

repo.dma.dp.ua

7	Internet Source	1 %
8	www.neuroquantology.com Internet Source	1 %
9	Lotta Gränse. "Electrophysiology and ocular blood flow in a family with dominant optic nerve atrophy and a mutation in the OPA1 gene", <i>Ophthalmic Genetics</i> , 12/1/2003 Publication	1 %
10	insights.ovid.com Internet Source	1 %
11	assets.researchsquare.com Internet Source	1 %
12	journals.lww.com Internet Source	1 %
13	www.researchsquare.com Internet Source	1 %
14	hdl.handle.net Internet Source	1 %
15	radiologykey.com Internet Source	1 %
16	Konstantinos Kintis, Costas Tsioufis, Alexandros Kasiakogias, Kyriakos Dimitriadis et al. "Noninvasive assessment of	1 %

haemodynamics in resistant hypertension",
Journal of Hypertension, 2017

Publication

17

scirp.org

Internet Source

1 %

18

Baris Afsar, Rengin Elsurur. "Increased renal resistive index in type 2 diabetes: Clinical relevance, mechanisms and future directions", Diabetes & Metabolic Syndrome: Clinical Research & Reviews, 2017

Publication

1 %

19

Lubas, Arkadiusz, Grzegorz Kade, and Stanisław Niemczyk. "Renal resistive index as a marker of vascular damage in cardiovascular diseases", International Urology and Nephrology, 2014.

Publication

1 %

20

Nitika Kumari, Natarajan Suresh, Josephine A.. "Significance of Bcl-2 expression in breast cancer", Biomedicine, 2022

Publication

1 %

21

secure.medicalletter.org

Internet Source

1 %

22

Luis Izquierdo, Ana M Rodríguez, Ramón A Sarquis, Diego Altamirano, Maria A Henriquez. "Intracorneal circular ring implant with

1 %

femtosecond laser: Pocket versus tunnel",
European Journal of Ophthalmology, 2021

Publication

23

Marwan Sofyan, Dewi Yunia Fitriani, Dewi Friska, Ray Wagi Basrowi, Ahmad Fuady. "B Vitamins, work - related stress and emotional mental disorders: a cross - sectional study among nurses in Indonesia", Nursing Open, 2022

Publication

1 %

24

ejrnm.springeropen.com

Internet Source

1 %

25

www.cardiologyres.org

Internet Source

1 %

26

www.pjmhsonline.com

Internet Source

1 %

27

www.pubmedcentral.nih.gov

Internet Source

1 %

28

"Clinical Implications", Hypertension, 2015

Publication

<1 %

29

Azar Nickavar, Baranak Safaeian, Hossein Zaeri, Mohammad Hadi Gharib, Tahereh Chaharnaiei. "Usefulness of Doppler ultrasound for the early diagnosis of diabetic nephropathy in type 1 diabetes mellitus", Journal of Ultrasound, 2021

Publication

<1 %

30 Tomasz Sternalski, Natalia Bajorek, Maciej Skubera, Małgorzata Urbańczyk-Zawadzka, Piotr Podolec, Lidia Tomkiewicz-Pająk. "Wielonarządowe powikłania odległe po radioterapii i chemioterapii chłoniaka Hodgkina", Folia Cardiologica, 2022
Publication

31 heart.bmj.com
Internet Source

32 oamjms.eu
Internet Source

33 www.irog.net
Internet Source

34 www.science.gov
Internet Source

35 "Biomarkers in Kidney Disease", Springer Science and Business Media LLC, 2016
Publication

36 Haijing Hu, Rumeng Ye, Lu Pang, Han Jiang, Kai Tian, Yang Gao, Yanli Ji, Pengwei Wan, Junbo Yang, Xiaoming Zou, Xingjun Tian. "After Life Effect of Endophytic Bacterium Bacillus Cereus on Litter Decomposition", Research Square Platform LLC, 2021
Publication

37 bsdwebstorage.blob.core.windows.net
Internet Source

<1 %

38

www.mdpi.com

Internet Source

<1 %

39

Lin, Y.H.. "The influence of estimated creatinine clearance on plasma homocysteine in hypertensive patients with normal serum creatinine", *Clinical Biochemistry*, 200702

Publication

<1 %

40

Ayşe Selcan Koc, Derya Demirtaş, Feride Fatma Gorgulu, Hilmi Erdem Sumbul. "Diurnal variation of renal resistive index over 24-hour period in hypertensive patients and healthy controls", *Abdominal Radiology*, 2018

Publication

<1 %

41

R. L. Kulynych, D. G. Rekalov. "Forecast and clinical-diagnostic significance of increased expression of beta-2microglobulin in the urine in the process of nephropathy formation in essential hypertension", *Journal of Education, Health and Sport*, 2020

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On