

The_Total_Edge_Irregular_Labeling.pdf

by

Submission date: 06-Jun-2023 10:02AM (UTC+0700)

Submission ID: 2109971536

File name: The_Total_Edge_Irregular_Labeling.pdf (420.3K)

Word count: 1319

Character count: 6444

The Total Edge Irregular Labeling of Network Constructed by Some Copies of Cycle on Three Vertices Corona a Vertex

Nurdin*, Jusmawati Massalesse and Besse Bunga Yulandary

Mathematics Department, Mathematics and Natural Sciences Faculty, Hasanuddin University, Makassar, Indonesia;
nurdin1701@unhas.ac.id

Abstract

Many networks have been found the total edge irregularity strength's. The aim of this paper is determined the total edge irregularity strength of network constructed by some copies of cycle on three vertices corona a vertex. The results of this paper that the total edge irregularity strength of network constructed by some copies of cycle on three vertices corona a vertex is $2m + 1$ for $m \geq 2$ where m is the number copies of cycle on three vertices.

Keywords: Corona Product of Graph, Irregular Labeling, Irregularity Strength

1. Introduction

A graph labeling is one of models to illustrate the approach for any network. A graph G consists of a finite set $V(G)$ of objects called vertices together with a set $E(G)$ of an ordered pairs of vertices; the elements of $E(G)$ are called edges. The edge e containing x and y is written $e = xy$, where x and y are called endpoints of e .

Formally, a graph labeling is a map that carries graph elements to the numbers (usually to the positive or non-negative integer). A labeling of a graph is called a vertex labeling, an edge labeling, or a total labeling, if the domain of this map is the vertex set, the edge set, or the union of vertex and edges set respectively¹⁷.

In 2002, Bača *et al.* introduced an edge irregular total k -labeling of any graph. A total labeling $\alpha : V \cup E \rightarrow \{1, 2, \dots, k\}$ is called a total edge irregular k -labeling of G if the weight of all edges is distinct. The weight of edge $e = xy$ under the total labeling α is $\alpha(e) + \alpha(x) + \alpha(y)$, where x and y are vertices in a graph.

For some positive integer k , any graph have a total edge irregular k -labeling. The interesting is the minimum positive integer number k for which G has a total

edge irregular k -labeling and it is called the total edge irregularity strength of G , denoted by $tes(G)$. Bača, *et al.* have determined the total edge irregularity strength of path, cycle, stars, wheels, and friendship graph².

The original motivation for the definition of the total edge irregularity strength came from irregular assignments and the irregularity strength of a graphs introduced in⁸

An irregular assignment for any graph G is a t -labeling of the edges

$$\lambda : E \rightarrow \{1, 2, 3, \dots, t\}$$

such that the weight of all vertices of G are distinct, where E is the edge set of G . The weight of vertex v in G under t -labeling λ is the sum of the label of v and the label of all edges incident to v . The smallest t for which there is an irregular assignment on G is called the irregularity strength of G .

Some results have been obtained for several classes of graphs. For instances, in^{3,4,6} determined the exact value of the irregularity strength of trees. For special kinds of tree, in⁷ have determined the irregularity strength of full d -ary trees.

*Author for correspondence

We are interested in the total edge irregular labeling mainly because of the following intriguing theorem posted by⁵ and conjecture posed by¹⁰.

Theorem A. Let $G = (V, E)$ be a graph with maximum degree Δ . Then

$$tes(G) \geq \max \left\{ \left\lceil \frac{|E(G)|+2}{3} \right\rceil, \left\lceil \frac{\Delta+2}{2} \right\rceil \right\}.$$

Conjecture B. For every graph $G = (V, E)$, different from K_5 , with the maximum degree Δ , then

$$tes(G) = \max \left\{ \left\lceil \frac{|E|+2}{3} \right\rceil, \left\lceil \frac{\Delta+1}{2} \right\rceil \right\}.$$

Many authors have been verified the conjecture. In^{13,14} we have determined the total edge irregularity strength of the corona product of a path with certain graphs and C_3 – snake. Besides that, in¹² have determined the total edge irregularity strength of a categorical product of two paths and certain family of graphs. In¹¹ determined the total edge irregularity strength of complete graphs and complete bipartite graphs. The total edge irregularity strength of the grids has determined by¹². In¹⁵ determined the total edge irregularity strength of subdivision of star. The total edge irregularity strength of the disjoint union of helm graphs have determined by¹⁶.

As main result in this paper, we determined the total edge irregularity strength of corona product of cycle with a vertex.

2. Main Result

The corona product is one of many graph operations in graph theory. Formally, the corona product of a graph G with a graph H , denoted by $G \odot H$, is the graph obtained by taking one copy G and $|V(H)|$ copies of H , and joining the i – th vertex of G with an edge to every vertex in i – th copy of H ¹⁰. Specially, in this paper we study about the corona product of m copies of cycle C_3 with a vertex, denoted by W_3^m .

First, we define the vertex set $V(W_3^m)$ and edge set $E(W_3^m)$ of W_3^m as follows:

$$V(W_3^m) = \{x, x_j^i | i = 1, 2, \dots, m; j = 1, 2, 3\} \text{ and}$$

$$E(W_3^m) = \{x_j^i x_{j+1}^i, x_1^i x_3^i, x_2^i x_3^i | i = 1, 2, \dots, m; j = 1, 2\}.$$

Theorem 1. For $m \geq 2$, $tes(W_3^m) = 2m + 1$.

Proof. By definition of the edge set of W_3^m , the number of edges of W_3^m is $6m + 2$. By Theorem A, we find that $tes(W_3^m) \geq 2m + 1$. To find that $tes(W_3^m) \leq 2m + 1$, $\{x_0 y_{i,j} | i = 1, \dots, m; j = 1, 2\}$, we construct a total edge irregular k -labeling for W_3^m where $k = 2m + 1$. Define a total labeling f for W_3^m as follows:

- 1. $f(x) = 2m + 1$,

- for $i = 1, 2, \dots, m$,

$$f(x_j^i) = \begin{cases} j, & \text{for } j = 1 \\ i, & \text{for } j = 2 \\ m + i, & \text{for } j = 3 \end{cases}$$

- for $i = 1, 2, \dots, m$,

$$f(x_j^i x_{j+1}^i) = \begin{cases} j, & \text{for } j = 1 \\ m + 2 - i, & \text{for } j = 2 \end{cases}$$

- for $i = 1, 2, \dots, m$,

$$f(x_1^i x_3^i) = 1,$$

- for $i = 1, 2, \dots, m$,

$$f(xx_j^i) = \begin{cases} m + i, & \text{for } j = 1 \\ 2m + 1, & \text{for } j = 2, 3 \end{cases}.$$

By the definition of the total labeling f , we find that $f : V \cup E \rightarrow \{1, 2, \dots, k\}$ where $k = 2m + 1$. The find that $f : V \cup E \rightarrow \{1, 2, \dots, k\}$ is the total edge irregular labeling, we have to show that the weight of all edges are distinct. By the definition of f , the weight of all edges of W_3^m as follows:

- for $i = 1, 2, \dots, m$

$$wt(x_1^i x_2^i) = 2 + i,$$

- for $i = 1, 2, \dots, m$

$$wt(x_1^i x_3^i) = m + 2 + i,$$

- for $i = 1, 2, \dots, m$

$$wt(x_2^i x_3^i) = 2m + i + 2,$$

- for $i = 1, 2, \dots, m$

$$wt(xx_1^i) = 3m + 2 + i,$$

- for $i = 1, 2, \dots, m$

$$wt(xx_2^i) = 4m + 2 + i,$$

- for $i = 1, 2, \dots, m$

$$wt(xx_3^i) = 5m + 2 + i.$$

Accordingly, we find that every two edges different in W_3^m for $m \geq 2$, $e_1 \neq e_2$, we have $wt(e_1) \neq wt(e_2)$. This showed that, the total labeling $f : V \cup E \rightarrow \{1, 2, \dots, k\}$, where $k = 2m + 1$, is a total edge irregular labeling for W_3^m . That is $tes(W_3^m) \leq 2m + 1$. \square

3. Conclusion

In this paper we verified that the conjecture posed by Ivančo and Jendrol' is true for the network constructed by some copies of cycle on three vertices corona a vertex.

4. References

1. Ahmad A, Bača M. Total edge irregularity strength of a categorical product of two paths, *Ars Combin.* 2014; 114:203–12.
2. Ahmad A, Bača M. Edge-irregular total labeling of certain family of graphs. *AKCE J Graphs Combin.* 2009; 6(1):21–9.
3. Aigner M, Triesch E. Irregular assignments of trees and forests. *SIAM J Discrete Math.* 1990; 3:439–49.
4. Amar D, Togni O. Irregularity strength of trees. *Discrete Math.* 1998; 190:15–38.
5. Bača M, Jendrol' S, Miller M, Ryan J. On irregular total labellings. *Discrete Math.* 2007; 307:1378–88.
6. Bohman T, Kravitz D. On the irregularity strength of trees. *J Graph Theory.* 2004; 45:241–54.
7. Cammack LA, Schelp RH, Schrag GC. Irregularity strength of full d-ary trees. *Congr Numer.* 1991; 81:113–9.
8. Chartrand G, Jacobson MS, Lehel J, Oellermann OR, Ruiz S, Saba F. Irregular networks. *Congr Numer.* 1988; 64:197–210.
9. Gallian JA. Dynamic survey of Graph labeling. *Electron J Combin.* 2015.
10. Ivančo J, Jendrol' S. Total edge irregularity strength of trees. *Discuss Math.* 2006; 26:449–56.
11. Jendrol' S, Miškuf J, Soták R. Total edge irregularity strength of complete graphs and complete bipartite graphs. *Discrete Math.* 2010; 310:400–7.
12. Miškuf J, Jendrol' S. On total edge irregularity strength of the grids. *Tatra Mt Math Publ.* 2007; 36:147–51.
13. Nurdin. The total edge and vertex irregularity strength of $nC_3 - snake$. *Far East Journal of Mathematical Sciences.* 2013; 78(2):205–10.
14. Nurdin, Salman AMN, Baskoro ET. The total edge-irregular strengths of the corona product of paths with some graphs. *J. Combin, Math Combin Comput.* 2008; 65:163–75.
15. Hinding N, Suardi N, Basir H. Total edge irregularity strength of subdivision of star. *Journal of Discrete Mathematical Sciences and Cryptography.* 2015; 18(6):869–75.
16. Siddiqui MK, Nurdin, Baskoro ET. Total edge irregularity strength of the disjointunion of helm graphs. *J Math Fund Sci.* 2012; 45(2):163–71.
17. Wallis WD. Magic graphs. Boston: Birkhäuser; 2001.

The_Total_Edge_Irregular_Labeling.pdf

ORIGINALITY REPORT

20%
SIMILARITY INDEX

%
INTERNET SOURCES

20%
PUBLICATIONS

%
STUDENT PAPERS

PRIMARY SOURCES

- 1** Akul Rana. "On the k -distant total labeling of graphs", Malaya Journal of Matematik, 2020
Publication **5%**
- 2** Susilawati, , Edy Tri Baskoro, and Rinovia Simanjuntak. "Total Vertex-Irregularity Labelings for Subdivision of Several Classes of Trees", Procedia Computer Science, 2015.
Publication **4%**
- 3** Jendrol, S.. "Total edge irregularity strength of complete graphs and complete bipartite graphs", Discrete Mathematics, 20100206
Publication **3%**
- 4** Dafik, D J Koesoemawati, I H Agustin, E Y Kurniawati, R Nisviasari. "On local vertex irregular reflexive coloring of graphs", Journal of Physics: Conference Series, 2022
Publication **2%**
- 5** Sarbari Mitra, Soumya Bhoumik. "On (a,d) -total edge irregular labelings", Discrete Mathematics, Algorithms and Applications, 2023
Publication **2%**

6

N. Krishnaveni. "Feature Selection Algorithms for Data Mining Classification: A Survey", Indian Journal of Science and Technology, 2019

Publication

1 %

7

Ahmad, Ali, Omar Bin Saeed Al-Mushayt, and Martin Bača. "On edge irregularity strength of graphs", Applied Mathematics and Computation, 2014.

Publication

1 %

8

K. Muthu Guru Packiam, T. Manimaran, A. ThuraiSwamy. "Chapter 23 Irregularity Strength of Corona of Two Graphs", Springer Science and Business Media LLC, 2017

Publication

1 %

9

Martin Bača, Muhammad Imran, Andrea Semaničová-Feňovčíková. "Irregularity and Modular Irregularity Strength of Wheels", Mathematics, 2021

Publication

1 %

Exclude quotes On

Exclude matches < 4 words

Exclude bibliography On