

Faisal_Amir_EEC-64_Artikel.pdf

by

Submission date: 13-Jan-2022 09:47PM (UTC+0700)

Submission ID: 1741133027

File name: Faisal_Amir_EEC-64_Artikel.pdf (465.6K)

Word count: 2707

Character count: 13529

Population dynamics of *Carcharhinus melanopterus* caught from Southern Makassar Strait, Indonesia

Faisal Amir*, Achmar Mallawa and Joeharnani Tresnati

Department of Fisheries, Faculty of Marine and Fisheries, Hasanuddin University, Makassar, 90245, South Sulawesi, Indonesia

30
(Received 15 August, 2020; Accepted 30 October, 2020)

ABSTRACT

21
Blacktip reef shark *Carcharhinus melanopterus* is the species dominant catch of the traditional fisheries using bottom gill nets and bottom longline in South Sulawesi Province, Indonesia. This study aims to determine several *C. melanopterus* population dynamics parameters include growth rate, fishing mortality, exploitation rate, and relative yield per recruitment. Data collection of fish length was carried out from July to September 2020. 28 Data were collected from all captured sharks landed at fish landing sites Paotere and Beba twice a week. Population growth is analyzed using the von Bertalanffy equation exponential growth, the value of L_{∞} , K , by ELEFAN method and t_0 by Pauly method. The total mortality (Z) was estimated from 37 catch curve, and the natural 35 mortality (M) was obtained from Pauly's empirical relationship based on 1 L_{∞} , K , and the mean temperature of the environment. FISAT-II software's help estimated the size of length 14 at first capture (L_c). Finally, estimating the optimum level of exploitation rate (E) was carried out using the relative yield-per-recruit model (Y'/R) of Beverton and Holt. The results showed that the population of 10 *C. melanopterus* in the Makassar Strait's southern waters had a low population growth rate with an infinity 10 length (L_{∞}) of 161.5 cm and a growth rate coefficient (K) of 0.25 yr^{-1} , during the t_0 value of -0.4247 yr. The total mortality (Z) of 1.05 yr^{-1} , the natural mortality (M) of 0.45 yr^{-1} , fishing mortality (F) 0.6 yr^{-1} , and the size of length at first capture (L_c) of 99.64 cm. The estimated rate of exploitation (E) was below the optimum rate of exploitation. It seems that the population of *C. melanopterus* in this area is under-exploited.

Key words : Blacktip reef shark, Population dynamics, Makassar Strait

Introduction

2
Blacktip reef sharks *Carcharhinus melanopterus* are commonly found throughout tropical coral reef and coastal habitats (Chin *et al.*, 2013a). It is one type of reef fish that has a high economic value in Indonesia, so that it is suspected that there has been a decline in the condition of its stock. Shark fisheries in the southern Makassar Strait have not been used commercially, but because the high demand for shark exports has changed, the shark turns out to be the primary target. Sharks are exploited throughout the year using traditional fishing tools such as bot-

tom long line and bottom gill net without a management policy. Although these fishing activities keep continuous, there is still very little research to reveal this problem.

The lower catch per unit effort (CPUE) indicates that shark fisheries resources have decreased (Fowler and Cavanagh, 2005). The growth of shark fisheries in Indonesia has exceeded production limits. This is indicated by the increasing difficulty of local fishers to catch sharks because the fishing locations are further away, the number of catches decreases, and the size that is caught is getting smaller (Fahmi and Dharmadi, 2013).

*Corresponding author email: faisalamir_unhas@yahoo.com

Shark's status is a potential change from vulnerable to overfishing. In order to recover the number of shark in one area, it takes an extended period. It is caused by some of the biological characteristics of sharks is a long life cycle reaching a maximum size of less than 180 cm total length, slow growth and sexual maturity with size at maturity (50%) both sexes: 95-110 cm and low fecundity (Compagno, 1984; Kyalo and Stephen, 2013; Stevens *et al.*, 2000). A report from (Mourier *et al.*, 2013) confirms that *C. melanopterus* males reached sexual maturity at 111 cm. The maximum age from wild-caught individuals is estimated to be 15 years; however, captive animals have lived for > 25 years (Chin *et al.*, 2013b). This study evaluates the condition of *C. melanopterus* stock in the southern Makassar Strait Waters using several population dynamics parameters, including population growth rate, fishing mortality rate, probability of capture, exploitation rate, and relative yield per recruitment.

Materials and Methods

Description of the study sites

The research location is focused on 2 fish landing site, which are the main landing sites for *C. melanopterus*. Location of study and area of sampling i.e at the site of fish landing Paotere, Makassar City, and the site of fish landing Beba, Regency Takalar,

Province South Sulawesi, Indonesia. Fish sampling locations were as shown as presented in Figure 1.

Data collection and data analysis

This study was conducted from July to September 2020 in the southern Makassar Strait Waters. *C. melanopterus* fish samples were obtained from the number of fishermen's catch that landed at the two fish landing sites using *in situ* measurements. Data that measured was the total length (TL: cm). Length frequency data is collected twice a week from the bycatch of bottom longline and bottom gill nets of the traditional fishery. Length data were grouped into 5 cm length groups, and the midpoint is used for the study.

In this study, population parameters as length asymptotic (L_{∞}) and growth coefficient (K), natural mortality (M) and fishing (F), rate of exploitation (E), and relative yield per recruitment (Y'/R) of *C. melanopterus* were estimated based on monthly length-frequency data using FiSAT II software (Gayanilo *et al.*, 2005).

The von Bertalanffy's growth curve (Sparre *et al.*, 1989) as follows: $L_t = L_{\infty}[1 - e^{-K(t-t_0)}]$, where: L_t = length of fish at time t (cm), L_{∞} = asymptotic length (cm), K = the growth coefficient (year^{-1}).

The value of asymptotic length and growth coefficient were estimated using monthly data percentage in the ELEFAN-I by the Response Surface method of FiSAT II. Estimates of growth parameters




Fig. 1. Research Location

were obtained from the L_{∞} and K pairs, which gave the highest R_n value ($R_n = 10ESP / ASP / 10$). t_0 - the hypothetical fish age at zero-length (yr). t_0 was determined using the empirical formula of Pauly (1980), as follows:

$$\text{Log}(-t_0) = -0.3922 - 0.2752 (\text{Log } L_{\infty}) - 1.038 (\text{Log } K)$$

The total mortality rate (Z) was calculated using the length converted catch curve method by FISAT II (Gayanilo *et al.*, 2005; Pauly, 1983). Length converted catch curves are created by plotting $\ln(N_i / \Delta t)$ against relative age t_i . A first estimate of Z is obtained when the following function is adjusted to the points of the right descending arm of the catch curve: $\ln(N_i / \Delta t) = a + b t_i$ where, N_i is the number of fish in length class i , Δt_i is the time needed for the fish to grow through length class i , t_i is the age (or the relative age) corresponding to the mid length of class i , and where b , with the sign changed, is an estimate of Z .

Natural mortality rate (M) was calculated using the empirical formula of Pauly (1980) with a mean annual surface temperature (T) of 28°C as follows: $M = \text{Exp}(-0.152 - 0.279 (\ln L_{\infty}) + 0.6543 (\ln K) + 0.4634 (\ln T^{\circ}\text{C}))$

Where: M = natural mortality rate (yr^{-1}), L_{∞} = Infinity Length (cm), K = coefficient of growth rate (year^{-1}), and T = average temperature of the water ($^{\circ}\text{C}$).

Mortality rate of fishing (F) estimated using the equation: $F = Z - M$ (Pauly, 1980; Pauly, 1983). As for the rate of exploitation (E) $E = F / Z$. The size of length at first capture (L_c) is analyzed based on the estimated logistic curve or the estimated selection ogive function ($S_{L_{est}}$) in the following equation (Sparre *et al.*, 1989):

$$S_{L_{est}} = \frac{1}{1 + \exp(S1 - S2 * L)} \quad L_c = S1 / S2$$

Where: $S_{L_{est}}$ - the estimated logistic curve or the estimated selection ogive function

$S1$ - the intercept, $S2$ - the slope in logistic curve.

L_c was computed using the FISAT-II tool (Gayanilo *et al.*, 2005).

Relative yield per recruit (Y'/R) model of Beverton and Holt (Sparre *et al.*, 1989) as follows:

$$(Y'/R) = E \cdot U^{M/K} \left(1 - \frac{3U}{1+m} + \frac{3U^2}{1+2m} - \frac{U^3}{1+3m} \right)$$




Fig. 2. Growth curve of *Carcharhinus melanopterus* in southern Makassar Strait Waters

Where:

$$U = 1 - \frac{Lc}{L_{\infty}}, \quad m = \frac{E}{M/K} = K / Z, \quad \text{and} \quad E = F/Z$$

Where: E = rate exploitation; U = the fraction of the growth to be completed after entry into the exploited phase;

K = coefficient growth rate (yr^{-1}); L_{∞} = asymptot fish Length (cm); L_c = size of the smallest class of fish caught (cm); and M = natural mortality rate (yr^{-1}).

Results

Growth - The value of the von Bertalanffy growth equation for *Carcharhinus melanopterus* was estimated that L_{∞} was 161.5 cm, K was 0.25 yr^{-1} , and R_n was 0.425 at $SS=2$ and $SL=77.5$. t_0 was -0.4217 yr . Based on K , L_{∞} , and t_0 values obtained above, the exponential growth equation of von Bertalanffy of *C. melanopterus* in the southern Makassar Strait Waters could be written as follows: $L_t = 161.5 [1 - \exp(-0.25(t + 0.4217))]$




Fig. 3. Graphic relative yield per the recruitment of *Carcharhinus melanopterus* in southern Makassar Strait Waters

$0.25^{(1+0.4217)}$]. In using von Bertalanffy growth equation, the age of every length of fish can be predicted easily (Figure 2), as follows:

Mortality and Exploitation Rate - Using the $L_{\infty} = 161.5$ cm and $K = 0.25$ yr⁻¹ obtained through ELEFAN I and $M = 0.45$ yr⁻¹ from Pauly's (1980) empirical relationship, the estimated value of total mortality (Z) obtained from the catch curve is $Z = 1.05$ yr⁻¹, which result in fishing mortality (F) = $1.05 - 0.45 = 0.60$ yr⁻¹. Hence the exploitation rate ($E = F/Z$) was 0.57. A value of $L_c = 99.64$ cm was obtained. **Relative Yield per Recruitment.** The yield per recruit was determined as a function of the exploitation rate, assuming L_c/L_{∞} and M/K are 0.6592 and 1.76, respectively. The plot of relative yield per recruit (Y'/R) against E is shown in (Figure 3), where the maximum Y'/R (0.041) was obtained at $E_{max} = 0.85$, and as the exploitation rate increases beyond this value, relative yield per recruit decreases. In this research the value of actual Y'/R obtained was 0.036 ($E = 0.57$) less than optimal value of Y'/R was 0.04 ($E = 0.85$) (Figure 3).

Discussion

The value of von Bertalanffy growth equation parameters (Figure 2) shows no difference compared with the L_{∞} and K values estimated by other researchers for the same species from different waters. A report from (Chin *et al.*, 2013b) said that the value of asymptotic length (L_{∞}) = 1585 mm LST; K (from logistic model) = 0.251 yr⁻¹ for *C. melanopterus* from north-eastern Australia. There is regional variation in growth, and the maximum size is generally less than 160 cm though individuals have been recorded up to 180 cm (Mourier *et al.*, 2013; Papastamatiou *et al.*, 2009). Different methods, different data, and factors of habitats' eco-biological conditions from time to time may be used to produce different results.

In this research, the rate of exploitation was 0.57. A statement from Mallawa (2012) explained that the exploitation rate is categorized high if $E > 0.5$. If the actual exploitation rate value is related to the maximum exploitation rate value, then the actual value for *C. melanopterus* in the Makassar Strait's southern waters is still below its optimum value ($E_{max} = 0.85$), so that its status is still underexploited. It is mainly carried out as by catch at bottom gill nets and bottom longline fisheries.

Based on the von Bertalanffy growth equation

obtained, the age of the *C. melanopterus* at the first time it was caught (L_c) was 3.417 year. The estimated L_c value above indicates that the average size of the *C. melanopterus* caught by the fishing gear is a group of young fish. To maintain the availability of *C. melanopterus* species stock, efforts are made to increase the L_c value so that the *C. melanopterus* minimum caught by the fishing gear has spawned with a total length > 111 cm. Report from (Chin *et al.*, 2013b; Mourier *et al.*, 2013) said that *C. melanopterus* for males matured at 4.2 yr (1050 mm L_{st}) and females at 8.5 yr (1335 mm L_{st}).

The current exploitation rate of *C. melanopterus* based on mortalities rate, $E = 0.57$, is lower than E_{max} and E_{MSY} based on relative yield per recruit analysis. Based on Figure 3, if we intend to keep the *C. melanopterus* population in equilibrium condition (Y'/R optimum), the rate of exploitation of the *C. melanopterus* population must be increased. A Beverton-Holt relative yield per recruit (Y'/R) model explored that the exploitation rate is underfishing less than 27.06% for E_{max} .

Conclusion

- The population of *Carcharhinus melanopterus* in the southern Makassar strait dominated by small fishes and grow was slowly.
- The values of dynamic population parameters were not to difference with the population of *C. melanopterus* in the specific fishing area, and the natural death less than the death because of the fishing.
- To maintain the population in equilibrium condition, the number (mass) of fishes recruited must be increasing by increasing the size of the fish caught.
- The rate of exploitation of *C. melanopterus* is currently low more than the optimal exploitation rate.

Acknowledgments

The authors wish to thank the leaders of Hasanuddin University and the research community service institute (LP2M). This work was supported in part by a grant from Basic Research of Hasanuddin University with contract number 1585/UN4.22/PT.01.03/2020 on 27 Mei 2020.

References

- Chin, A., Heupel, M. R., Simpfendorfer, C. A. and Tobin, A. J. 2013a. Ontogenetic movements of juvenile blacktip reef sharks: evidence of dispersal and connectivity between coastal habitats and coral reefs. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 23 (3) : 468–474.
- Chin, A., Simpfendorfer, C., Tobin, A. and Heupel, M. 2013b. Validated age, growth and reproductive biology of *Carcharhinus melanopterus*, a widely distributed and exploited reef shark. *Marine and Freshwater Research*. 64(10) : 965–975.
- Compagno, L. J. V. 1984. FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2. Carcharhiniformes. *FAO Fisheries Synopsis*. 125(4): 251–655.
- Fahmi and Dharmadi 2013 [Status of Shark Fisheries and Aspects of Management]. *Oseana*, Volume XXX, Number 1, 2005: 1-8. (in Indonesia)
- Fowler, S. L. and Cavanagh, R. D. 2005. *Sharks, rays and chimaeras: the status of the Chondrichthyan fishes: status survey* (Vol. 63). IUCN.
- Gayanilo, F. C., Sparre, P. and Pauly, D. 2003. FAO-ICLARM stock assessment tool (FiSAT II) user's guide. *FAO Computerized Information Series (Fisheries)*. 8 : 266.
- Kyalo, K. B. and Stephen, N. 2013. *Shark bycatch-small scale tuna fishery interactions along the Kenyan coast*. IOTC–2013–WPEB09–13.
- Mallawa, A. 2012. *Aspects of Fishing and Catch Per Unit Effort of Skipjack in Luwu Waters, Bone Bay*. 325–330.
- Mourier, J., Mills, S. C. and Planes, S. 2013. Population structure, spatial distribution and lifehistory traits of blacktip reef sharks *Carcharhinus melanopterus*. *Journal of Fish Biology*. 82 (3) : 979–993.
- Papastamatiou, Y. P., Caselle, J. E., Friedlander, A. M. and Lowe, C. G. 2009. Distribution, size frequency, and sex ratios of blacktip reef sharks *Carcharhinus melanopterus* at Palmyra Atoll: a predator dominated ecosystem. *Journal of Fish Biology*. 75(3) : 647–654.
- Pauly, D. 1983. Some Simple Methods for the Assessment of Tropical Fish Stock. *FAO Fisheries Technical Paper*, 254, 52.
- Pauly, D. 1980. On the inter-relationships between natural mortality, growth performance and mean environmental temperature in 175 fish stock. *Journal du Conseil*. 39(3) : 175-192.
- Sparre, P.E., Ursine, and S.C. Venema. 1989. Introduction to Tropical Fish Stock Assessment. Part I Manual. *FAO Fisheries Technical 306/1*. Rome. 337p
- Stevens, J. D., Bonfil, R., Dulvy, N. K. and Walker, P. A. 2000. The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal of Marine Science*. 57(3) : 476–494.
-

ORIGINALITY REPORT

22%

SIMILARITY INDEX

16%

INTERNET SOURCES

21%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

- 1 M. Magdy, HALFAWY, El, AMIN, Amal, M. and RAMADAN, Amal, M.. "Growth and reproduction of female brushtooth lizardfish *Saurida undosquamis* (Richardson) from the Gulf of Suez, Egypt", Tubitak, 2007.
Publication 2%
 - 2 Submitted to The University of the South Pacific
Student Paper 2%
 - 3 Submitted to Universitas Hasanuddin
Student Paper 2%
 - 4 A Damora, T Firdiyanti, A Rahmah, R M Aprilla, M A Chaliluddin. "Population dynamics of Indian scad (*Decapterus russelli*) in the northern and western waters of Aceh", IOP Conference Series: Earth and Environmental Science, 2021
Publication 1%
 - 5 docobook.com
Internet Source 1%
-

6

Internet Source

1 %

7

Submitted to University of Malaya

Student Paper

1 %

8

worldwidescience.org

Internet Source

1 %

9

G Bintoro, T D Lelono, D P Ningtyas. " Biological aspect of mackerel scad (Cuvier, 1833) in Prigi waters Trenggalek Regency East Java Indonesia ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

1 %

10

Edwine Yongo, Simon Agembe, Nicholas Outa, Monica Owili. " Growth, mortality and recruitment of Nile perch () in Lake Victoria, Kenya ", Lakes & Reservoirs: Science, Policy and Management for Sustainable Use, 2018

Publication

1 %

11

J Tresnati, A Yanti, N Rukminasari, Irmawati, Suwarni, I Yasir, P Y Rahmani, R Aprianto, A Tuwo. " Sex ratio, maturity stage and fist maturity of yellowfin parrotfish Schultz, 1958 in Wallace line at Spermonde Archipelago, South Sulawesi ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

1 %

12 P.O Bannerman. "Stock assessment of the big-eye grunt (*Brachydeuterus auritus*, Val.) fishery in Ghanaian coastal waters", Fisheries Research, 20021230

Publication

1 %

13 Aswathy Vijaya Krishna, Parvathi Ammini. " Population characteristics of in Vembanad Lake, India ", Lake and Reservoir Management, 2017

Publication

<1 %

14 Rueda, M.. "Population dynamics and fishery of the fresh-water clam *Polymesoda solida* (Corbiculidae) in Cienaga Poza Verde, Salamanca Island, Colombian Caribbean", Fisheries Research, 19981201

Publication

<1 %

15 jifro.ir
Internet Source

<1 %

16 mail.scialert.net
Internet Source

<1 %

17 Cliff A. Robb. "Paying for College: Advice from Current Students to Incoming Freshmen", Family and Consumer Sciences Research Journal, 2011

Publication

<1 %

18 Fatma A. Abdel Razek, Evelyn Ragheb, Rabab S. El-Deeb, Hamdy Omar Ahmed. "Growth

<1 %

pattern and stock assessment of Jinga shrimp *Metapenaeus affinis* (H. Milne Edwards, 1837) (Decapoda, Penaeidae) from the southeastern Mediterranean of the Egyptian coasts", *The Egyptian Journal of Aquatic Research*, 2022

Publication

19

M. Njiru, P. Nzungi, A. Getabu, E. Wakwabi, A. Othina, T. Jembe, S. Wekesa. "Are fisheries management, measures in Lake Victoria successful? The case of Nile perch and Nile tilapia fishery", *African Journal of Ecology*, 2007

Publication

20

adoc.pub

Internet Source

<1 %

21

en.wikipedia.org

Internet Source

<1 %

22

eprints.cmfri.org.in

Internet Source

<1 %

23

www.sid.ir

Internet Source

<1 %

24

Faisal Amir, Achmar Mallawa, Joeaharnani Tresnati. "Size Structure and Sex Ratio of Black Tip Reef Shark (*Carcharhinus melanopterus*) Landed in the TPI Paotere of Makassar City and TPI Beba of Takalar

<1 %

- 25 J. Dulčić. "Age, growth and mortality of brown comber, *Serranus hepatus* (Linnaeus, 1758) (Pisces: Serranidae), in the eastern Adriatic (Croatian coast)", *Journal of Applied Ichthyology*, 4/2007 $<1\%$
- Publication
-

- 26 R Sala, R Bawole, F Runtuboi, Mudjirahayu, I A Wopi, J Budisetiawan, Irwanto. " Population dynamics of the yellowstripe scad (Cuvier, 1833) and Indian mackerel (Cuvier, 1816) in the Wondama Bay Water, Indonesia ", *IOP Conference Series: Earth and Environmental Science*, 2018 $<1\%$
- Publication
-

- 27 Ria Faizah, Lilis Sadiyah, Moh. Fauzi. "POPULATION PARAMETER AND REPRODUCTIVE BIOLOGY OF INDIAN MACKEREL *Rastrelliger kanagurta* (Cuvier, 1817) CAUGHT BY LIFT NET IN KWANDANG WATERS, NORTH GORONTALO", *Indonesian Fisheries Research Journal*, 2018 $<1\%$
- Publication
-

- 28 citeseerx.ist.psu.edu $<1\%$
- Internet Source
-

29

Internet Source

<1 %

30

www.frontiersin.org

Internet Source

<1 %

31

zsp.com.pk

Internet Source

<1 %

32

Submitted to Universitas Brawijaya

Student Paper

<1 %

33

Andrew Chin, Colin Simpfendorfer, Andrew Tobin, Michelle Heupel. "Validated age, growth and reproductive biology of *Carcharhinus melanopterus*, a widely distributed and exploited reef shark", *Marine and Freshwater Research*, 2013

Publication

<1 %

34

Baset Abdul, Liu Qun, Liao Baochao, Waris Abdul, Yanan Han, Qingqing Zhang. "Population dynamics of Rainbow Sardines, *Dussumieria acuta* (Valenciennes, 1847) from Pakistani waters", *International Journal of Aquaculture and Fishery Sciences*, 2020

Publication

<1 %

35

G.A. Charles, K. Sivashanth. "Population Dynamics of Squid *Sepioteuthis lessoniana* (Lesson, 1830) from the Northern Coast of Sri Lanka", *Journal of Fisheries and Aquatic Science*, 2010

<1 %

36

King, Michael. "Stock Structure and Abundance", Fisheries Biology Assessment and Management King/Fisheries Biology Assessment and Management, 2013.

Publication

<1 %

37

Kwarfo-Apegyah, K, PK Ofori-Danson, and FKE Nunoo. "Exploitation Rates and Management Implications for the Fisheries of Bontanga Reservoir in the Northern Region of Ghana", West African Journal of Applied Ecology, 2009.

Publication

<1 %

38

Marcus Rodrigues da Costa, Raquel Rennó Mascarenhas Martins, Acácio Ribeiro Gomes Tomás, Rafael de Almeida Tubino et al. "Biological aspects of *Mugil liza* Valenciennes, 1836 in a tropical estuarine bay in the southwestern Atlantic", Regional Studies in Marine Science, 2021

Publication

<1 %

39

N Abiaobo, M Udo. "The Population Dynamics of the Mudskipper, *Periophthalmus barbarus* (LINNEAUS 1766) (TELEOSTEI, GOBIIDAE) and the Implication for Conservation and Management in the Mangrove Swamp of Iko River Estuary, Southeastern Nigeria", Journal of Applied Life Sciences International, 2017

Publication

<1 %

40

Niamaimandi, N.. "Population dynamic of green tiger prawn, *Penaeus semisulcatus* (De Haan) in Bushehr coastal waters, Persian Gulf", Fisheries Research, 200709

Publication

<1 %

41

researchonline.jcu.edu.au

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On