

BIODIVERSITY OF MARINE TUNICATES IN SAMALONA WATERS, SANGKARANG ARCHIPELAGO, INDONESIA

by

FILE	ET_AL-SPERMONDE-2018-26-31_BIODIVERSITY_OF_MARINE_TUNICATES.PDF (741.89K)		
TIME SUBMITTED	29-MAY-2019 02:16PM (UTC+0700)	WORD COUNT	4271
SUBMISSION ID	1137317635	CHARACTER COUNT	22482

BIODIVERSITY OF MARINE TUNICATES IN SAMALONA WATERS, SANGKARANG ARCHIPELAGO, INDONESIA

Magdalena Litaay^{1*}, Slamet Santosa¹, Eva Johannes¹, Rosana Agus¹, Willem Moka¹,
Jennyta Dhewi Darmansyah Tanjung¹

Submitted: 19 February 2018 Accepted: 26 February 2018

ABSTRACT

The study aims to know the biodiversity and community structure of marine tunicate in Samalona waters. The present study is part of biodiversity assessment for marine resources of Sangkarang Archipelago SW Makassar Indonesia. Field campaign was conducted from October to November 2016. Sample collection was done at 3 and 7 m depth by using Line Intersection Transect (LIT) method combined with a quadrat (plot). Two 50 m transects were placed parallel to a shore line at three stations (sta.) at Samalona waters. A quadrat (plot) (2.5 m x 2.5 m) was placed side by side of the line transect and all tunicates in the transect was recorded, identified, counted and photographed. Samples were collected by using SCUBA and under water camera. Environmental parameters including water temperature, salinity, dissolved oxygen, clarity, current and wind speed, were measured *in situ*. Data were analysed using ecological indices including species composition and density, Shannon Wiener species diversity, Evenness, and Morisita Indices. The result indicates that there are 18 species of tunicates present at 3 m as well as 7 m depth of Samalona waters. Result of the ecological analysis shows that species diversity can be categorized as moderate and there were no dominant species. Environmental parameters indicates that water quality at Samalona waters was in good condition to support tunicates.

Key words: ascidian, chordata, coral reefs, Spermonde

INTRODUCTION

Marine invertebrates as major group living in coral reefs of the Indo-Pacific region are rich for secondary metabolite and are targeted for studying lead compound as marine drug discovery (Sabdon and Radjasa 2008). Radjasa *et al.* (2011) stated that coral reef ecosystem is a source for bioactive compound origin from its associated biotas such as sponge, ascidians, mollusks, bryozoans and cnidarian.

Biodiversity of marine biotas has pushed the discovery of marine natural products that can be developed as a therapeutic candidate. Marine tunicates were potential for inoculum source for endo-symbiotic that can produce anti-bacterial and anti-fungi (Karthikeyan *et al.* 2009; Litaay *et al.* 2015; Christine *et al.* 2015; Nurfadillah *et al.* 2015; Sardiani *et al.* 2015; Tahir *et al.* 2016). Tunicates are also potential antiviral (Murti and Agrawal, 2010), anti-cancer (Shaala and Youssef, 2015), inhibitor and induces apoptosis of breast (MCF-7; MDA-MB) cancer cells and also used for phase II cancer treatment (Zelek *et al.* 2006; Michaelson *et al.* 2012; Atmaca *et al.* 2013), as inhibitor of breast cancer cells by JNK dependent apoptosis (Gonzalez-Santiago *et al.* 2006), breast and prostate cancer (Kalimuthu *et al.* 2014). One of the bioactive compounds produced by tunicate is used to cure refractory soft-tissue sarcomas (Sinko *et al.* 2012). Sangkarang Archipelago which was previously known as Spermonde is located South West off Makassar,

consisting more than a hundred islands. Various marine biodiversity in Sangkarang area have been studied (Moll 1983; Verheij 1993; Massin 1999; Renema and Simon 2001; de Voogd *et al.* 2006; Pogoreutz *et al.* 2012; Priosambodo *et al.* 2014). However, diversity of marine tunicates in the Sangkarang region is less studied (Fikruddin, 2013; Mawaleda, 2014). In order to explore bio-prospecting of marine tunicate, a basic research on species diversity is needed. Information on biodiversity and distribution of tunicate in this region will provide useful baseline data to support sustainable use of marine resources.

MATERIALS AND METHODS

Sample Collection

Sampling of tunicates was done at three different sites (sta.) of Samalona waters, Sangkarang Archipelago South Sulawesi Indonesia (Figure 1). SCUBA was used in collecting sample and LIT




Figure 1. Location at Samalona waters, red spot indicating the sampling sites. sta 1 (5°7.390'S; 119°20.576'E), sta 2 (5°7.513'S; 119°20.330'E) and sta 3 (5°7.699'S; 119°20.506'E).

¹Department of Biology, Hasanuddin University
Jl Perintis Kemerdekaan Km 10. Makassar 90245,
Indonesia.

*Magdalena Litaay
Email: mlitaay@fmpipa.unhas.ac.id

at 3 m and 7 m depth at three stations (Figure 1). In each station, a 50 m line transect was placed parallel to shore line and a quadrat (2.5 m x 2.5 m) was applied side by side of transects. All tunicates in the quadrat were recorded, identified, counted and photographed. Sub-sample was taken and brought to laboratory for identification purposes. Environmental parameter was measured *in situ*. Underwater camera Canon GIS was used to obtain images of species and habitat (English *et al.* 1997; Brower *et al.* 1998). Water parameter including type of substrate, pH, temperature, current, salinity, clarity and dissolved oxygen were measured during sampling. Identification of tunicates was based on main morphological characters according to Kott (2005), Page and Kelly (2013), WoRMS (2017).

Data Analysis

Data analysis were based on ecological indices including density, Shanon Wiener diversity index, Dominance Index, Dispersion Index of Morisita (Odum, 1993) as follows

$$D_i = n_i / A$$

Where;
 D_i = Density species i
 n_i = Total no of individuals of species i in all quadrats
 A = Total area of quadrats (plot)

Species composition = (No of individual sp- i / Total no of individual) x 100 %

Dominance Index (Odum, 1993)

$$C = \sum (n_i / N)^2$$

Where;
 n_i = No of individual species i
 N = Total no of individuals of all species
 C = Dominance Index
 Criteria :
 $0 < C \leq 0.5$ = Low
 $0.5 < C \leq 0.75$ = Moderate
 $0.75 < C \leq 1.00$ = High

Shanon-Wiener Diversity Index

$$H = -\sum (n_i / N) \ln (n_i / N)$$

Where;
 n_i = Number of species - i
 N = Total number of species
 H' = Diversity Index
 Criteria :
 $0 < H' \leq 2.0$ = Low
 $2 < H' \leq 3.0$ = Modertae
 $3.0 < H' \leq 4.0$ = High

Morisita Index

$$I_d = n \frac{\sum X^2 - N}{N(N-1)}$$

Where;
 n = total number of plot/transect = $\sum F(x)$
 N = Total no of individual inside plot/transect = $\sum [F(X)] (X)$

$\sum X^2$ = Sum of the squares of the number of individual inside plot = $\sum [F(X)] (X^2)$

Criteria :
 $I_d < 1.0$ = Randomly distributed
 $I_d = 1.0$ = Uniformly distributed
 $I_d > 1.0$ = Clumped distributed

Correspondance Analysis (CA) was used to observe the relationship between tunicates with water parameters (SPSS ver 23).

RESULTS AND DISCUSSION

Species Composition

The result shows that there are 18 species of tunicate present at reef flat (3 m) and reef slope (7 m) of Samalona waters (Table 1). The highest percentage of species diversity was found at 3 m depth at sta. 3 (83.3%) and lower in Sta 1 and 2 (77.2 %). At 7 m depth, we found similar number of tunicate species at sta. 2 and 3 (72,2%), yet higher compared to sta. 1. We recorded 7 families and number of species of tunicates (in brackets) as follows Clavelinidae (3), Didemniidae (3), Diazonidae (1), Styelidae (4), Ascidiidae (1), Perophoridae (3), Pyuridae (3) (Table 1).

Species Richness

The result of species richness is provided in Table 2. Table 2 shows that *D. molle* were more abundant at 3 m depth reef flat in all stations. This species is small in size, living solitary or forming a colony. Hence, *D. molle* can be found in a big colony at different habitats and depths. Previous finding indicated that *Didemnum* can grow at different habitats (Bullard and Whitlatch 2008; Carman *et al.*, 2010).

Diversity, Morisita and Dominance Indices

Table 3 shows value of species diversity and Morisita indices at different sites and depths in Samalona waters. Based on criteria used in this study, diversity of tunicate is classified as low at 3 m depth at sta. 1 and sta. 2, while moderate at other depths and sites. Values of Morisita index indicates that tunicates are randomly distributed at all depths and sites (Table 3).

Table 4 shows the value of dominance index ranges from 0 to 0.68. High values were recorded for *Didemnum molle* at 3 m depth at sta. 2 as well as sta. 1. This indicates a moderate dominance of particular species at those sites. As also described in Table 4, value of dominance index was less than 0.5 or close to zero.

Table 1. The occurrence of tunicates at different sites and depth of Samalona water

Tunicates Family & species	Sta 1		Sta 2		Sta 3	
	3 m	7 m	3 m	7 m	3 m	7 m
<i>Clavelinidae</i>						
<i>Clavelina robusta</i> Kott 1990 (<i>white spot</i>)	+	+	+	+	+	+
<i>Clavelina lepadiformis</i> Muller, 1776	+	+	+	+	+	+
<i>Oxycornia fascicularis</i> Drasche, 1882	-	+	+	+	+	+
<i>Didemnidae</i>						
<i>Didemnum molle</i> Hermann, 1986	+	+	+	+	+	+
<i>Lissoclinum patella</i> Gottschaldt 1898	-	+	-	-	+	-
<i>Trididemnum della</i> Ritter & Forsyth, 1917	+	+	+	+	+	+
<i>Diazoniidae</i>						
<i>Rhopalea crassa</i> Hermann, 1880	+	+	+	+	+	+
<i>Styelidae</i>						
<i>Polycarpa papillata</i> Sluiter, 1886	+	+	+	+	+	+
<i>Polycarpa aurata</i> Quoy & Gaimard, 1834	+	+	+	+	+	+
<i>Polycarpa nigricans</i> Heller, 1878	-	+	+	+	+	+
<i>Polycarpa spongiabilis</i> Traustedt 1883	+	-	-	-	+	-
<i>Asciidiidae</i>						
<i>Ascidia sydneisis</i> Monniot F. 1898	-	+	+	+	+	+
<i>Perophoridae</i>						
<i>Pherophora sp (soft blue)</i>	+	-	-	-	-	-
<i>Pherophora sp (orange)</i>	+	-	-	+	-	+
<i>Perophora amectens</i> Ritter, 1893	+	-	+	-	+	-
<i>Pyuridae</i>						
<i>Pyura molina</i> Blainville, 1824	-	+	-	-	+	+
<i>Halocynthia verrill</i> Dumosa Simpson, 1885	+	-	+	+	+	+
<i>Microcosmus juinii</i> Drasche 1884	+	-	+	-	-	-
Total percentage (%)	77.7	66.6	77.7	72.2	83.3	72.2

Table 2. Species richness at different sites and depths of Samalona waters (ind/500m²)

No.	Tunicates species	Sta 1		Sta 2		Sta 3	
		3 m	7 m	3 m	7 m	3 m	7 m
1	<i>Clavelina robusta</i> Kott 1990 (<i>white spot</i>)	0.126	0.054	0.004	0.058	0.042	0.016
2	<i>Clavelina lepadiformis</i> Muller, 1776	0.04	0.014	0.008	0.004	0.372	0.04
3	<i>Oxycornia fascicularis</i> Drasche, 1882	0	0.03	0.006	0.054	0.052	0.03
4	<i>Didemnum molle</i> Hermann, 1986	2.286	0.79	2.16	0.564	0.872	0.416
5	<i>Lissoclinum patella</i> Gottschaldt 1898	0	0.028	0.012	0.036	0.026	0.004
6	<i>Trididemnum della</i> Ritter & Forsyth, 1917	0.01	0.002	0.048	0.012	0.012	0.008
7	<i>Rhopalea crassa</i> Hermann, 1880	0.012	0.01	0.002	0.032	0.01	0.014
8	<i>Polycarpa papillata</i> Sluiter, 1886	0.352	0.052	0.068	0.09	0.12	0.056
9	<i>Polycarpa aurata</i> Quoy & Gaimard, 1834	0.178	0.25	0.092	0.212	0.344	0.096
10	<i>Polycarpa nigricans</i> Heller, 1878	0.134	0.128	0.186	0.078	0.206	0
11	<i>Polycarpa spongiabilis</i> Traustedt 1883	0.008	0	0	0	0.01	0
12	<i>Ascidia sydneisis</i> Monniot F. 1898	0	0.008	0.002	0	0.004	0.002
13	<i>Pherophora sp (soft blue)</i>	0.006	0	0	0	0	0
14	<i>Pherophora sp (orange)</i>	0.004	0	0.002	0	0.002	0
15	<i>Perophora amectens</i> Ritter, 1893	0.004	0	0	0.002	0	0.002
16	<i>Pyura molina</i> Blainville, 1824	0	0.132	0	0	0.094	0.024
17	<i>Halocynthia verrill</i> Dumosa Simpson, 1885	0.018	0	0.016	0.074	0.002	0.002
18	<i>Microcosmus juinii</i> Drasche 1884	0.002	0	0.002	0	0	0

Table 3. Value of Diversity and Morisita Indices

Sites	Depth (m)	Diversity Index	Morisita Index
Sta 1	3	0.99	0.35
	7	1.55	0.53
Sta 2	3	0.73	0.25
	7	1.68	0.58
Sta 3	3	1.67	0.57
	7	1.35	0.46

Table 4. The Result of Dominance Analysis

No	Tunicate Species	Sta 1		Sta 2		Sta 3	
		3 m	7 m	3 m	7 m	3 m	7 m
1	<i>Clavelina robusta</i> Kott 1990 (white spot)	0.002	0.001	0	0.002	0	0.001
2	<i>Clavelina lepadiformis</i> Muller, 1776	0	0	0	0	0.029	0.003
3	<i>Oxycomia fascicularis</i> Drasche, 1882	0	0	0	0.002	0.001	0.002
4	<i>Didemnum molle</i> Hermann, 1886	0.517	0.278	0.686	0.215	0.162	0.343
5	<i>Lissoclinum patella</i> Gottschaldt 1898	0	0	0	0.001	0	0
6	<i>Trididemnum della</i> Ritter & Forsyth, 1917	0	0	0	0	0	0
7	<i>Rhopalea crassa</i> Hermann, 1880	0	0	0	0.001	0	0
8	<i>Polycarpa papillata</i> Sluiter, 1886	0.012	0.001	0.001	0.005	0.003	0.006
9	<i>Polycarpa aurata</i> Quoy & Gaimard, 1834	0.003	0.028	0.001	0.030	0.025	0.018
10	<i>Polycarpa nigricans</i> Heller, 1878	0.002	0.007	0.005	0.004	0.009	0
11	<i>Polycarpa spongiabilis</i> Traustedt 1883	0	0	0	0	0	0
12	<i>Ascidia sydneinsis</i> Monniot F. 1898	0	0	0	0	0	0
13	<i>Pherophora</i> sp (soft blue)	0	0	0	0	0	0
14	<i>Pherophora</i> spp (orange)	0	0	0	0	0	0
15	<i>Perophora annectens</i> Ritter, 1893	0	0	0	0	0	0
16	<i>Pyura molina</i> Blainville, 1824	0	0.008	0	0	0.002	0.001
17	<i>Halocynthia verrill</i> Dumosa Simpson, 1885	0	0	0	0.004	0	0
18	<i>Microcosmus junii</i> Drasche 1884	0	0	0	0	0	0

Environmental Parameters

The range of environmental parameters in study sites (Table 5) are as follows: temperature 30-31°C, salinity 31-32.5‰, dissolved oxygen (DO) 6.1-6.2 ppm, wind velocity 5.3-10.5 km/h, current 0.1-1.1 km/h, clarity 8.2-11.2 m, and pH 8.4. In general, water parameters are preferable for marine invertebrates.

Table 5. Environment parameters at Samalona waters

Water Parameters	sta 1	sta 2	sta 3
Temperature °C	31	30	30
Salinity ‰	31	31.3	32.5
DO (ppm)	6.2	6.2	6.1
Wind Velocity (km/h)	10.5	9.8	5.3
Current (km/h)	0.1	1.1	0.7
Clarity (m)	8.2	10	11.2
pH	8.4	8.4	8.4

Relationship Between Tunicates With Environment Parameters

The result of Correspondence Analysis (CA) between species composition and environmental parameters is given in Figure 2.




Figure 2. The relationship between species composition with environmental parameters (Note: 1-18 tunicates species as indicated in Table 1; 19 = temperature, 20 = salinity, 21 = DO, 22 = wind velocity, 23 = current, 24 = clarity, 25 = pH)

As shown in figure 2, distribution of most of the tunicate are related to environmental parameters as they are close to center of the quadrant. While *Clavelina lepadiformis* (2), *Ascidia sydneinsis* (12), and *Halocynthia verrill* (17) are different compare to other ascidians as indicated in Figure 2. These species were found solitary living amongst brain coral (*A. sydneinsis*), *H. verrill* in small size and abundant at 3 m. This may indicate that other factors may influence the distribution of these particular tunicates.

Tunicates from Sangkarang archipelago is less documented. This study was the first record for marine tunicate from Samalona waters of this archipelago. This study indicates that a number of tunicate species for this area is less than recorded from Baranglompo island of Sangkarang. Mawaleda (2014) found 33 species of tunicates at coral reef areas of Baranglompo waters. However, this study shows more tunicates species are found in Samalona waters compared to those recorded for other islands of Sangkarang: 7 species (Lae-Lae), 9 species (Bone Batang) and 10 species (Badi) waters (Fikruddin, 2013). Tunicate class Ascidiacea is the most diverse group, as 700 species have been recorded in the Australian waters, while Thaliacea is having less than 100 species worldwide and Appendicularia is about 60 species shown worldwide (Kott, 2005). Kiuru *et al.* 2014 estimated that marine and coastal environments host about 90% of all organisms living on earth. Here, we found 18 species tunicate at Samalona waters, this number is still low compare to a worldwide known tunicates.

Marine resources particularly in coastal areas and small islands received main pressure from natural catastrophic and anthropogenic activities. Human activities in main land as use of unfriendly fishing method, overfishing, pollution, coastal development and global warming has have big impact on marine

resources. These also can contribute to loss of biodiversity of marine resources. Sangkarang archipelago has more than a hundred islands, biodiversity of tunicates in this area are still questionable. Therefore, more studies on biodiversity of tunicates in this area is needed.

5

ACKNOWLEDGEMENTS

Funding for this project was provided to Magdalena Litaay from Hasanuddin University through the research scheme of Indonesia Maritime Specific Topic II (BMIS II) 2016. Thanks to Nenis Sardiani, Reignildis Regina, Ilham, Muh Nurdin, Wahyulfatwatul, Ayub Wirabuana, Bachtiar Anas, Marjuni from Biological Celebes Diving Club (BCDC Unhas) for their assistance with field work.

REFERENCES

- Atmaca H, Bozkurt E, Uzunoglu S, Uslu R, Karaca B. 2013. A diverse induction of apoptosis by trabectedin in MCF-7 (HER2- /ER+) and MDA-MB-453 (HER2+/ER-) breast cancer cells. *Toxicology Letters* 221:128-36.
- Brower JE, ZR JH, Von Ende CN. 1998. *Field and Laboratory Methods for General Ecology*, Mc Graw Hill Company.
- Bullard SG and Whitlatch RB. 2009. In situ growth of the colonial ascidian *Didemnum vexillum* under different condition. *Aquatic Invasion* 4, 275-278.
- Carman MR, Morris JA, Karney RC, Grunden DW. 2010. An initial assessment of native and invasive tunicates in shellfish aquaculture of the northeast coast. *Journal Applied Ichthyology* 26:8-11.
- Christine G, Litaay M, Gobel RB, Dwyana Z. 2015. Potency of tunicate *Polycarapa aurata* as inoculum source for endosymbiotic bacteria; characterization of isolates. In: Tahir (eds). *Proceeding National Seminar on Marine and Fisheries*. FIKP Unhas. Makassar, May 2015. [Indonesian]
- de Voogd NJ, Cleary DFR, Hoeksema BW, Noor A, van Soest RWM. 2006. Sponge beta diversity in the Sangkarang Archipelago, SW Sulawesi, Indonesia. *Marine Ecological Progress Series* 309:131-142.
- English S, Wilkinson C, Baker V. 1997. *Survey Manual for Tropical Marine Resources*. Australian Institute of Marine Science. Townsville
- Fikruddin MBAbdH. 2013. Distribution and diversity of Tunicate (Ascidacea) at different habitat at waters of Badi, Bonebatang and Laelae Island. BSc Hon Thesis FIKP Hasanuddin University.
- Gonzalez-Santiago L, Suarez Y, Zarich N, Munoz-Alonso M, Cuadrado A, Martinez T, Goya, Iradi A, Saez-Tormo G, Maier J. 2006. Aplidin® induces JNK-dependent apoptosis in human breast cancer cells via alteration of glutathione homeostasis, Rac1 GTPase activation, and MKP-1 phosphatase downregulation. *Cell Death & Differentiation* 13:1968-81
- Karthikeyan MM, Ananthan G, Balasubramanian T. 2009. Antimicrobial Activity of Crude Extracts of Some Ascidiaceans (Urochordata: Ascidiacea), from Palk Strait, (Southeast Coast of India). *World Journal of Fish and Marine Sciences* 1 (4): 262-267.
- Kalimuthu S, Venkatesan J and Kim Se-Kwon. 2014. Marine Derived Bioactive Compounds for Breast and Prostate Cancer Treatment: A Review. *Current Bioactive Compounds* 10: 62-74.
- Kiuru P, Valeria D'Auria M, Muller CD, Tammela P, Vuorela H, Yli-Kauhaluoma J. 2014. Exploring Marine Resources for Bioactive Compounds. *Planta Med* 2014; 80: 1234-1246. DOI [http://dx.doi.org/ 10.1055/s-0034-1383001](http://dx.doi.org/10.1055/s-0034-1383001) Published online September 9, 2014
- Kott P. 2005. *Catalogue of Tunicata in Australian Waters*. Queensland Museum, Brisbane, Australia. ISBN 0 642 56842 1. 301 p.
- Litaay M, Christine G, Gobel RB, Dwyana Z. 2015. Bioactivity of tunicate *Polycarapa aurata* symbion as antimicrobe. In: Ohee (eds.) *Proceeding the 23rd National Seminar of Indonesia Biology Society*. Jayapura, 18 September 2015.
- Massin 1999. Reef Dwelling Holothuroidea (Echinodermata) of The Sangkarang Archipelago (South-West Sulawesi, Indonesia). *Zoologische Verhandelingen* 329. National Natuurhistorisch Museum. Leiden
- Mawaleda R. 2014. Distribution and habitat preference Urochordata class Ascidiacea at coral reefs area of Barrang Lompo island of Makassar. BSc Hon Thesis FIKP Hasanuddin University.
- Michaelson M, Bellmunt J, Hudes G, Goel S, Lee R, Kantoff P, Stein C, Lardelli P, Pardos I, Kahatt C. 2012. Multicenter phase II study of trabectedin in patients with metastatic castrationresistant prostate cancer. *Annals of Oncology* 23: 1234-40
- Misset J. 2006. A phase II study of Yondelis®(trabectedin, ET 743) as a 24-h continuous intravenous infusion in pretreated advanced breast cancer. *Britainia Journal Cancer* 94.

- Moll H. 1983. Zonation and diversity of scleractinian on reefs of South Sulawesi Indonesia. [Thesis]. Netherland: Leiden University.
- Murti Y and Agrawal T. 2010. Marine derived pharmaceuticals development of natural health products from marine biodiversity. International Journal of ChemTech Research 2 (4): 2198-2217
- Nurfadillah A, Litaay M, Gobel RB, Haedar N. 2015. Potency of tunicate *Polycarpa aurata* as inoculum source of sebagai sumber endosimbyotic fungsi that produce antimicrobe. Jurnal Alam and Lingkungan 6 (12): 10-16.
- Page M and Kelly M. 2013. Awesome Ascidiens, a Guide to the Sea Squirts of New Zealand. TC Media Ltd.53 pp
- Priosambodo D, Kneer D, Asmus H, Zamani NP, von Juterzenka K, Litaay M, Soekendarsi E. 2014. Community analysis of burrower shrimp in Bone Batang seagrass bed South Sulawesi. Proceeding of The First International Conference on Science (ICOS-1) 2014 ISBN 978-602-72198-0-9: 209 -219.
- Pogoreutz, C, H. Asmus, H. Anelt, D. Kneer, M. Litaay. 2012. The influence of canopy structure and tidal level on fish assemblages in tropical Southeast Asian seagrass meadows. Eustarine Coastal and Shelf Science. 107 (2): 58-68.
- Odum EP. 1993. General Ecology. Indonesia version: Dasar-Dasar Ekologi (Tjahjono Samingan; penyunting B. Srigandono). Yogyakarta, Gadjah Mada University Press. 667 pp.
- Radjasa OK, Vaske YM, Navarro G, Vervoort HC, Tenney K, Linington RG, Crews P. 2011. Highlights of marine invertebrate-derived biosynthetic products: Their biomedical potential and possible production by microbial associants Bioorganic & Medicinal Chemistry 19: 6658-6674
- Renema WT and Simon R. 2001. Larger foraminifera distribution on a mesotrophic carbonate shelf in SW Sulawesi (Indonesia). Palaeogeography, Palaeo climatology, Palaeoecology 175:125-146
- Sabdono A and Radjasa OK. 2008. Microbial symbionts in marine sponges: Marine natural product factory. Journal Coastal Development 11: 57-61.
- Sardiani N, Litaay M, Gobel RB, Dwyana Z. 2015. Potency of tunicate *Rhopalaea sp* as source of bacterial inoculum that produce antibacterial; isolates characterization. J. Alam dan Lingkungan 6: 11.
- Shaala LA and Diaa Youssef TA. 2015. Identification and Bioactivity of Compounds from the Fungus *Penicillium sp.* CYE-87 Isolated from a Marine Tunicate. Marine Drugs 2015, 13, 1698-1709; doi:10.3390/md13041698.
- Sinko J, Rajchard J, Balounova Z, Fikotova L. 2012. Biologically active substances from water invertebrates: a review. Veterinarni Medicina, 57(4): 177-184
- Tahir E, Litaay M, Gobel RB, Haedar N, Prisambodo D, Syahribulan. 2016. Potency of tunicate *Rhopalaea crassa* as inoculum source of endosymbiont fungi that produce antimicrobe. Marine Science Journal Spermonde 2(2): 33-37 [Indonesian]
- WoRMS. 2017. World Register of Marine Species. <http://www.marinespecies.org/aphia.php?p=taxdetails&id=146420>.
- Verheij E. 1993. Marine plants on the reefs of the Sangkarang Archipelago, SW Sulawesi, Indonesia: aspect of taxonomy, floristics and ecology. [Dissertation] Leiden: Rijksherbarium / Hortus Botanicus

FORMAT MANUSKRIP
Jurnal Ilmu Kelautan SPERMONDE

I. Petunjuk umum

- Manuskrip diformat pada kertas A4 (21 x 30cm), menggunakan Bahasa Indonesia baku sesuai dengan kaidah Bahasa Indonesia yang baik dan benar
- Semua halaman dilengkapi nomor halaman dan tersusun teratur
- Menggunakan Times New Roman font 12, tanpa menggunakan huruf italic, bold atau tipe penulisan lain yang tidak standar. Jangan meratakan margin kanan. Setiap paragraf baru diberi indentasi
- Gunakan spasi 2 untuk keseluruhan manuskrip termasuk halaman judul, abstrak, daftar pustaka, keterangan tabel dan gambar.
- Margin kiri, kanan, atas dan bawah adalah 2,5 cm (1 inchi)
- Rasio seperti contoh pada unit diharapkan menggunakan garis miring (contoh mg/L)
- Nama ilmiah seharusnya dilengkapi dengan nama umum pada judul dan saat untuk pertama kali dituliskan pada abstrak dan pada teks. Author dari nama ilmiah tidak menyertai genus atau nama spesies kecuali bila dibutuhkan penjelasan lebih lanjut.
- Tuliskan dengan lengkap angka satu hingga sepuluh, kecuali bila diikuti oleh unit pengukuran (empat ikan, 4 ekor ikan, 4kg, 14 ikan). Hindari memulai kalimat dengan angka. Gunakan '1.000' untuk '1000'; 0,13 dan % (bukan 'persen').
- Untuk menuliskan waktu (jam), gunakan sistem 24 jam (13.30 bukan 01.30; 02.30 bukan '2.30'). Data kalender dituliskan dalam susunan hari, bulan, tahun (14 Januari 2014).
- Semua referensi yang dikutip pada teks, harus terdata pada Daftar Pustaka manuskrip, begitu pula sebaliknya.
- Sitasi literatur pada teks mengikuti sistem "nama dan tahun":
 1. Untuk seorang penulis: Hamzah (2013) atau (Hamzah, 2013)
 2. Untuk dua orang penulis: Hamzah dan Amran (2013) atau (Hamzah dan Amran, 2013)
 3. Tiga atau lebih penulis: Hamzah et al. (2013) atau (Hamzah et al., 2013)
 4. Menggunakan manuskrip yang telah diterima untuk dipublikasi: Hamzah (in press) atau (Hamzah, in press).
 5. Sangat diharapkan untuk menghindari penggunaan data atau komunikasi dengan orang untuk dimasukkan ke dalam manuskrip. Bila itu dirasa sangat perlu, maka dituliskan: A. Hamzah (Universitas Hasanuddin, unpublished data) atau A. Hamzah (Universitas Hasanuddin, komunikasi pribadi).
 6. Bila menggunakan multisitasi, gambar dan tabel, gunakan semicolon: (Hamzah, 2013; Amran,2010) (Tabel 1; Gambar 2).
- Manuskrip tersusun atas: halaman judul, halaman abstrak, teks (pendahuluan-simpulan/saran), daftar kepustakaan, daftar tabel dan gambar yang ada, tabel

II. Petunjuk Lainnya dan Template

Dapat dilihat dan diunduh di Website Jurnal Ilmu Kelautan SPERMONDE: <http://journal.unhas.ac.id/index.php/jiks>
atau dapat menghubungi Dewan Redaksi di Email: journalspermonde@mar-sci.unhas.ac.id

Jurnal Ilmu Kelautan

SPERMONDIE

SPERMONDE Vol. 4 No. 1 Hal. 1-47 Makassar, Maret 2018

ISSN: 2460-0156

EISSN: 2614-5049

- Alimuddin Hamzah Assegaf, Wasir Samad, Sakka* 1-6
Some Characteristics of Atmospheric Boundary Layer Over Makassar
- Alinda. N. Hasanah, Nita Rukminasari, Budiman Yunus, Dewi Yanuarita, Jamaluddin Jompa, Suharto* 7-11
The Effect of Temperature on Zooxanthellae of Isopora Palifera and Acropora Hyacinthus From Karanrang Island, Indonesia
- Chair Rani, M. Natsir Nessa, Jamaluddin Jompa, Ahmad Faisal, Shinta Werorilangi, Akbar Tahir* 12-19
Temporal Dynamics of Eutrophication Level and Sedimentation Rate in Coral Reef Area of Spermonde and Sembilan Islands, South Sulawesi
- Esa Fajar Hidayat, Sri Pujiyati, Ali Suman, Totok Hestirianoto* 20-25
Distribution of Pelagic Fish In South China Sea Using Geostatistical Approach
- Magdalena Litaay, Slamet Santosa, Eva Johannes, Rosana Agus, Willem Moka, Jennyta Dhewi Darmansyah Tanjung* 26-31
Biodiversity of Marine Tunicates at Samalona Waters, Sangkarang Archipelago, Indonesia
- Muhammad Banda Selamat, Mahatma Lanuru, Amir Hamzah Muhiddin* 32-38
Spatial Composition of Benthic Substrate around Bontosua Island
- Muh. Farid Samawi, Shinta Werorilangi, Rahmadi Tambaru, Rastina* 39-42
Accumulation of Heavy Metals in Hard Coral *Porites lutea* at Spermonde Archipelago, South Sulawesi
- Wasir Samad, Muhammad Anshar Amran, Amir Hamzah Muhiddin, Rahmadi Tambaru* 43-47
Spatial-Temporal Distribution of Chlorophyll-a in The Southern Part of The Makassar Strait



BIODIVERSITY OF MARINE TUNICATES IN SAMALONA WATERS, SANGKARANG ARCHIPELAGO, INDONESIA

ORIGINALITY REPORT

% **3**

SIMILARITY INDEX

% **1**

INTERNET SOURCES

% **1**

PUBLICATIONS

% **2**

STUDENT PAPERS

PRIMARY SOURCES

1

www.thieme-connect.de

Internet Source

% **1**

2

Submitted to University College London

Student Paper

<% **1**

3

Miura, M.. "Effects of sprouting and canopy states on the structure and dynamics of a *Castanopsis cuspidata* var. *sieboldii* sapling population in an old-growth evergreen broad-leaved forest", *Forest Ecology and Management*, 20030915

Publication

<% **1**

4

adc.bmj.com

Internet Source

<% **1**

5

www.int-res.com

Internet Source

<% **1**

6

Submitted to University of the Philippines Los Banos

Student Paper

<% **1**

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES

< 5
WORDS