

8. Sustainability approach in cropping intensity (CI) 400 through optimizing the dosage of compost and chemical fertilizers to early-maturing rice varieties based on multivariate analysis

by master 11

Submission date: 22-Feb-2024 09:50AM (UTC+0700)

Submission ID: 2245235076

File name: 8_1-s2.0-S2666154323004143-main.pdf (1.94M)

Word count: 11062

Character count: 55989



Sustainability approach in cropping intensity (CI) 400 through optimizing the dosage of compost and chemical fertilizers to early-maturing rice varieties based on multivariate analysis

Yunus Musa^a, Muh Farid^{a,*}, Nasaruddin Nasaruddin^a, Muhammad Fuad Anshori^a, Ahmad Fauzan Adzima^b, Muhammad Farid Maricar^c, Amran Sulaiman^d, Yi Cheng Huang^e, Hari Iswoyo^a, Abd Haris Bahrhun^a, Adnan Adnan^f

^a Department of Agronomy, Hasanuddin University, Makassar, 90245, Indonesia

^b Department of Soil Science, Faculty of Agriculture, Hasanuddin University, Makassar, 90245, Indonesia

^c Department of Civil Engineering, Faculty of Engineering, Hasanuddin University, Makassar, 90245, Indonesia

^d Department of Agribusiness, Faculty of Agriculture, Hasanuddin University, Makassar, 90245, Indonesia

^e Tainan International Cooperation and Development Fund (Taiwan/CDF), Taipei, 111047, Taiwan

^f Department of Biology, State University of Makassar, Makassar, 90222, Indonesia

ARTICLE INFO

Keywords:

CI 400

Early-maturity varieties

Fertilizer dosage

Multivariate analysis

Sustainable rice

ABSTRACT

The development of early maturing rice varieties in the cropping intensity 400 program is one of the alternatives in overcoming the challenges of rice production against land conversion and dynamic climate change. Optimization of the program is feasible through composting and reduction of chemical and synthetic substances in order to avoid excessive soil damage. Therefore, evaluation in the combination of compost and chemical fertilizers is important with evaluation criteria through multivariate analysis. This research aimed to identify evaluation criteria, fertilizer packages and potential of early maturing varieties in supporting the cropping intensity 400 concept on rice. This research was conducted in two locations in South Sulawesi, Indonesia with different environmental conditions ... This study employed a nested split-plot design with a group randomized design as the environmental design. The main plot was the fertilizer package (FP) consisting of 3 levels. The subplots consisted of 7 rice varieties. Each combination of fertilizer package and variety was repeated three times. The results showed that reducing the dose of NPK fertilizer together with compost application is generally effective in the growth of early maturing rice. This study also revealed the characters of chlorophyll *a*, harvest age, panicle length, 1000-grain weight and yield were effective in evaluating the effect of fertilizer packages on early maturing varieties. The 75 % dose of N:P:K and 3 tons.ha⁻¹ of compost is effective fertilizer packages to the growth and production of early maturing rice in supporting cropping intensity 400. The Cakrabuana, and Inpari 13 varieties are good early maturing varieties used with the FP3 fertilization packages. Conclusively, the results of this research are important recommending optimization of rice cultivation with the CI 400 concept, especially in the South Sulawesi region.

1. Introduction

Rice is the main food and has been the main source of carbohydrates for most of the world's population, especially in Asia. Asia dominates the top five largest rice production countries by contributing 72.6 % of the world production. These countries are China (29.5 %), India (23.8

%), Bangladesh (7.0 %), Indonesia (6.9 %), and Vietnam (5.4 %) [1]. Moreover, the development of rice in Asia has always been a priority issue for the sake of food security. In Indonesia, rice production has experienced a relatively fluctuating increase every year [2]. However, this increase in production is still considered insignificant and unstable, consequently, Indonesia occasionally must import to cover the shortage

* Corresponding author.

E-mail addresses: 54.yunusmusa@gmail.com (Y. Musa), farid_deni@yahoo.co.id (M. Farid), nasaruddin@gmail.com (N. Nasaruddin), fuad.anshori@unhas.ac.id (M.F. Anshori), fauzanadzima@unhas.ac.id (A.F. Adzima), farid.river@unhas.ac.id (M.F. Maricar), andiamransulaimanmp@gmail.com (A. Sulaiman), fred10141988@gmail.com (Y.C. Huang), hariiswoyo@unhas.ac.id (H. Iswoyo), harisbahrhun@yahoo.com (A.H. Bahrhun), adnan@unm.ac.id (A. Adnan).

<https://doi.org/10.1016/j.jafr.2023.100907>

Received 12 September 2023; Received in revised form 14 November 2023; Accepted 1 December 2023

Available online 6 December 2023

2666-1543/© 2023 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

of rice demand [2–4]. This phenomenon is caused by the rapid development and population growth every year, which has an impact on the decrease in rice planting area [5–7]. In addition, the challenge of climate change is also an obstacle for stable rice production [5]. Climate change can trigger various abiotic and biotic stresses that can be detrimental and result in crop failure in various regions [8–10]. Therefore, optimizing solutions to increase rice production in Indonesia always become priority effort to adapt to these two challenges.

The assembly of early maturing rice varieties is one option to face the challenges of development and climate change. Early maturing varieties yields under 100 days after planting; hence they are potential for optimizing the planted area per year [11–13]. According to Sutardi et al. [2], the development of early maturing varieties can optimize the potential harvest index four times per year (cropping intensity (CI) 400) in a farm. It enables the farmland to be more productive in supporting food security and the economy. In addition, early maturing varieties are alternative tolerance to avoid and minimize the impact of climate change [11,13]. Some early maturing rice varieties that have been developed are Inpari 13, Inpari 19, Cakrabuana, Padjajaran, and M70D. The five varieties have been reported to be optimal for achieving CI 400 in Indonesia, so they have become a favorite choice in supporting rice production in several regions [14–17]. However, the development of CI 400 on these varieties does not guarantee positive response. There are several issues that are taken into consideration in the optimization of CI 400. One of them is the aspect of environment and sustainability.

Intensive cultivation under the CI 400 concept causes soil properties to decline, especially physical [18–20] and biological [21–23]. Intensive cultivation makes tillage more frequent [18]. This causes soil structure to deteriorate and organic matter to decrease [19,20,24,25], so that high clay soils become increasingly sticky during the rainy season and hard during the dry season [26]. These conditions make it difficult for rice crops to grow and produce [24]. In addition, the process of mineralization and decomposition by microorganisms becomes faster, so that the potential of ecosystems and nutrients in supporting plant growth cannot be optimized [23,27]. Therefore, optimization through the concept of cultivation needs to be supportive to the genetic potential of early maturing varieties in the CI 400 concept [28].

The use of compost is one solution to improve soil quality and plant growth. Compost is the result of residual organic matter decomposed by soil microorganisms [28–30]. The description of organic matter can improve soil aggregates, so that the soil structure becomes crumblier with an optimal proportion between soil micro and macro pores [19,20,31,32]. In addition, the compost will enrich the availability of essential nutrients that are easily absorbed by plants, especially microelements [18,27,30,33]. The decomposition that occurs will also indirectly multiply the colonies of decomposer microorganisms, so that when compost is applied it will further enrich the diversity of soil microorganisms that support the decomposition process in soil organic matter [27,34,35]. These advantages are expected to strengthen the concept of CI 400 [19], especially when combined with a reduction in chemical fertilizer doses. The application of a combination of compost and reduced doses of chemical fertilizers has also been reported to be effective in several rice varieties [2,32,36,37]. However, these tests on early maturing varieties have not been reported comprehensively. Therefore, it is necessary to apply a combination of compost and chemical fertilizer doses to support the growth and production of early maturing rice varieties.

Evaluation of a combination of compost and chemical fertilizer dose-based fertilizer packages requires a systematic approach. The evaluation needs to consider the environmental effects of the study, so testing more than one environment is necessary [36,37]. However, multi-environment-based evaluation leads to a more comprehensive and complex level of data complexity. This requires a precise approach in assessing such evaluations, one of which is through multivariate analysis. Multivariate analysis can simplify, describe, categorize and reduce data with large dimensions to be more easily understood [38,39]. This analysis is very instrumental in determining potential character traits

that represent the diversity of data to be examined in the evaluation process [39,40]. Several studies have reported the effectiveness of the multivariate analysis concept in the evaluation and selection process, both for strains [39,41–43] and combinations of cultivation technology packages [30,44]. Therefore, the use of multivariate analysis becomes necessary to be applied in the evaluation of fertilizer packages of compost and chemical fertilizer doses on the growth and production of early maturing rice. This concept will determine optimal evaluation characteristics in assessing the potential interaction between compost and chemical fertilizers on early maturing varieties, both in general and specific terms. This means the results of this interaction can be recommended as a fertilizer basic for cultivating early maturing varieties in supporting CI 400. The purpose of this study is to identify evaluation criteria fertilizer packages and early maturing varieties that are potential in supporting rice growth and production, especially in the CI 400 concept.

2. Materials and methods

This research was conducted in two locations with different environments from August to November 2022. The first location was Tangke Bajeng village, Limbong sub-district, Gowa Regency, South Sulawesi province, Indonesia at an altitude of 20 m above sea level, with coordinates 5°18'03.98"S 119°26'30.85"E. Tropical climate type which has a temperature of 30°-33 °C with andosol soil type. The second location was Apala village, Barebbo sub-district, Bone Regency at coordinates 4°36'40"S 120°17'52"E. Precipitation of the two locations were shown in Table 1. The selection of the two locations was based on the representation of climate zones in South Sulawesi, namely the western and eastern zones. Apart from that, both had farmers who are experts in aspects of cultivation, including organic farming. Therefore, these two optimal environments were applied as a first step in optimizing crop intensity 400. Table 2 showed the description of both environments.

2.1. Research design

This study employed a nested split-plot design, with 7 replicates were nested within locations, with a group randomized design as the environmental design. The main plot was the fertilizer package (FP) consisting of 5 levels, namely P1 = Nitrogen (N), Phosphorus (P), and Potassium (K) (200:100:100), P2 = NPK (200:100:100) + 2 tons.ha⁻¹ of compost, P3 = 3/4 NPK (150:75:75) + 3 tons.ha⁻¹ of compost, P4 = 1/2 NPK (100:50:50) + 4 tons.ha⁻¹ of compost, and P5 = 1/4 NPK (50:25:25) + 5 tons.ha⁻¹ of compost. The subplots consisted of 7 rice varieties (V), namely Padjajaran (V1), Cakrabuana (V2), Inpari 13 (V3), Inpari 19 (V4), M70D (V5), Ciherang (V6), and Inpari 32 (V7). Each combination of fertilizer package and variety was repeated three times, resulting in 105 experimental units. Each experimental unit was planted with a plot size of 3.5 m × 3.5 m.

2.2. Research procedure

The research procedure followed the general procedure in rice cultivation. First, soil preparation and tillage were carried out by tamping and plowing the land with a tractor. After that, beds were set for the nursery of each variety. The nursery started by soaking the rice seeds

Table 1
Precipitation of both environments (Bone and Gowa) during study in 2022.

| Environment | Precipitation (mm) in 2022 | | | |
|-------------|----------------------------|-----------|---------|----------|
| | August | September | October | November |
| Bone | 231 | 297 | 296 | 80 |
| Gowa | 41.7 | 47.9 | 389.6 | 337.1 |

Table 2
Description of experimental locations.

| Parameter | Unit | Bone | | Gowa | |
|--------------------------|-------------------------------------|------------|-------|--------|-------|
| | | Before | FP3 | before | FP3 |
| Texture | | Dusty Clay | | Clay | |
| Clay | % | 37 | 41 | 13 | 18 |
| Dust | % | 53 | 56 | 40 | 46 |
| Sand | % | 10 | 8 | 47 | 45 |
| C-Organic | % | 1.89 | 2.06 | 0.72 | 0.98 |
| Total Nitrogen | % | 0.15 | 0.17 | 0.13 | 0.15 |
| pH | | 6.17 | 6.75 | 6.24 | 6.35 |
| P Olsen | cmol (+) kg ⁻¹ | 9.1 | 12.9 | 31 | 32 |
| Ca | Ppm cmol (+) kg ⁻¹ | 0.24 | 0.24 | 0.14 | 1.56 |
| Mg | cmol (+) kg ⁻¹ | 6.01 | 8.75 | 7.96 | 12.42 |
| Na | cmol (+) kg ⁻¹ | 2.51 | 3.01 | 3.3 | 3.39 |
| Cation Exchange Capacity | me 100g ⁻¹ | 0.078 | 0.048 | 0.02 | 0.018 |
| | | 20.46 | 21.42 | 11.42 | 18.36 |

with clean water 4: 24 h and then spread on the seedbeds according to the variety label. After the seedlings were 15 days old, they were planted with a spacing of 20 cm × 20 cm in each plot, where each planting point consisted of 3 seedlings. The seedlings were then maintained by replanting, weeding, irrigating, fertilizing, and controlling pests and diseases.

Replanting was performed at the age of 7–14 days after transplanting (DAP). Dead and discarded plants were replaced with new plants from the rest of the nursery according to the variety. Weeding was performed at 15 and 30 days after transplanting by pulling weeds and applying herbicides. Irrigation was carried out 20 days after transplanting (6 days after initial fertilization), water was flowed into the experimental field until the water level reached ± 5 cm. Irrigation was stopped at the time of the second fertilization and the soil was allowed to heavily saturated but un-inundated (local term: macak-macak). Watering was done again 5 days after fertilization. At the age of ±45 DAP, water level was raised to ±10 cm from the soil surface to suppress the growth of new tillers.

Fertilization was done by applying compost and Phonska® NPK fertilizer combined with Urea adjusted to the dose of the fertilization package treatment. Compost was given 2 weeks before planting according to the treatment. Meanwhile, NPK fertilizer was given 2 times in stages. The first fertilizer was given after 14 DAP of planting and the second fertilizer was given after 45 DAP according to the treatment dose. Pest and disease control was carried out in the morning by spraying Regent® insecticides with fipronil 50sc active ingredient when an attack occurred. Rat pest control is carried out by mechanical control by destroying rat nests around rice field bunds and applying rodenticides made from active kumatetralil 0.75tp. Disease control was applied by spraying each experimental plot evenly with Explore® Fungicide with Difenoconazole active ingredient at the age of ±45–60 DAP.

Harvesting was done when 2/3 of the panicle has entered the physiological maturity phase (yellowing straw) and the rice grains at the base of the panicle have hardened. Harvesting was performed manually with a sickle and threshed using a rice thresher. Sample plants were harvested first, be 15: harvesting the rest of the plot. The observed characters include plant height, number of productive tillers, harvest 15: chlorophyll a, chlorophyll b, panicle length, total grain number, percentage of filled grain, flag leaf length, 1000-grain weight, and the yield.

2.3. Data analysis

The determination of evaluation criteria was based on three multivariate analyses, namely factor analysis, PCA biplot analysis, and cluster

gram heatmap. The use of the three analyses together has not been widely reported in determining evaluation criteria. However, some research reports show the effectiveness of each analysis in determining selection or evaluation criteria. In general, factor analysis can reduce the diversity of big data to parameters with low internal covariance in influencing a multivariate dimension [39,45,46]. This advantage is very instrumental in determining characters that have a variety of characteristics in the process of evaluating treatment combinations [38,39]. The reduction is further sharpened by PCA biplot and clustergram analysis. Both analyses are often used together in the identification of clustering of objects and parameters [47–50]. PCA biplot analysis can show how the direction and magnitude of variance of one parameter is with respect to another in two dimensions [50]. This makes it possible to see the relationship and clustering of variance between characters in a multivariate manner. In addition, clustering is further supported by clustergram heatmap analysis. This analysis is able to map and visualize simply the grouping pattern of interactions between objects and parameters based on the accumulation of similarity of the middle value, so that information on the parameters that determine the grouping of objects can be identified [49,51]. The combination of the two analyses can recommend characters that can effectively reflect the diversity of data in the evaluation process. Therefore, the combination of the three analyses in determining 12: evaluation criteria is considered effective, especially in evaluating the effect of fertilizer packages on the growth and production of early maturing rice. The use of the three analyses utilized Rstudio 3.6.1 software with factoextra and gplots package.

The predetermined evaluation 26: criteria were deepened by further analysis based on the results of analysis of variance (ANOVA). Further analysis used the least significant distance (LSD) test at the 5 % error level. The evaluation was more focused on the source of diversity of the interaction, especially the interaction between the three. However, characters that do not have significant interactions between the three factors will continue with further tests on the interaction between the remaining two factors. ANOVA and further tests were conducted using STAR 2.0.1 software.

3. Results

3.1. Identification of agronomy evaluation criteria toward fertilizer technology package to variety growth

The factor analysis results show a total variance percentage of 70 %. Based on this analysis, there are 4 main factors with different characteristics of the characterizing parameters (Table 3). Factor 1 is characterized by chlorophyll a and chlorophyll b which have variance factor

Table 3

Factor analysis to rice growth and yield component on the combination of fertilizer package and varieties.

| Variable | Factor1 | Factor2 | Factor3 | Factor4 | Communality |
|----------------|---------|---------|---------|---------|-------------|
| PH | 0.131 | 0.23 | 0.101 | 0.018 | 0.665 |
| NPT | 0.218 | -0.04 | -0.004 | -0.01 | 0.42 |
| DH | 0.137 | 0.074 | -0.356 | 0.113 | 0.613 |
| Chl_a | 0.313 | -0.058 | -0.101 | -0.069 | 0.859 |
| Chl_b | 0.309 | -0.059 | -0.052 | -0.047 | 0.833 |
| FFL | 0.067 | 0.279 | 0.028 | -0.398 | 0.686 |
| PL | -0.074 | 0.399 | 0.123 | 0.147 | 0.797 |
| NTG | -0.123 | 0.364 | -0.087 | 0.013 | 0.663 |
| PFG | -0.201 | 0.142 | -0.366 | -0.268 | 0.598 |
| W1000 | -0.008 | 0.023 | 0.082 | 0.817 | 0.836 |
| Yield | 0.026 | -0.096 | -0.711 | -0.059 | 0.805 |
| Variance (Var) | 3.0774 | 2.2928 | 1.2879 | 1.1168 | 7.7749 |
| % Var | 0.28 | 0.208 | 0.117 | 0.102 | 0.7 |

Notes: PH = 10: plant height, NPT = number of productive tillers, DH = days to harvest 2: g, Chl a = chlorophyll a, Chl b = chlorophyll b, FFL = flag leaf length, PL = panicle length, NTG = number of total grains, PFG = percentage of filled grains, W1000 = Weight of 1000 grains.

loading values of 0.313 and 0.309, respectively. Factor 2 is characterized by the characters of panicle length and total grain number which have variance factor loading values of 0.399 and 0.364, respectively. Factor 3 is characterized by three characters namely harvest time, percentage of filled grains, and the yield. The three characters have loading values of -0.356 , -0.366 , and -0.711 , respectively. Meanwhile, Factor 4 is only characterized by one character i.e. 1000-grain weight, with a loading factor value of 0.817.

The clustering of selected characters in the factor analysis was done by principal component analysis (PCA) and clustergram heatmap. Based on the PCA (Fig. 1), the yield had a low variance contribution compared to the eight selected characters in the factor analysis. In contrast, the highest contribution of diversity belonged to chlorophyll *a* and *b*, but the variance of both was not in the same direction as the variance of the yield. The characters of total grain number, panicle length, harvest age, and 1000-grain weight have the same direction of variation as the yield. Among these characters, total grain number and panicle length have a large degree of diversity in the group.

Based on the clustergram heatmap analysis (Fig. 2), there are three main groups that divide the grouping of treatment combinations. The first group consisted of yield, panicle length and 1000-grain weight. The second group consisted of percentage of filled grains, chlorophyll *b*, harvest time and total grain number. Meanwhile, the third group consisted of only one member of the growth character i.e. chlorophyll *a*.

3.2. Evaluation effectiveness of fertilizer technology package to variety growth based on selected agronomy evaluation criteria

The results of analysis of variance (ANOVA) showed that chlorophyll *a* had an interaction among location, fertilizer package and variety (Table 4). In the first location (Bone), FP2 and FP3 were the most varied fertilizer packages among varieties, while FP1 and FP5 had relatively

unchanged responses among early maturing varieties. Variety of Padjajaran had the highest chlorophyll in FP2, but not significantly different from Cakrabuana, Inpari 13, and Inpari 19. Inpari 13 had the highest chlorophyll in FP3 but was not significantly different from almost all varieties, except Padjajaran. Based on the second location (Gowa), FP1 and FP3 were fertilizer packages with different response on chlorophyll *a* among early maturing varieties. For both fertilizer packages, Ciherang remained the variety with the highest chlorophyll content i.e. 234.68 for FP1 and 230.35 for FP3.

The results of ANOVA showed that harvest time had a significant interaction between location and fertilizer package, as well as between location and variety (Table 5). The fertilizer package treatment in Bone showed that the FP4 and FP5 treatments had a longer harvest age than the FP1, FP2, and FP 3 treatments. On the other hand, in Gowa, all fertilizer packages showed the same harvest time. In addition, based on the source of variance, variety M70D had the fastest average harvest time in both Bone (91.3 days) and Gowa (83.00 days).

The results of ANOVA for panicle length showed interaction among environment, fertilizer package and variety (Table 6). In Bone, FP2 and FP5 showed very varied responses compared to other treatments. However, the FP3 package still has good potential in explaining the diversity of responses between varieties based on panicle length characteristics. For the FP2, The Padjajaran variety was the variety with the longest panicles, while for FP5, the Inpari 19 was variety with the longest panicles compared to other varieties. Although, these results were not significant compared to most varieties. Meanwhile, in Gowa, FP1 and FP3 were optimal technology packages in explaining the diversity of responses of each variety to fertilizer treatment. At the FP1 level, Inpari 19 variety had the longest panicles, although the responses were not significant compared to most the varieties. At the FP3 level, Ciherang was the variety with the longest panicles, but it was not significantly different from Inpari 19 and Inpari 32.





Fig. 1. Principal component biplot analysis on selected characters based on factor analysis. (DH = days to harvesting, Chl *a* = chlorophyll *a*, Chl *b* = chlorophyll *b*, PL = panicle length, NTG = number of total grains, PFG = percentage of filled grains, W1000 = Weight of 1000 grains).



2. heatmap clustergram analysis to the combination of fertilizer package and varieties based on selected characters of factor analysis. (DH = days to harvesting, Chl a = chlorophyll a, Chl b = chlorophyll b, PL = panicle length, NTG = number of total grains, PFG = percentage of filled grains, W1000 = Weight of 1000 grains).

Table 4
Chlorophyll a performance is based on interaction of environment, fertilizer package and variety.

| Variety | Fertilizer Packages | | | | | | | | | |
|------------|---------------------|-------|--------|-------|--------|-------|--------|-------|--------|-------|
| | FP1 | | FP2 | | FP3 | | FP4 | | FP5 | |
| | chl a | notif | chl a | notif | chl a | notif | chl a | notif | chl a | notif |
| Bone | | | | | | | | | | |
| Padjajaran | 277.25 | p | 262.65 | p | 238.44 | q | 314.32 | p | 258.65 | p |
| Cakrabuana | 274.14 | p | 236.9 | pqr | 245.81 | pq | 255.39 | q | 249.43 | p |
| Inpari 13 | 258.21 | p | 259.77 | pq | 271.63 | p | 249.44 | q | 250.97 | p |
| Inpari 19 | 252.39 | p | 248.65 | pqr | 262.87 | pq | 311.55 | p | 238.92 | p |
| M70D | 255.75 | p | 228.96 | r | 259.42 | pq | 244.63 | q | 243.6 | p |
| Ciherang | 268.6 | p | 229.38 | qr | 255.79 | pq | 273.47 | q | 256.45 | p |
| Inpari 32 | 259.19 | p | 254.91 | pqr | 262.62 | pq | 258.72 | q | 248.04 | p |
| Gowa | | | | | | | | | | |
| Padjajaran | 166.23 | bcd | 211.29 | a | 225.02 | ab | 129.22 | c | 133.09 | c |
| Cakrabuana | 136.94 | d | 164.63 | b | 196.79 | bc | 179.13 | b | 174.25 | b |
| Inpari 13 | 176.07 | bc | 201.65 | a | 154.75 | d | 124.64 | c | 139.83 | c |
| Inpari 19 | 158.45 | cd | 189.21 | ab | 172.04 | cd | 171.1 | b | 173.79 | b |
| M70D | 139.39 | d | 159.74 | b | 195.69 | bc | 134.66 | c | 144.6 | bc |
| Ciherang | 234.68 | a | 202.7 | a | 230.35 | a | 188.14 | b | 139.84 | c |
| Inpari 32 | 196.74 | b | 198.58 | a | 186.96 | c | 237.13 | a | 243.92 | a |

Notes: FP = fertilizer package, chl a = chlorophyll a, notif = notifications, the same letter in one column has same response.

The results of ANOVA of weight of 1000-grain showed that there was an interaction among location, fertilizer package and variety (Table 7). The fertilizer package treatments had relatively similar patterns of variation among genotypes. However, FP1, FP2 and FP3 were fertilizer packages that supported a potential weight of 1000 grains. In Bone, Padjajaran was the variety with the heaviest 1000-grain weight in the FP1 treatment. For FP2, Inpari 13 was the variety with the heaviest 1000-grain weight and for FP3, Inpari 32 had the heaviest 1000-grain weight. However, the best varieties in each FP were not significantly different from almost all varieties. Meanwhile, in Gowa, Inpari 32 was the variety with the heaviest 1000-grain weight at the FP1 level. However, this weight was not significantly different from all varieties, except

Inpari 13. At the FP2 level, Padjajaran was the variety with the heaviest 1000-grain weight and was not significantly different from all varieties, except Cakrabuana. At FP3 level, M70D was the variety with the heaviest 1000-grain weight, but was not significantly different from Padjajaran, Inpari 13 and Ciherang.

The ANOVA of the yield characters showed that there was a significant interaction among environment, fertilizer packaging and variety (Table 8). For this character, all levels of fertilizer packages had relatively similar varying response patterns. However, FP1, FP2 and FP3 were fertilizer packages with good productivity potential. In Bone, Inpari 13 (6.51 tons ha⁻¹, 7.38 to 14 a-1), Ciherang (7.89 tons ha⁻¹, 8.11 tons ha⁻¹), and Inpari 32 (7, 22 tons ha⁻¹, 7.55 tons ha⁻¹) showed

Table 5

Days to harvesting performance is based on interaction of environment x fertilizer package and environment x variety.

| Treatment | Bone | | Gowa | |
|--------------------|-------|-------|-------|-------|
| | Days | notif | Days | notif |
| Fertilizer Package | | | | |
| FP1 | 100.9 | q | 97.6 | p |
| FP2 | 100.9 | q | 97.6 | p |
| FP3 | 101.0 | q | 97.6 | p |
| FP4 | 101.4 | p | 97.6 | p |
| FP5 | 101.5 | p | 97.6 | p |
| Variety | | | | |
| Padjajaran | 97.4 | d | 95.0 | b |
| Cakrabuana | 97.1 | d | 95.0 | b |
| Inpari 13 | 99.1 | c | 95.0 | b |
| Inpari 19 | 93.2 | e | 95.0 | b |
| M70D | 91.3 | f | 83.0 | c |
| Ciherang | 117.0 | a | 110.0 | a |
| Inpari 32 | 112.9 | b | 110.0 | a |

Notes: FP = fertilizer package, notif = notifications, the same letter in one column has same response.

good yield response to FP1 and FP2 fertilizer packages. Inpari 32 variety (7.33 tons ha⁻¹) had the highest yield in FP3 but was not significantly different from Cakrabuana (5.71 tons ha⁻¹), Inpari 13 (6.56 tons ha⁻¹), and Ciherang (7 0.11 tons ha⁻¹). Meanwhile, based on the results in Gowa, all varieties had almost the same relative response to each fertilizer package regarding yield. For FP1 fertilizer package, Inpari 19 variety (4.81 tons ha⁻¹) showed the highest yield response, but was not significantly different from all other varieties. For FP2, FP3 fertilizer packages, the Cakrabuana variety (7.05, and 8.20-tons ha⁻¹) was the variety with the highest yield, but was not significantly different from the Padjajaran variety (6.56, 7.46-, and 5.81-tons ha⁻¹).

4. Discussion

Based on factor analysis, PCA biplot, and clustergram heatmap; the characters of harvest time, chlorophyll *a* content, panicle length, 1000-grain weight and the yield are characters that can be used in evaluating the effect of fertilizer packages on the growth and production of early maturing rice. The five characters meet the standards of characterization on factor analysis, 24% with the characters of chlorophyll *b* content, total grain number, and percentage of filled grains. The determination is based on the loading factor value that exceeds 0.3. This is in line with the

Table 6

Panicle length performance is based on interaction of environment, fertilizer package and variety.

| Variety | Fertilizer Packages | | | | | | | | | |
|------------|---------------------|-------|---------|-------|---------|-------|---------|-------|---------|-------|
| | FP1 | | FP2 | | FP3 | | FP4 | | FP5 | |
| | PL (cm) | notif | PL (cm) | notif | PL (cm) | notif | PL (cm) | notif | PL (cm) | notif |
| Bone | | | | | | | | | | |
| Padjajaran | 24.82 | p | 26.24 | p | 23.81 | pq | 24.28 | p | 24.11 | pqr |
| Cakrabuana | 21.59 | q | 24.3 | pqr | 25.92 | p | 23.26 | p | 21.96 | qr |
| Inpari 13 | 24.05 | pq | 24.68 | pq | 24.33 | pq | 25.00 | p | 24.5567 | pq |
| Inpari 19 | 22.38 | pq | 24.81 | pq | 23.63 | pq | 25.62 | p | 25.5133 | p |
| M70D | 23.47 | pq | 21.39 | r | 22.61 | q | 24.45 | p | 21.2733 | r |
| Ciherang | 24.48 | pq | 21.99 | qr | 23.38 | pq | 22.81 | p | 22.4667 | pqr |
| Inpari 32 | 24.41 | pq | 24.61 | pq | 25.88 | p | 22.98 | p | 24.9667 | pq |
| Gowa | | | | | | | | | | |
| Padjajaran | 22.11 | c | 24.55 | ab | 21.66 | bc | 24.22 | ab | 23.77 | a |
| Cakrabuana | 23.89 | abc | 22.66 | b | 19.55 | c | 22.66 | b | 22.77 | a |
| Inpari 13 | 23.11 | abc | 24.11 | ab | 23.33 | b | 23.44 | b | 25.11 | a |
| Inpari 19 | 25.88 | a | 24.33 | ab | 24.11 | ab | 26.89 | a | 23.55 | a |
| M70D | 22.33 | bc | 24.44 | ab | 23.11 | b | 22.55 | b | 24.11 | a |
| Ciherang | 25.33 | ab | 26.67 | a | 26.56 | a | 24.88 | ab | 23.33 | a |
| Inpari 32 | 24.67 | abc | 24.89 | ab | 24.66 | ab | 24 | ab | 22.33 | a |

Notes: FP = fertilizer package, PL = panicle length, notif = notifications, the same letter in one column has same response.

Table 7

Weight of 1000 grains performance is based on interaction of environment, fertilizer package and variety.

| Variety | Fertilizer Packages | | | | | | | | | |
|------------|---------------------|-------|-----------|-------|-----------|-------|-----------|-------|-----------|-------|
| | FP1 | | FP2 | | FP3 | | FP4 | | FP5 | |
| | W1000 (g) | notif | W1000 (g) | notif | W1000 (g) | notif | W1000 (g) | notif | W1000 (g) | notif |
| Bone | | | | | | | | | | |
| Padjajaran | 28.07 | p | 27.43 | pq | 25.33 | q | 24.33 | pq | 26.90 | pq |
| Cakrabuana | 25.50 | pq | 27.63 | pq | 24.43 | q | 27.33 | p | 24.63 | pq |
| Inpari 13 | 24.17 | pq | 29.13 | p | 25.63 | q | 22.87 | q | 23.84 | qr |
| Inpari 19 | 22.50 | q | 26.33 | pq | 26.37 | q | 27.27 | p | 19.67 | s |
| M70D | 22.90 | q | 23.93 | q | 27.40 | pq | 22.93 | q | 20.43 | rs |
| Ciherang | 26.23 | pq | 27.33 | pq | 27.83 | pq | 27.47 | p | 23.37 | qrs |
| Inpari 32 | 25.60 | pq | 27.60 | pq | 30.77 | p | 26.23 | pq | 28.20 | p |
| Gowa | | | | | | | | | | |
| Padjajaran | 29.11 | ab | 31.11 | a | 29.44 | ab | 27.44 | ab | 30.11 | a |
| Cakrabuana | 29.33 | ab | 26.66 | b | 25.55 | b | 23.55 | b | 24.11 | b |
| Inpari 13 | 26.33 | b | 30.89 | a | 28.89 | ab | 28.89 | a | 27.33 | ab |
| Inpari 19 | 27.00 | ab | 28.66 | ab | 25.89 | b | 26.11 | ab | 25.44 | b |
| M70D | 30.22 | ab | 30.00 | ab | 31.55 | a | 28.00 | a | 26.44 | ab |
| Ciherang | 30.64 | a | 28.64 | ab | 28.30 | ab | 25.64 | ab | 25.30 | b |
| Inpari 32 | 30.97 | a | 28.97 | ab | 26.97 | b | 26.64 | ab | 26.30 | ab |

FP = fertilizer package, W1000 = weight of 1000 grains, notif = notifications, the same letter in one column has same response.

Table 8
The yield performance is based on interaction of environment, fertilizer package and variety.

| Variety | Fertilizer Packages | | | | | | | | | |
|------------|---------------------|-------|--------------|-------|--------------|-------|--------------|-------|--------------|-------|
| | FP1 | | FP2 | | FP3 | | FP4 | | FP5 | |
| | Yield (t/ha) | notif | Yield (t/ha) | notif | Yield (t/ha) | notif | Yield (t/ha) | notif | Yield (t/ha) | notif |
| Bone | | | | | | | | | | |
| Padjajaran | 5.71 | qrs | 5.18 | q | 4.98 | q | 4.87 | pq | 3.87 | q |
| Cakrabuana | 5.31 | rs | 5.35 | q | 5.71 | pq | 5.29 | pq | 4.78 | pq |
| Inpari 13 | 6.51 | pqr | 7.38 | p | 6.56 | pq | 6.47 | p | 5.98 | p |
| Inpari 19 | 4.22 | s | 5.51 | q | 5.29 | q | 4.33 | q | 4.67 | pq |
| M70D | 4.89 | rs | 5.38 | q | 5.00 | q | 4.89 | pq | 4.62 | pq |
| Ciherang | 7.89 | p | 8.11 | p | 7.11 | p | 6.56 | p | 6.33 | p |
| Inpari 32 | 7.22 | pq | 7.55 | p | 7.33 | p | 6.33 | p | 6.11 | p |
| Gowa | | | | | | | | | | |
| Padjajaran | 4.50 | a | 6.56 | ab | 7.46 | a | 5.41 | ab | 5.81 | ab |
| Cakrabuana | 4.05 | a | 7.05 | a | 8.20 | a | 6.10 | a | 6.47 | a |
| Inpari 13 | 4.55 | a | 5.07 | bc | 5.49 | b | 4.32 | b | 3.83 | c |
| Inpari 19 | 4.81 | a | 5.15 | bc | 5.60 | b | 4.58 | ab | 4.76 | bc |
| M70D | 4.64 | a | 3.85 | cd | 5.80 | b | 4.42 | b | 4.27 | bc |
| Ciherang | 3.89 | a | 4.15 | cd | 5.12 | b | 5.03 | ab | 4.22 | bc |
| Inpari 32 | 3.61 | a | 3.39 | d | 5.60 | b | 4.52 | ab | 4.33 | bc |

FP = fertilizer package, notif = notifications, the same letter in one column has same response.

concept of Yong and Pieace [52], Farid et al. [42], Sakinah et al. [40], Momen et al. [38] and Anshori et al. [39]. Furthermore, the determination of the five selected characters is based on the interaction of PCA biplot analysis and clustergram heatmap. The selection of 1000-grain weight and panicle length as evaluation criteria is due to the fact that both characters have a strong relationship with the yield as the main evaluation criteria. This is the main supporting basis in the evaluation process in this study. The choice of chlorophyll *a* as an evaluation criterion is due to the wide diversity of this character compared to other characters. Although, the direction of chlorophyll *a* diversity is different from the yield. In addition, this character is independent in the process of grouping objects in clustergram analysis, so the use of chlorophyll *a* can be taken into consideration in the evaluation process. In contrast, chlorophyll *b* was not included as an evaluation criterion, although the diversity of the character was also high and became one of the determinants of object grouping. This is based on the fact that chlorophyll *b* has a pattern and the amount of variation that is relatively the same as chlorophyll *a*. This is ineffective considering that the ratio of chlorophyll *b* is also lower than chlorophyll *a*, which is 3:1 [53,54]. Meanwhile, harvest age was chosen as an evaluation criterion because this character has a diversity that is relatively in the same direction as the yield and has a grouping pattern that is relatively similar to chlorophyll *a* and *b*. In addition, harvest age is also in line with the special characteristics of early maturing rice [11–13], so this character is prioritized as an evaluation criterion. Therefore, these five characters are considered suitable to be the evaluation criteria in this study.

Based on the results of further tests on chlorophyll *a* content, the interaction between the environment, fertilizer package and varieties have a very dynamic pattern. There is no consistent pattern and effect between each combination. However, in general, there are two things that characterize the results of this study on chlorophyll *a* character, namely the influence of the environment and the application of compost fertilizer. The diversity of chlorophyll content patterns is more dynamic in the Bone environment compared to Gowa environment. This is due to the different rainfall between the two environments, especially in the vegetative phases. Rainfall affects the solubility and mobility of elements from soil to plants and plant cell division [55,56]. However, the main aspect that determined the pattern of variation was the fertilizer package. Compost fertilizer application shows a very diverse pattern, especially in supporting chlorophyll content due to changes in chemical fertilizer doses. In general, compost contains complete nutrients for growth, especially in the microelements Fe, Mn and Mo [57–59]. These three elements are very instrumental in supporting the potential chlorophyll content along with nitrogen [60–62]. In addition, compost also

contains many microorganisms that are good for plant growth [63–66]. The advantages of compost make the chlorophyll content pattern more dynamic, especially when combined with chemical fertilizers, such as in FP3 with enough high consistent in both environments. This fertilizer packages were able to relatively increase the potential chlorophyll content in early maturing rice (V1–V5). Based on the overall explanation, the application of compost fertilizer, especially in FP3, can be recommended to increase the chlorophyll content of early maturing rice, especially under optimal rainfall.

Based on the results of the follow-up test, the environment with less rainfall had an earlier harvest time than the environment with more rainfall. This also influenced the insignificant effect of the fertilizer package on harvest time. In contrast, areas with high rainfall affected harvest time, where fertilizer packages with optimized nutrition, such as FP1 and FP2, had a relatively earlier harvest age. In general, harvest time is highly dependent on the heat unit received by the plant [67–70]. In the presence of high intensity rain, the heat unit per day becomes lower [67,68,70], so that the plant growth becomes slower than areas with low rainfall intensity. In addition, the effect of fertilization also affects the optimization of the interaction between nutrients and their heat unit capacity. This makes the harvest age pattern between fertilizer package levels different. However, the optimization does not affect the interaction between the fertilizer package and the variety or in other words, the changes in harvest time are static or additive. In contrast, the harvest time in areas with low rainfall is largely determined by its genetic potential. This causes the effect of fertilizer packages to have no significant effect on harvest time [71,72]. The variety with a consistent early harvest time in both environments is the M70D variety. According to Monica et al. [17] and Rismawati et al. [16], this variety is known to have a super early age, so the harvest age is consistently faster than other varieties in both environments.

Based on the results of further tests on panicle length characters, compost fertilizer application showed the best interaction on several varieties in both environments (Bone and Gowa). The response was clearly visible when compared to the fertilizer package level without compost application (FP1). This pattern also occurred in the 1000-grain weight character. Compost application relatively increased the 1000-grain weight potential compared to no compost application in both environments, especially in the FP2 and FP3 fertilizer packages. In general, compost dosing will affect the potential of yield components, such as panicle length and 1000-grain weight [73–75]. Both characters are highly related as part of the yield components. Several studies have also reported the relationship between panicle length and 1000-grain weight and the closeness of both to the main character of the yield

[39,76], so the interaction pattern of the two characters is relatively similar in this study. Although specifically, the best varieties on both characters are different in the interaction analysis. In the character of panicle length, varieties Inpari 13, and Ciherang are stable potential varieties in FP2 and FP3 fertilization packages in both environments. Conversely, on the character of 1000-grain weight, Varieties Inpari 13, and M70D became potential early maturing rice varieties in fertilization packages FP2 and FP3 in both environments. In addition, the decrease in chemical fertilizer doses on the growth of both characters can still be supported by compost fertilization [30,77,78] especially in FP3. This indicates that compost can make up for the lack of NPK chemical fertilizer doses in early maturing rice varieties. Based on this, the FP2 and FP3 fertilization packages can be recommended in supporting the yield components of early maturing rice varieties.

Based on further tests of the yield character, FP2 and FP3 fertilization packages are the best fertilization packages on both environments. In addition, varieties Cakrabuana and Inpari 13 became early maturing rice varieties with the same or higher yield potential than check varieties Ciherang and Inpari 32 in both fertilizer packages (FP2 and FP3). This result further confirms the application of compost fertilizer to be critical for early maturing rice varieties that have been described in the characters of chlorophyll *a*, panicle length and 1000-grain weight. Compost fertilizers that are rich in micro-nutrients and positive microorganisms for crops will support the production of these crops, despite a decrease in the dose of chemical fertilizers [30,77]. In addition, these results can also strengthen the design of the CI 400 concept in paddy fields, especially in the FP3 fertilization package. Reducing the dose of NPK chemical fertilizer by 25 % with the addition of compost by 3 tons/ha can balance the loss of potential from chemical fertilizers in supporting plant growth and productivity. Furthermore, these results are also supported by changes in the soil physicochemical properties which showed changes in several aspects for the better (Table 2). This fertilization package can also minimize the risk of damage to the physical properties of rice fields in the CI 400 concept [2,29,79]. Based on the explanation above, compost application in FP2 and FP3 is recommended in the cultivation of early maturing rice. Meanwhile, the combination of FP3 fertilization package with Cakrabuana, or Inpari 13 varieties is recommended in supporting rice cultivation in optimizing the CI 400 concept.

21 5. Conclusion

30
The results of this study concluded that the chlorophyll *a*, harvest age, panicle length, weight of 1000 grains, and the yield can be used to increase the effectiveness of agronomical evaluation in early maturity rice variety. FP2 (100 % dose of N:P:K and 2 tons.ha⁻¹ of compost) and FP3 (75 % dose of N:P:K and 3 tons.ha⁻¹ of compost) packages are recommended as effective fertilizer packages in supporting the growth and production of early maturing rice. For FP3 package, it is also effective to be applied in CI 400 planning. Varieties of Cakrabuana and Inpari 13 are early maturing varieties that are well used in FP2 and FP3 fertilization packages. Although this research shows good results, this research has limitations in the number of seasons and locations that do not represent the middle climate zone in South Sulawesi. Further research regarding these limitations needs to be carried out to support an effective policy in optimizing the development of early maturing rice, including the concept of CI 400 in Indonesia.

3 Credit authorship contribution statement

Yunus Musa: Conceptualization, software, validation, formal analysis, writing-original draft preparation.

Muh Farid: Conceptualization, methodology, validation, resources, supervision, and funding acquisition.

Nasaruddin Nasaruddin: validation, data curation, and writing review and editing.

Muhammad Fuad Anshori: validation, data curation, writing-

original draft preparation, formal analysis, and investigation.

Ahmad Fauzan Adzima: validation and investigation.

Muhammad Farid Maricar: validation, and data curation.

Amran Sulaiman: supervision, and funding acquisition.

Yi Cheng Huang: Conceptualization, resources, and funding acquisition.

Hari Iswoyo: validation, translation, and writing review and editing.

Abd Haris Bahrin: writing review and editing.

Adnan Adnan: writing review and editing.

5 Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgements

8
We are grateful to the Ministry of Education, Culture, Research, and Technology of Republic of Indonesia through the scheme of UNHAS Fundamental Research Collaborative with grant number: 00323/UN4.22/PT.01.03/2023.

References

- [1] M.A. Al Mamun, S.A.I. Nihad, M.A.R. Sarkar, M.A. Aziz, M.A. Qayum, R. Ahmed, et al., Growth and trend analysis of area, production and yield of rice: a scenario of rice security in Bangladesh, *PLoS One* 16 (2021), e0261128, <https://doi.org/10.1371/journal.pone.0261128>.
- [2] Sutardi, Y. Apriyana, P. Rejekiingrum, A.D. Alifia, F. Ramadhani, V. Darwis, N. Setyowati, D.E.D. Setyono, Gunawan, A. Malik, S. Abdullah, Muslimin, W. Wibawa, J. Triastono, Yusuf, F.D. Arianti, A.Y. Fadwiwati, The transformation of rice crop technology in Indonesia: innovation and sustainable food security, *Agronomy* 13 (2023) 1–14, <https://doi.org/10.3390/agronomy13010001>.
- [3] A.W. Putra, J. Supriatna, R.H. Koestoer, T.E.B. Soesilo, Differences in local rice price volatility, climate, and macroeconomic determinants in the Indonesian market, *Sustain. Times* 13 (2021), <https://doi.org/10.3390/su13084465>.
- [4] A. Quartina Pudjiastuti, G. Mekse Korri Arisena, A.A. Keswari Krisnandika, Rice import development in Indonesia, *SOCA J. Sos. Ekon. Pertan.* 15 (2021) 390, <https://doi.org/10.24843/soca.2021.v15.i02.p14>.
- [5] I.A. Rumanti, A. Hairmansis, Y. Nugraha, Nafisah, U. Susanto, P. Wardana, R. E. Subandiono, Z. Zaini, H. Sembiring, N.I. Khan, R.K. Singh, D.E. Johnson, A. M. Stuart, Y. Kato, Development of tolerant rice varieties for stress-prone ecosystems in the coastal deltas of Indonesia, *Field Crops Res.* 223 (2018) 75–82, <https://doi.org/10.1016/j.fcr.2018.04.006>.
- [6] A.T. Widayati, Nurbani, W.B. Prasetyo, S.E. Manurung, A.M. Pebriadi, The effect of rice field conversion and its impact on food availability in North Kalimantan Province, *IOP Conf. Ser. Earth Environ. Sci.* 824 (2021), <https://doi.org/10.1088/1755-1315/824/1/012059>, 0–8.
- [7] A.R. Daulay, K.P. Eka Intan, B. Barus, N.B. Pramudya, Rice land conversion into plantation crop and challenges on sustainable land use system in the east tanjung jabung regency, *Procedia - Soc. Behav. Sci.* 227 (2016) 174–180, <https://doi.org/10.1016/j.sbspro.2016.06.059>.
- [8] A. Raza, A. Razzaq, S.S. Mehmood, X. Zou, X. Zhang, Y. Lv, J. Xu, Impact of climate change on crops adaptation and strategies to tackle its outcome: a review, *Plants* 8 (2019), <https://doi.org/10.3390/plants8020034>.
- [9] S.I. Zandalinas, F.B. Fritsch, R. Mittler, Global warming, climate change, and environmental pollution: recipe for a multifactorial stress combination disaster, *Trends Plant Sci.* 26 (2021) 588–599, <https://doi.org/10.1016/j.tplants.2021.02.011>.
- [10] B.K. Singh, M. Delgado-Baquerizo, E. Egidi, E. Guirado, J.E. Leach, H. Liu, P. Trivedi, Climate change impacts on plant pathogens, food security and paths forward, *Nat. Rev. Microbiol.* (2023), <https://doi.org/10.1038/s41579-023-00900-7>.
- [11] Y. Li, X. Wu, W. Xu, Y. Sun, Y. Wang, G. Li, P. Xu, High-Throughput physiology-based stress response phenotyping: advantages, applications and prospective in horticultural plants, *Hortic. Plant J.* 7 (2021) 181–187, <https://doi.org/10.1016/j.hpj.2020.09.004>.
- [12] I. Noviana, Y. Haryati, R. Sari, N. Sunandar, Adaptation to climate change by using drought tolerant and early maturing rice varieties in Majalengka Regency, *IOP Conf. Ser. Earth Environ. Sci.* 648 (2021), <https://doi.org/10.1088/1755-1315/648/1/012118>.

- [13] Y. Yun, Changes in the growth and yield of an extremely early-maturing rice variety according to transplanting density, *Agric. For.* 13 (2023), <https://doi.org/10.3390/agriculture13030717>.
- [14] A. Damiri, T. Hidayat, L. Harta, L. Ivanti, W. Mikasari, A. Afrizon, S.N. Permedi, E. Fauzi, Growth performance of four varieties of rice paddy through PTT technology approach in seluma district, bengkulu province, *PANGAN* 31 (2022) 209–216.
- [15] R. Purba, Y. Giometri, Performance of yields and profits in rice farming with the introduction of varieties in banten, *J. Ilmu Pertanian. Indones.* 22 (2017) 13–19, <https://doi.org/10.18343/jipi.22.1.13>.
- [16] Rismawati, G.R. Sadimantara, Usman, S. Leomo, D. Erawan, S. Samai, Effectiveness of NPK fertilizer on the growth of several new superior upland rice lines (*Oryza sativa* L.) crosses from Southeast Sulawesi, *IOP Conf. Ser. Earth Environ. Sci.* 977 (2022), <https://doi.org/10.1088/1755-1315/977/1/012010>.
- [17] M.F. Danilevicz, P.E. Bayer, B.J. Nestor, M. Bennamoun, D. Edwards, Resources for image-based high-throughput phenotyping in crops and data sharing challenges, *Plant Physiol* 187 (2021) 699–715, <https://doi.org/10.1093/plphys/kiab301>.
- [18] H. Zheng, W. Liu, J. Zheng, Y. Luo, R. Li, H. Wang, H. Qi, Effect of long-term tillage on soil aggregates and aggregate-associated carbon in black soil of northeast China, *PLoS One* 13 (2018) 1–18, <https://doi.org/10.1371/journal.pone.0199523>.
- [19] S. Mondal, D. Chakraborty, Global meta-analysis suggests that no-tillage favourably changes soil structure and porosity, *Geoderma* 405 (2022), 115443, <https://doi.org/10.1016/j.geoderma.2021.115443>.
- [20] A. El Mekkaoui, R. Moussadek, R. Mrabet, A. Douaik, R. El Haddadi, O. Boulhal, M. Elomari, M. Ganoudi, A. Zouahri, S. Chakiri, Effects of tillage systems on the physical properties of soils in a semi-arid region of Morocco, *Agric. For.* 13 (2023) 1–14, <https://doi.org/10.3390/agriculture13030683>.
- [21] J. Zhou, H. Chen, J. Zhou, X. Fu, H. Ye, H.T. Nguyen, Development of an automated phenotyping platform for quantifying soybean dynamic responses to salinity stress in greenhouse environment, *Comput. Electron. Agric.* 151 (2018) 319–330, <https://doi.org/10.1016/j.compag.2018.06.016>.
- [22] J.K. Dhaliwal, M.J. Singh, S. Sharma, N. Gupta, S.S. Kukal, Medium-term impact of tillage and residue retention on soil physical and biological properties in dry-seeded rice-wheat system in north-west India, *Soil Res.* 58 (2020) 468–477, <https://doi.org/10.1071/SR19238>.
- [23] S. Sahoo, P. Mukhopadhyay, J. Mowrer, P.P. Maity, A. Maity, A.K. Sinha, P. Sow, S. Rakesh, Tillage and N-source affect soil fertility, enzymatic activity, and crop yield in a maize-rice rotation system in the Indian Terai zone, *Front. Environ. Sci.* 10 (2022) 1–17, <https://doi.org/10.3389/fenvs.2022.983973>.
- [24] S. Gupta Choudhury, S. Srivastava, R. Singh, S.K. Chaudhari, D.K. Sharma, S. K. Singh, D. Sarkar, Tillage and residue management effects on soil aggregation, organic carbon dynamics and yield attribute in rice-wheat cropping system under reclaimed sodic soil, *Soil Tillage Res.* 136 (2014) 76–83, <https://doi.org/10.1016/j.still.2013.10.001>.
- [25] F.A. Al-Wazzan, S.A. Muhammad, Effects of conservation and conventional tillage on some soil hydraulic properties, *IOP Conf. Ser. Earth Environ. Sci.* 1060 (2022), <https://doi.org/10.1088/1755-1315/1060/1/012002>.
- [26] F. Zulhaedar, M. Nurudin, E. Hanudin, Minimum tillage on vertisols lowland increases rice production and soil properties, *Ilmu Pertanian, Agricultural Sci* 8 (2023) 21, <https://doi.org/10.22146/ipas.75314>.
- [27] A.M. Gajda, E.A. Czyż, A. Klimkowicz-Pawlas, Effects of different tillage intensities on physicochemical and microbial properties of a eutric fluvisol soil, *Agronomy* 11 (2021), <https://doi.org/10.3390/agronomy11081497>.
- [28] K. Azim, B. Soudi, S. Boukhari, C. Perissol, S. Roussos, I. Thami Alami, Composting parameters and compost quality: a literature review, *Org. Agric.* 8 (2018) 141–158, <https://doi.org/10.1007/s13165-017-0180-z>.
- [29] C. Brock, M. Oltmanns, C. Matthes, B. Schmehe, H. Schaaf, D. Burghardt, H. Horst, H. Spieß, Compost as an option for sustainable crop production at low stocking rates in organic farming, *Agronomy* 11 (2021) 1–17, <https://doi.org/10.3390/agronomy11061078>.
- [30] A. Yassi, M. Farid, M.F. Anshori, H. Mughtar, R. Syamsuddin, A. Adnan, The integrated minapadi (Rice-Fish) farming system: compost and local liquid organic fertilizer based on multiple evaluation criteria, *Agronomy* 13 (2023), <https://doi.org/10.3390/agronomy13040978>.
- [31] Z. Zhao, C. Zhang, F. Li, S. Gao, J. Zhang, Effect of compost and inorganic fertilizer on organic carbon and activities of carbon cycle enzymes in aggregates of an intensively cultivated Vertisol, *PLoS One* 15 (2020) 1–16, <https://doi.org/10.1371/journal.pone.0229644>.
- [32] C.A. Phares, S. Akaba, Co-application of compost or inorganic NPK fertilizer with biochar influences soil quality, grain yield and net income of rice, *J. Integr. Agric.* 21 (2022) 3600–3610, <https://doi.org/10.1016/j.jia.2022.07.041>.
- [33] M. Sultana, M. Jahiruddin, M. Rafiqul Islam, M. Mazibur Rahman, M. Anwarul Abedin, Effects of nutrient enriched municipal solid waste compost on soil fertility, crop yield and nutrient content in brinjal, *Eurasian J. Soil Sci.* 10 (2021) 191–198, <https://doi.org/10.18393/ejss.880503>.
- [34] L.P. Luo, B. Han, X.P. Yu, X.Y. Chen, J. Zhou, W. Chen, Y.F. Zhu, X.L. Peng, Q. Zou, S.Y. Li, Anti-metastasis activity of black rice anthocyanins against breast cancer: analyses using an ErbB2 positive breast cancer cell line and tumoral xenograft model, *Asian Pac. J. Cancer Prev. APJCP* 15 (2014) 6219–6225, <https://doi.org/10.7314/APJCP.2014.15.15.6219>.
- [35] T. Razavipour, S.S. Moghaddam, S. Doaei, S.A. Noorhosseini, C.A. Damalas, Azolla (*Azolla filiculoides*) compost improves grain yield of rice (*Oryza sativa* L.) under different irrigation regimes, *Agric. Water Manag.* 209 (2018) 1–10, <https://doi.org/10.1016/j.agwat.2018.05.020>.
- [36] A. Iqbal, L. He, I. Ali, S. Ullah, A. Khan, K. Akhtar, S. Wei, S. Fahad, R. Khan, L. Jiang, Co-incorporation of manure and inorganic fertilizer improves leaf physiological traits, rice production and soil functionality in a paddy field, *Sci. Rep.* 11 (2021) 1–16, <https://doi.org/10.1038/s41598-021-89246-9>.
- [37] M. Anisuzzaman, M.Y. Rafii, N.M. Jaafar, S. Izan Ramlee, M.F. Iqbal, M.A. Haque, Effect of organic and inorganic fertilizer on the growth and yield components of traditional and improved rice (*Oryza sativa* L.) genotypes in Malaysia, *Agronomy* 11 (2021) 1830, <https://doi.org/10.3390/agronomy11091830>.
- [38] M. Momen, M. Bhatta, W. Hussain, H. Yu, G. Morota, Modeling multiple phenotypes in wheat using data-driven genomic exploratory factor analysis and Bayesian network learning, *Plant Direct* 5 (2021) 1–12, <https://doi.org/10.1002/pld3.304>.
- [39] M.F. Anshori, B.S. Purwoko, I.S. Dewi, W.B. Suwarno, S.W. Ardie, Salinity tolerance selection of doubled-haploid rice lines based on selection index and factor analysis, *AIMS Agric. Food.* 7 (2022) 520–535, <https://doi.org/10.3934/agrfood.2022032>.
- [40] A.I. Sakinah, Y. Musa, M. Farid, A. Hairmansis, M.F. Anshori, N. Nasaruddin, Rice selection criteria based on morphological and image-based phenotyping under drought-and salinity-stress conditions, *Sabroa J. Breed. Genet.* 54 (2022) 686–699, <https://doi.org/10.54910/sabroa2022.54.4.1>.
- [41] R. Alsabah, B.S. Purwoko, I.S. Dewi, Y. Wahyu, Selection index for selecting promising doubled haploid lines of black rice, *Sabroa J. Breed. Genet.* 51 (2019) 430–441, <https://sabroajournal.org/wp-content/uploads/2020/01/SABRAO-J-Breed-Genet-514-430-441-Purwoko.pdf>.
- [42] M. Farid, N. Nasaruddin, Y. Musa, M.F. Anshori, I. Ridwan, J. Hendra, G. Subroto, Genetic parameters and multivariate analysis to determine secondary traits in selecting wheat mutant adaptive on tropical lowlands, *Plant Breed. Biotechnol.* 8 (2020) 368–377, <https://doi.org/10.9787/PBB.2020.8.4.368>.
- [43] R. Padjung, M. Farid, Y. Musa, M.F. Anshori, A. Nur, A. Masnenong, Drought-adapted maize line based on morphophysiological selection index, *Biodiversitas* (2021), <https://doi.org/10.13057/biodiv/d220951>.
- [44] A.D. Mochammad, R. Padjung, M. Farid, A.H. Bahrun, M. Fuad Ansho, N. I. Ridwan, A. Nur, M. Taufik, Interaction of genetic and cultivation technology in maize prolific and productivity increase, *Pakistan, J. Biol. Sci.* 24 (2021) 716–723, <https://doi.org/10.3923/pjbs.2021.716.723>.
- [45] Z. Krmáčová, S. Krmáč, M. Barančoková, Analysis of interrelations structure in agro-systems using the factor analysis technique (FA), *Land* 12 (2023), <https://doi.org/10.3390/Land12020272>.
- [46] J.R. do A. Rocha, J.C. Machado, P.C.S. Carneiro, Multitrait index based on factor analysis and ideotype-design: proposal and application on elephant grass breeding for bioenergy, *GCB Bioenergy* 10 (2018) 52–60, <https://doi.org/10.1111/gcbb.12443>.
- [47] W. Taratima, C. Chuanchumkan, P. Maneerattanarungroj, A. Trunjaruen, P. Theerakulpiset, A. Dongsansuk, Effect of heat stress on some physiological and anatomical characteristics of rice (*Oryza sativa* L.) cv. KDML105 callus and seedling, *Biology* 11 (2022) 1–14, <https://doi.org/10.3390/biology11111587>.
- [48] A.A.L. Stansluos, A. Öztürk, G. Niedbala, A. Türkoğlu, K. Hallıoğlu, P. Szulc, A. Omrani, T. Wojciechowski, M. Piekutowska, Genotype-Trait (GT) biplot analysis for yield and quality stability in some sweet corn (*Zea mays* L. Saccharata sturt.) genotypes, *Agronomy* 13 (2023) 1538, <https://doi.org/10.3390/agronomy13061538>.
- [49] A.S. Rahman, A. Rahman, Application of principal component analysis and cluster analysis in regional flood frequency analysis: a case study in new South Wales, Australia, *Water (Switzerland)* 12 (2020) 1–26, <https://doi.org/10.3390/w12030781>.
- [50] B. Sarabi, J. Ghashghaie, Evaluating the physiological and biochemical responses of melon plants to NaCl salinity stress using supervised and unsupervised statistical analysis, *Plant Stress* 4 (2022), 100067, <https://doi.org/10.1016/j.stress.2022.100067>.
- [51] M.F. Anshori, B.S. Purwoko, I.S. Dewi, W.B. Suwarno, S.W. Ardie, Cluster heatmap for detection of good tolerance trait on doubled-haploid rice lines under hydroponic salinity screening, *IOP Conf. Ser. Earth Environ. Sci.* 484 (2020), <https://doi.org/10.1088/1755-1315/484/1/012001>.
- [52] A.G. Yong, S. Pearce, A beginner's guide to factor analysis: focusing on exploratory factor analysis, *Tutor. Quant. Methods Psychol.* 9 (2013) 79–94, <https://doi.org/10.20982/tqmp.09.2.p079>.
- [53] J. Zhang, C. Han, Z. Liu, Absorption spectrum estimating rice chlorophyll concentration: preliminary investigations, *J. Plant Breed Crop Sci.* 1 (2009) 223–229, <http://www.academicjournals.org/jpbs>.
- [54] N.S.H. Kumiawan, I.A.P. Kirana, A.S. Abidin, A. Jupri, S. Widyastuti, A. Hernawan, A. Nikmatullah, H. Sunarpi, E.S. Prasedya, Analysis of leaf chlorophyll content of paddy plants during vegetative stage grown in soil media containing macroalgae organic fertilizer, *IOP Conf. Ser. Earth Environ. Sci.* 913 (2021), <https://doi.org/10.1088/1755-1315/913/1/012025>.
- [55] P.D. Johan, O.H. Ahmed, L. Omar, N.A. Hasbullah, Phosphorus transformation in soils following co-application of charcoal and wood ash, *Agronomy* 11 (2021) 1–25, <https://doi.org/10.3390/agronomy11102010>.
- [56] D. Neina, The role of soil pH in plant nutrition and soil remediation, *Appl. Environ. Soil Sci.* 2019 (2019), <https://doi.org/10.1155/2019/5794869>.
- [57] A. Rahmat, W.S. Ramadhani, D. Prasetyo, S.N. Aini, H. Yanfika, A. Mutolib, The composition of micronutrients and toxic elements in household waste compost uses black soldier fly larvae (*Hermetia illucens*), *IOP Conf. Ser. Earth Environ. Sci.* 739 (2021), <https://doi.org/10.1088/1755-1315/739/1/012004>.
- [58] T. Rodríguez-Espinoza, J. Navarro-Pedreño, I. Gómez Lucas, M.B. Almdendo Candel, A. Pérez Gimeno, A.A. Zorpas, Soluble elements released from organic wastes to increase available nutrients for soil and crops, *Appl. Sci.* 13 (2023), <https://doi.org/10.3390/app13021151>.

- [59] Y. Gao, S. Wang, W. Tan, B. Xi, Dynamics of nutrient elements and potentially toxic elements during composting with different organic wastes, *Front. Sustain. Food Syst.* 7 (2023) 1–10, <https://doi.org/10.3389/fsufs.2023.1181392>.
- [60] A.K. Sadvakasova, B.D. Kossalbayev, A.I. Token, M.O. Bauenova, J. Wang, B. K. Zayadan, H. Balouch, S. Alwaseel, Y.K. Leong, J.S. Chang, S.I. Allakhverdiev, Influence of Mo and Fe on photosynthetic and nitrogenase activities of nitrogen-fixing cyanobacteria under nitrogen starvation, *Cells* 11 (2022) 1–22, <https://doi.org/10.3390/cells11050904>.
- [61] F.S. Saquee, S. Diakite, N.J. Kavhiza, E. Pakina, M. Zargar, The efficacy of micronutrient fertilizers on the yield formulation and quality of wheat grains, *Agronomy* 13 (2023), <https://doi.org/10.3390/agronomy13020566>.
- [62] M. Imran, C. Hu, S. Hussain, M.S. Rana, M. Riaz, J. Afzal, O. Aziz, A.M. Elyamine, M.A. Farag Ismael, X. Sun, Molybdenum-induced effects on photosynthetic efficacy of winter wheat (*Triticum aestivum* L.) under different nitrogen sources are associated with nitrogen assimilation, *Plant Physiol. Biochem.* 141 (2019) 154–163, <https://doi.org/10.1016/j.plaphy.2019.05.024>.
- [63] X. Liu, Y. Shi, L. Kong, L. Tong, H. Cao, H. Zhou, Y. Lv, Long-term application of bio-compost increased soil microbial community diversity and altered its composition and network, *Microorganisms* 10 (2022), <https://doi.org/10.3390/microorganisms10020462>.
- [64] H. Miyamoto, K. Shigeta, W. Suda, Y. Ichihashi, N. Nihei, M. Matsuura, A. Tsuboi, N. Tominaga, M. Aono, M. Sato, S. Taguchi, T. Nakaguma, N. Tsuji, C. Ishii, T. Matsushita, C. Shindo, T. Ito, T. Kato, A. Kurotani, H. Shima, S. Moriya, S. Wada, S. Horiuchi, T. Satoh, K. Mori, T. Nishiuchi, H. Miyamoto, H. Kodama, M. Hattori, H. Ohno, J. Kikuchi, M.Y. Hirai, An agroecological structure model of compost—soil—plant interactions for sustainable organic farming, *ISME Commun* 3 (2023) 1–12, <https://doi.org/10.1038/s43705-023-00233-9>.
- [65] A. Sharma, T.N. Saha, A. Arora, R. Shah, L. Nain, Efficient microorganism compost benefits plant growth and improves soil health in calendula and marigold, *Hortic. Plant J.* 3 (2017) 67–72, <https://doi.org/10.1016/j.hpj.2017.07.003>.
- [66] A. Aguilar-Paredes, G. Valdés, N. Araneda, E. Valdebenito, F. Hansen, M. Nuti, Microbial community in the composting process and its positive impact on the soil biota in sustainable agriculture, *Agronomy* 13 (2023), <https://doi.org/10.3390/agronomy13020542>.
- [67] Komariah, D.D.A. Pitaloka, I. Batubara, W. Nurcholis, A. Sandrawati, A. Setyawati, J. Syamsiyah, W.S. Dewi, The effects of soil temperature from soil mulching and harvest age on phenol, flavonoid and antioxidant contents of Java tea (*Orthosiphon aristatus* B.), *Chem. Biol. Technol. Agric.* 8 (2021) 1–13, <https://doi.org/10.1186/s40538-021-00256-1>.
- [68] A. Pramudia Suciandini, Variability of harvest time of vegetable at different four altitudes based on heat unit analysis, *IOP Conf. Ser. Earth Environ. Sci.* 648 (2021), <https://doi.org/10.1088/1755-1315/648/1/012071>.
- [69] G.V.P.J.T. Parthasarathi, Impact of crop heat units on growth and developmental physiology of future crop production: a Review, *Res. Rev. J. Crop Sci. Technol.* 2 (2013) 11–18, <http://www.stmjournals.com/sci/index.php?journal=RRJoCST&page=article&op=view&path%5B%5D=311%5Cn>.
- [70] K. Ba, V.P. Chimmad, Studies on morpho-phenological traits and heat unit accumulation in chickpea genotypes under different temperature regimes, – 2956, *J. Pharmacogn. Phytochem.* 7 (2018) 2956–2961.
- [71] E.A. Waraich, R. Ahmad, A. Halim, T. Aziz, Alleviation of temperature stress by nutrient management in crop plants: a review, *J. Soil Sci. Plant Nutr.* 12 (2012) 221–244, <https://doi.org/10.4067/S0718-95162012000200003>.
- [72] A. Giri, S. Heckathorn, S. Mishra, C. Krause, Heat stress decreases levels of nutrient-uptake and -assimilation proteins in tomato roots, *Plants* 6 (2017) 443–448, <https://doi.org/10.3390/plants6010006>.
- [73] R.C. Medina Litardo, S.J. García Bendezi, M.D. Carrillo Zenteno, I.B. Pérez-Almeida, L.L. Parismoreno, E.D. Lombeida García, Effect of mineral and organic amendments on rice growth and yield in saline soils, *J. Saudi Soc. Agric. Sci.* 21 (2022) 29–37, <https://doi.org/10.1016/j.jssas.2021.06.015>.
- [74] J. Sung, W. Kim, T.K. Oh, Y.S. So, Nitrogen (N) use efficiency and yield in rice under varying types and rates of N source: chemical fertilizer, livestock manure compost and food waste-livestock manure compost, *Appl. Biol. Chem.* 66 (2023), <https://doi.org/10.1186/s13765-022-00766-y>.
- [75] R. Hindersah, A.M. Kalay, A. Talahaturuson, Rice yield grown in different fertilizer combination and planting methods: case study in Buru Island, Indonesia, *Open Agric* 7 (2022) 871–881, <https://doi.org/10.1515/opag-2022-0148>.
- [76] G.R. Sadimantara, D.N. Yusuf, E. Febrianti, S. Leomo, Muhidin, The performance of agronomic traits, genetic variability, and correlation studies for yield and its components in some red rice (*Oryza sativa*) promising lines, *Biodiversitas* 22 (2021) 3994–4001, <https://doi.org/10.13057/biodiv/d220947>.
- [77] T.A. Urmi, M.M. Rahman, M.M. Islam, M.A. Islam, N.A. Jahan, M.A.B. Mia, S. Akhter, M.H. Siddiqui, H.M. Kalaji, Integrated nutrient management for rice yield, soil fertility, and carbon sequestration, *Plants* 11 (2022) 1–17, <https://doi.org/10.3390/plants11010138>.
- [78] N. Marlina, N. Gofar, A.H.P.K. Subakti, A.M. Rahim, Improvement of rice growth and productivity through balance application of inorganic fertilizer and biofertilizer in inceptisol soil of lowland swamp area, *Agri* 36 (2014) 48–56, <https://doi.org/10.17503/agrivita-2014-36-1-p048-056>.
- [79] M. Siedt, A. Schäffer, K.E.C. Smith, M. Nabel, M. Roß-Nickoll, J.T. van Dongen, Comparing straw, compost, and biochar regarding their suitability as agricultural soil amendments to affect soil structure, nutrient leaching, microbial communities, and the fate of pesticides, *Sci. Total Environ.* 751 (2021), 141607, <https://doi.org/10.1016/j.scitotenv.2020.141607>.

8. Sustainability approach in cropping intensity (CI) 400 through optimizing the dosage of compost and chemical fertilizers to early-maturing rice varieties based on multivariate analysis

ORIGINALITY REPORT

9%

SIMILARITY INDEX

%

INTERNET SOURCES

9%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Mamush Masha, Mengie Belayneh, Elias Bojago, Silas Tadiwos, Amanuel Dessalegn. "Impact of land-use and topography on soil physicochemical properties in the Wamancho watershed, southern Ethiopia", *Journal of Agriculture and Food Research*, 2023
Publication 1%
- 2** A Susila, R Oelviani, W Haryanto, E Winarni, S Jauhari. "Growth and results test of adaptive superior rice varieties through integrated components in the rainfed of Central Java", *IOP Conference Series: Earth and Environmental Science*, 2021
Publication 1%
- 3** Muhammad Fuad Anshori, Bambang Sapta Purwoko, Iswari Saraswati Dewi, Willy Bayuardi Suwarno, Sintho Wahyuning Ardie. "Systematic selection to adaptive doubled haploid rice lines under different 1%

environments of submergence screening methods", Journal of Agriculture and Food Research, 2023

Publication

4

Amir Yassi, Muh Farid, Muhammad Fuad Anshori, Hamdani Muchtar, Rajuddin Syamsuddin, Adnan Adnan. "The Integrated Minapadi (Rice-Fish) Farming System: Compost and Local Liquid Organic Fertilizer Based on Multiple Evaluation Criteria", Agronomy, 2023

Publication

1 %

5

Hyojin Lee, Il-Ho Choi, Kyung-Ran Hwang. "A comprehensive review of machine learning prediction in the production of bio-oil from lignocellulose via pyrolysis", Research Square Platform LLC, 2024

Publication

1 %

6

Mu'minah, Baharuddin, Hazarin Subair, Fahrudin. "Isolation and Screening Bacterial Exopolysaccharide (EPS) from Potato Rhizosphere in Highland and the Potential as a Producer Indole Acetic Acid (IAA)", Procedia Food Science, 2015

Publication

<1 %

7

Muhammad Farid, Nasaruddin ., Yunus Musa, Ifayanti Ridwan, Muhammad Fuad Ansho. "Effective Screening of Tropical Wheat Mutant

<1 %

Lines under Hydroponically Induced Drought Stress Using Multivariate Analysis Approach", Asian Journal of Plant Sciences, 2020

Publication

8

Muh Farid, Muhammad Fuad Anshori, Riccardo Rossi, Feranita Haring et al. "Combining Image-Based Phenotyping and Multivariate Analysis to Estimate Fruit Fresh Weight in Segregation Lines of Lowland Tomatoes", Agronomy, 2024

Publication

<1 %

9

Novaty Eny Dunga, Nurlina Kasim, Muhammad Fuad Anshori, Chalil Gibran Muryadi. "Enhancement of Butternut Squash (*Cucurbita moschata*) Growth and Yield Through Sugarcane Waste and Banana Hump Extract Application", Asian Journal of Plant Sciences, 2023

Publication

<1 %

10

Marcos Denilson Melo Soares, Maila Vieira Dantas, Geovani Soares de Lima, Valeska Karolini Nunes Oliveira et al. "Physiology and yield of 'Gaúcho' melon under brackish water and salicylic acid in hydroponic cultivation", Arid Land Research and Management, 2022

Publication

<1 %

11

Asnelly Ridha Daulay, K.P. Eka Intan, Baba Barus, N. Bambang Pramudya. "Rice Land

<1 %

Conversion into Plantation Crop and Challenges on Sustainable Land Use System in the East Tanjung Jabung Regency", *Procedia - Social and Behavioral Sciences*, 2016

Publication

12

Forita Dyah Arianti, Miranti Dian Pertiwi, Joko Triastono, Heni Purwaningsih et al. "Study of Organic Fertilizers and Rice Varieties on Rice Production and Methane Emissions in Nutrient-Poor Irrigated Rice Fields", *Sustainability*, 2022

Publication

<1 %

13

Adinda Asri Laraswati, Rusnadi Padjung, Muh Farid, Nasaruddin Nasaruddin, Muhammad Fuad Anshori, Amin Nur, Andi Isti Sakinah. "Image Based-Phenotyping and Selection Index Based on Multivariate Analysis for Rice Hydroponic Screening under Drought Stress", *Plant Breeding and Biotechnology*, 2021

Publication

<1 %

14

Azinwi Tamfuh P, Kamga Pangop CR, Douanla Tapindje DG, Boukong A, Tabi FO, Cho-Ngwa F, Bitom D. "Effect of dolomite amendment of acid Andosols on the performance of two green beans varieties in the Cameroon Western Highlands", *International Journal of Advanced Geosciences*, 2019

Publication

<1 %

15

Gusti Ray Sadimantara, Muhidin, Edi Cahyono. "Genetic analysis on some agromorphological characters of hybrid progenies from cultivated paddy rice and local upland rice", *Advanced Studies in Biology*, 2014

Publication

<1 %

16

N. T. Son, C. F. Chen, C. R. Chen, L. Y. Chang, S. H. Chiang. "RICE YIELD ESTIMATION THROUGH ASSIMILATING SATELLITE DATA INTO A CROP SIMULATION MODEL", *ISPRS - International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 2016

Publication

<1 %

17

Md. Abdullah Al Mamun, Sheikh Arafat Islam Nihad, Md. Abdur Rouf Sarkar, Md. Abdullah Aziz et al. "Growth and trend analysis of area, production and yield of rice: A scenario of rice security in Bangladesh", *PLOS ONE*, 2021

Publication

<1 %

18

Muhammad Fuad Anshori, Bambang Sapta Purwoko, Iswari Saraswati Dewi, Sintho Wahyuning Ardie, Willy Bayuardi Suwarno. "A New Approach to Select Doubled Haploid Rice Lines under Salinity Stress Using Indirect Selection Index", *Rice Science*, 2021

Publication

<1 %

19

"Sustainable Solutions for Elemental Deficiency and Excess in Crop Plants", Springer Science and Business Media LLC, 2020

Publication

<1 %

20

Binoy Kumar Show, Suraj Panja, Richik GhoshThakur, Aman Basu et al. "Optimisation of Anaerobic Digestate and Chemical Fertiliser Application to Enhance Rice Yield—A Machine-Learning Approach", Sustainability, 2023

Publication

<1 %

21

Nawal Ababsa, Sonia Boudjabi, Haroun Chenchouni. "Biochar Amendments Changed Soil Properties and Improved Cereal Crop Growth Under Salt Stress", Journal of Soil Science and Plant Nutrition, 2023

Publication

<1 %

22

P E R Prahardini, E Fidiyawati, S S Antarlina, T Sudaryono. "Characteristic of grain and rice quality from five superior varieties support food sustainability", IOP Conference Series: Earth and Environmental Science, 2023

Publication

<1 %

23

Tahmina Akter Urmi, Md. Mizanur Rahman, Md. Moshiul Islam, Md. Ariful Islam et al. "Integrated Nutrient Management for Rice

<1 %

Yield, Soil Fertility, and Carbon Sequestration", Plants, 2022

Publication

24

Wira Hadianto, Bambang Sapta Purwoko, Iswari Saraswati Dewi, Willy Bayuardi Suwarno, Purnama Hidayat, Iskandar Lubis. "Agronomic performance, yield stability and selection of doubled haploid rice lines in advanced yield trials", AIMS Agriculture and Food, 2023

Publication

<1 %

25

Yu-Xin Wang, Xie-Yang Liu, Hui-Hui Di, Xiao-Song He, Yue Sun, Song Xiang, Zhan-Bin Huang. "The mechanism of microbial community succession and microbial co-occurrence network in soil with compost application", Science of The Total Environment, 2024

Publication

<1 %

26

Zehuai Chen, Chunlei Chen, Yingpan Yang, Xingpeng Wang, Hanghai Zhou, Chunfang Zhang. "Rhamnolipids supplement in salinized soils improves cotton growth through ameliorating soil properties and modifying rhizosphere communities", Applied Soil Ecology, 2024

Publication

<1 %

27

"Applied Computational Intelligence and Mathematical Methods", Springer Science and Business Media LLC, 2018

Publication

<1 %

28

"Biology of Composts", Springer Science and Business Media LLC, 2020

Publication

<1 %

29

David E. Johnson, Elizabeth Humphreys.
"Enhancing the productivity and sustainability of cropping systems in the coastal zones of tropical deltas of Asia", Field Crops Research, 2021

Publication

<1 %

30

Muh Farid, Nasaruddin Nasaruddin, Yunus Musa, Muhammad Fuad Anshori, Ifayanti Ridwan, Jekvy Hendra, Gatot Subroto.
"Genetic Parameters and Multivariate Analysis to Determine Secondary Traits in Selecting Wheat Mutant Adaptive on Tropical Lowlands", Plant Breeding and Biotechnology, 2020

Publication

<1 %

31

Muhammad Fuad Anshori, Andi Dirpan, Trias Sitaresmi, Riccardo Rossi et al. "An overview of image-based phenotyping as an adaptive 4.0 technology for studying plant abiotic stress: A bibliometric and literature review", Heliyon, 2023

<1 %

32

Peng Li, Duo Ying, Jing Li, Jianjun Deng, Chunkai Li, Shanyi Tian, Gen Zhao, Choufei Wu, Jiaguo Jiao, Min Jiang, Feng Hu. "Global-scale no-tillage impacts on soil aggregates and associated carbon and nitrogen concentrations in croplands: A meta-analysis", *Science of The Total Environment*, 2023

Publication

<1 %

33

Elena Ambros, Olga Kotsupiy, Evgeniya Karpova, Ulyana Panova, Alexander Chernonosov, Elena Trofimova, Boris Goldenberg. "A Biostimulant Based on Silicon Chelates Enhances Growth and Modulates Physiological Responses of In-Vitro-Derived Strawberry Plants to In Vivo Conditions", *Plants*, 2023

Publication

<1 %

34

Poulomi Nandy, Sanjib Kumar Das, Jagadish Chandra Tarafdar. "Effect of Integrated Nutrient Management and Foliar Spray of Zinc in Nanoform on Rice Crop Nutrition, Productivity and Soil Chemical and Biological Properties in Inceptisols", *Journal of Soil Science and Plant Nutrition*, 2022

Publication

<1 %

35

Yeotae Yun. "Changes in the Growth and Yield of an Extremely Early-Maturing Rice Variety

<1 %

According to Transplanting Density", Agriculture, 2023

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On